



GLOBAL JOURNAL OF MEDICAL RESEARCH: C
MICROBIOLOGY AND PATHOLOGY
Volume 14 Issue 1 Version 1.0 Year 2014
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Determination of the Causative Agents of Bacteremia in Children under 5 Years and their Susceptibility Pattern to the Antibiotics

By Dr. Abhineet Mehrotra & Dr. Shailendra Mishra

Career Institute of Medical Sciences, India

Abstract- Objective: To determine the causative agents of bacteremia in children under 5 years and their susceptibility to the commonly used antibiotics.

Methods: This was a cross-sectional study on children (aged from 1 day to 5 years) admitted to the paediatric ward. The patients included all newborn babies and children admitted with fever and suspected of having sepsis. All the included children were clinically diagnosed for septicemia following strict aseptic precautions and the blood sample was taken. Blood culture were done by standard method.

Results: The overall incidence of bacteremia was 23.1%. The incidence of bacteremia was higher among the children of age group 13-60 month (38.1%) than <1 month (23.4%) and 1-12 month (12.5%). The male (25.8%) children were affected than females (19.6%).

Keywords: bacteremia, incidence, children, bacterial isolates.

GJMR-C Classification : NLMC Code: WC 240



Strictly as per the compliance and regulations of:



© 2014. Dr. Abhineet Mehrotra & Dr. Shailendra Mishra. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Determination of the Causative Agents of Bacteremia in Children under 5 Years and their Susceptibility Pattern to the Antibiotics

Dr. Abhineet Mehrotra ^α & Dr. Shailendra Mishra ^σ

Abstract- Objective: To determine the causative agents of bacteremia in children under 5 years and their susceptibility to the commonly used antibiotics.

Methods: This was a cross-sectional study on children (aged from 1 day to 5 years) admitted to the paediatric ward. The patients included all newborn babies and children admitted with fever and suspected of having sepsis. All the included children were clinically diagnosed for septicemia following strict aseptic precautions and the blood sample was taken. Blood culture were done by standard method.

Results: The overall incidence of bacteremia was 23.1%. The incidence of bacteremia was higher among the children of age group 13-60 month (38.1%) than <1 month (23.4%) and 1-12 month (12.5%). The male (25.8%) children were affected than females (19.6%). The *Klebsiella species* organism was the most common organism which was 44.4% followed by *Staphylococcus Aureus* (14.8%) and *Coagulase Negative staphylococci (CONS)* (11.1%). The percentage of other organism was less than 10%. The percentage of gram negative bacterial isolate was among 70.4% of the samples and gram negative was 29.6%.

Conclusion: The resistance of the recovered *Klebsiella spp.* isolates to a number of antimicrobial agents was determined, and a pattern of multi-resistance was observed which may explain the prevalence of these isolates in pediatric bacteremia.

Keywords: bacteremia, incidence, children, bacterial isolates.

I. INTRODUCTION

Bacteremia is the presence of viable bacteria in the circulating blood. Bacteria may enter the blood stream giving rise to bacteremia from an existing focus of infection from a site with the commensally flora or by direct inoculation of contaminated materials into the vascular system. These organisms are often cleared from the blood within minutes, so the bacteremia is silent and transient, but if the immune system is overwhelmed or evaded, organisms persist in the blood and bacteremic symptoms would arise. Bacteremia should be distinguished from septicemia in which signs and symptoms of severe diseases are present¹. Neonates are particularly vulnerable to infections

because of their weak immune barrier. Several risk factors have been identified both in the neonates and children which makes them susceptible to infections². Children with septicemia present with fever, difficulty in breathing, tachycardia, malaise, refusal of feeds or lethargy³.

Studies of bloodstream infections in children admitted to African hospitals suggest that the prevalence of bacterial bloodstream infections among inpatients with fever or clinical sepsis exceeds that described in wealthier regions^{4,5}. Bloodstream infections continue to be the major cause of morbidity and mortality despite advance in antimicrobial therapy and supportive care⁶. Fever in infants younger than 1 year old, especially those younger than 3 months, can signal a serious infection⁷.

The aim of this study was to determine the causative agents of bacteremia in children under 5 years and their susceptibility to the commonly used antibiotics.

II. MATERIAL AND METHODS

This was a cross-sectional study approved by the ethical committee of the institute. In this study, 117 blood samples were collected from children (aged from 1 day to 5 years) admitted to the paediatric ward of a teaching hospital in north India. The patients included all newborn babies and children admitted with fever and suspected of having sepsis. Children with fever less than 5 days and with known clinical condition such as malignancies, tuberculosis etc. were excluded from the study. The consent was taken from parent/guardian of each children before enrolling in the study.

A total of 128 children were included in this study. All the included children were clinically diagnosed for septicemia following strict aseptic precautions and the blood sample was taken. One milliliter (neonates) and 5 ml (children) blood were collected, inoculated into 10 ml and 50 ml, respectively of brain heart infusion broth. The culture bottles were incubated at 37°C aerobically and periodic subcultures were done onto Mac Conkey's agar, blood agar and chocolate agar after overnight incubation on day three, day four and finally on day seven⁸. The growth obtained was identified by conventional biochemical tests and the antibiotic sensitivity testing was performed on Mueller-Hinton agar plates by Kirby-Bauer disc diffusion method. Zone

Author ^α: Assistant Professor, Department of Microbiology, Career Institute of Medical Sciences, Lucknow, UP, India.

e-mail: abhineetmehrotra@gmail.com

Author ^σ: Junior Resident Department of Pharmacology, Lucknow ERA's Medical College, Lucknow, UP, India.

diameter was measured and interpreted as per the Clinical and Laboratory Standards Institute (CLSI) guidelines as used by Tiwari et al⁸. Bacterial sensitivity was tested for the following antimicrobials: amikacin, amoxicillin clavulanic acid, ampicillin, aztreonam, cefotaxime, ceftazidime, ceftriaxone, cephalexin, ceftoxitin, ciprofloxacin, gentamicin, imipenem, meropenem, piperacillin-tazobactam, tobramycin, linezolid and vancomycin.

Methicillin resistance in *Staphylococcus aureus* (MRSA) was tested using Mueller-Hinton agar with 4% NaCl with ceftoxitin disc (30 micrograms) by Kirby-Bauer disc diffusion method. A zone size of >22 mm was considered sensitive and < 21 was considered as resistant. Suspected extended-spectrum beta lactamases (ESBLs) producing organisms were confirmed by double disk synergy test⁸. Detection of plasmid-mediated AmpC was done by the AmpC disk test and the isolates showing reduced susceptibility to carbapenems (imipenem and meropenem) were selected for detection of metallo-beta lactamases (MBLs) enzymes by imipenem-EDTA disk method. For quality control of disc diffusion tests ATCC control strains of *E. coli* ATCC 25922, *S. aureus* ATCC 25923 and *P. aeruginosa* ATCC 27853 strains were used. The results were expressed as percentages. Microsoft excel was used for the interpretation of these results.

III. RESULTS

Table-1 presents the percentage of children of Bacteremia according to demographic profile. The overall incidence of bacteremia was 23.1%. The incidence of bacteremia was higher among the children of age group 13-60 month (38.1%) than <1 month (23.4%) and 1-12 month (12.5%). The male (25.8%) children were affected than females (19.6%).

The *Klebsiella species* organism was the most common organism which was 44.4% followed by *Staphylococcus aureus* (14.8%) and *Coagulase Negative staphylococci (CONS)* (11.1%). The percentage of other organism was less than 10% (Fig.1).

The percentage of gram negative bacterial isolate was among 70.4% of the samples and gram positive was 29.6% (Fig.2).

Table-2 depicts antibacterial resistance pattern of the gram negative blood stream isolates. *Klebsiella spp* organism was resistance to ampicillin (3), amoxyclav (5), amikacin & gentamycin (2) and one each to cotrimoxazole, ciprofloxacin, cefotaxime & aztreonam. The frequency of other organism resistance to most of drugs was one. The frequency gram positive isolates resistance to most of the drugs was one (Table-3).

IV. DISCUSSION

WHO guidelines for management of acute illness in children (Integrated Management of Childhood

Illness) recommend use of an appropriate antibacterial drug in addition to an antimalarial drug in children with certain signs of severe illness⁹. Despite considerable progress in hygiene, antimicrobial therapy, and supportive treatment, blood stream infections remain important causes of morbidity and mortality which may reach to 20%-30%¹⁰. Microbiologic culturing of blood is the only available means for diagnosis of these infections and allows for successful recovery of bacteria in 99% in patients with bacteremia of septicemia¹¹. An American review covering a 50-years period has shown major changes in the etiology of neonatal septicemia¹².

In this study, the incidence of bacteremia was higher among male children. Similar findings had been reported in southern state of India⁸. Some other studies had also reported higher incidence of bacteremia in male children^{13, 2}. The incidence of bacteremia was higher among 13-60 month (38.1%) in this study which was contradictory to the studies by Tsering et al¹⁴ and Meremkwer et al.² in which incidence of bacteremia was more common among newborns. In this study, the culture positivity rate was found to be 23.1% (23/117). Almost similar rate had been reported in a Indian study in which blood culture positivity rate of 25%⁸. Other studies had also reported similar positivity rate^{15, 14}.

The percentage of gram negative organism found in this study was similar to other studies also^{16, 15}. In the present study, the *Klebsiella species* was the commonest isolate associated with bacteremia which was similar to study by Al-Charrakh et al¹⁷. Many studies have been shown that Gram positive organisms were the mainly *Staphylococcus aureus* as the most frequently isolated bacteria causing bacteremia^{18, 19}.

There was varying number of the gram negative and positive organisms resistance to different drugs in this study. Prabhu et al²⁰ reported that the gram negative organisms showed maximum resistance to ampicillin. However, Tiwari et al⁸ reported that the gram positive organisms showed 77.78% resistance to penicillin but were 100% sensitive to linezolid and vancomycin. Among the 6 *Staphylococcus aureus*, 2(33.33%) were detected as Methicillin resistant *Staphylococcus aureus* (MRSA).

One of the limitations of this study is lesser sample size. Studies on larger sample size is recommended for better interpretation of the results. The studies on the community acquired blood stream infections is also needed at present.

V. CONCLUSION

The resistance of the recovered *Klebsiella spp.* isolates to a number of antimicrobial agents was determined, and a pattern of multiresistance was observed which may explain the prevalence of these isolates in pediatric bacteremia.

Source of funding: None

Conflict of interest: None

REFERENCES RÉFÉRENCES REFERENCIAS

- Campbell A.G.M. and Mclutosh N., Forfar and Arnill's text book of pediatrics, 5th Ed. ELBS Churchill Livingston Educational, low priced books, Scheme funded by British government, 1341-1343 (1998).
- Meremkwere MM, Nwachukwu CE, Asuquo AE, Okebe J, Utsalo SJ. Bacterial isolates from blood cultures of children with suspected septicaemia in Calabar, Nigeria. BMC Infect Dis. 2005; 5:110-5.
- Nwadioha SI, Nwokedi EOP, Kashibu E, Odimayo MS, Okwori EE. A review of bacterial isolates in blood cultures of children with septicaemia in a Nigerian tertiary Hospital. Afr J Microbiol Res. 2010;4:222-5.
- Bahwere P, Levy J, Hennart P, et al. Community-acquired bacteremia among hospitalized children in rural central Africa. Int J Infect Dis. 2001; 5:180-88.
- Idoko JA, Lawande RV, Mohammed I. A prospective study of septicaemia in Zaria, northern Nigeria. East Afr Med J. 1986; 63:515-21.
- Al-Waznee W.S., Study on septicemia in immunocompromised patients, M.Sc. thesis, College of Science, University of Babylon, (2001).
- Torpy J.M., Lynn C. and Glass R.M., Fever in infants, JAMA, 2004; 291, 1284-1286.
- Tiwari Devendra Kumar, Golia Saroj, Sangeetha K.T., Vasudha C.L. A Study on the Bacteriological Profile and Antibigram of Bacteremia in Children Below 10 Years in a Tertiary Care Hospital in Bangalore, India. Journal of Clinical and Diagnostic Research. 2013 Dec, Vol-7(12): 2732-2735.
- Who, Unicef. Integrated management of childhood illness chart booklet—standard. Geneva: World Health Organization; 2008.
- Washington, J.A. and Istrup, D.M.. Blood cultures issues and controversies. Rev. Infect. Dis., 1986; 8: 792-802.
- Wagner, G.E.. Bacteremia and septicemia. In: Kingsbury, D.T., and Wagner. G.E. (eds.) Microbiology. 2nd edition. John Wiley & Sons. Inc., USA. pp. 315-320; 1990.
- Fredman, R.M., Ingram, D.L., Gross, I., Ehrenkranz, R.A., Warshaw, J.B., and Baltimore, R.S., A half century of neonatal sepsis at Yale, 1929 to 1978. Am. J. Dis. Child., 1981; 135: 140-144.
- Baltimore RS, Bogue CW. Infections Disease - Foci of Infection. In: Burg FD, Polin PA, Ingelfinger JR, Gershon AA, Editors. Gellis & Karan's current pediatric therapy. 17th Edition. USA: Elsevier Science; 2002; 39-256.
- Tsering D C, Chanchal L, Pal R, Kar S. Bacteriological Profile of Septicemia and the Risk Factors in Neonates and Infants in Sikkim. J Glob Infect Dis., 2011 Jan- Mar; 3(1): 42-45.
- Sharma M., A. Yadav, N. Goel, U. Chaudary. Microbial profile of septicemia in children. Ind Jr for the practicing doctor. Vol 5, No.4 (2008-09-2008-10).
- Ali Z. Neonatal bacterial septicaemia at the Mount Hope Women's Hospital, Trinidad. Ann Trop Paediatr. 2004; 24(1):41-4.
- Al-Charrakh Alaa H, Al-Muhana, Al-Saadi Zainab. Bacterial Profile of Blood Stream Infections In Children Less Than Three Years Old. J. Babylon Univ. 2005.10(30): 481-85.
- Koneman EW, Allen SD, Janda WM, Schreckember PC, Winn WC. Koneman's Colour Atlas and text book of Diagnostic Microbiology. 6th edition. Newyork: Lippincott; 2006; 97-99.
- Mohammad A. Bacteremia among Jordanian children at Princess Rahmah Hospital: Pathogens and antimicrobial susceptibility patterns. Iran J Microbiol. 2010; 2(1): 22-26.
- Prabhu K, Bhat S, Rao S. Bacteriologic profile and antibiogram of blood culture isolates in a pediatric care unit. J Lab Physicians. 2010; 2:85-8.

Table 1 : Percentage of children of Bacteremia according to demographic profile

	No. of children assessed	Bacteremia	
		No.	%
Age group			
<1 month	64	15	23.4
1-12 months	32	4	12.5
13-60 month	21	8	38.1
Gender			
Male	66	17	25.8
Female	51	10	19.6
Total	117	27	23.1

Table 2 : Antibacterial resistance pattern of the gram negative blood stream isolates

Antibiotics	Klebsiella spp (n=12)		E.coli (n=2)		Pseudomonas aeruginosa (n=2)		Acinetobacter baumannii (n=1)		S.typhi (n=2)		Citrobacter freundii (n=1)	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Ampicillin	3	25.0	1	50.0	NT	-	NT	-	2	100.0	1	100.0
Amoxyclav	5	41.7	0	0.0	NT	-	NT	-	1	50.0	0	0.0

Amikacin	2	16.7	1	50.0	1	50.0	0	0.0	0	0.0	1	100.0
Cotrimoxazole	1	8.3	1	50.0	NT	-	0	0.0	1	50.0	0	0.0
Gentamycin	2	16.7	0	0.0	1	50.0	1	100.0	0	0.0	0	0.0
Tobramycin	NT	-	NT	-	1	50.0	0	0.0	NT	-	NT	-
Ciprofloxacin	1	8.3	0	0.0	2	100.0	1	100.0	1	50.0	1	100.0
Cefotaxime	1	8.3	1	50.0	NT	-	NT	-	1	50.0	0	0.0
Ceftriaxone	2	16.7	0	0.0	NT	-	NT	-	0	0.0	0	0.0
Ceftazidime	NT	-	NT	-	0	0.0	1	100.0	NT	-	NT	-
Aztreonam	1	8.3	0	0.0	0	0.0	0	0.0	NT	-	0	0.0
Imipenem	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

NT-Not Tested

Table 3 : Antibacterial resistance pattern of gram positive blood stream isolates

Antibiotics	Staphylococcus aureus (n=4)		CONS (n=3)	
	No.	%	No.	%
Penicillin	2	50.0	1	33.3
Amoxyclav	1	25.0	1	33.3
Cefoxitin	1	25.0	2	66.7
Erythromycin	2	50.0	1	33.3
Linezolid	1	25.0	0	0.0
Vancomycin	0	0.0	1	33.3




Figure 1 : Distribution of organisms isolated from blood culture




Figure 2 : Prevalence of gram negative and gram positive bacterial isolates





This page is intentionally left blank