

Clinical Significance of Perceived Occupational Stress Influencing Body Mass and Osteopenia: A Pilot Study

Dr. Suresh.D.R¹ and anil kumar²

1

Received: 14 October 2011 Accepted: 5 November 2011 Published: 15 November 2011

Abstract

Introduction: Studies demonstrating perceived occupational stress, obesity osteopenia among industrial workers are very limited. This pilot study was conducted to analyze the inter-relationships, if any, between stress, obesity and osteopenia among industrial workers.

Materials and Methods: All patients who attended the health camps aging from 18 years to 70 years of either sex were the subjects of the study. They were subjected to General Physical Examination along with measurement of anthropometric parameters (Body Mass Index, Waist to hip ratio) Blood pressure and Systemic examination. Perceived Stress Scale (4 point) accepted worldwide for assessment of Stress was given in the form of Clinical Questionnaire to the subjects. Blood samples from the patients were subjected to random blood sugar and lipid profile. Bone Scan was done to assess the Bone Density as a marker of Osteoporosis. Comparison of the parameters obtained were done using appropriate statistical methods. Correlations were done using Pearson's correlation co-efficient. All statistical tests were conducted at 5

Index terms— Perceived stress scale; Osteopenia; Obesity; Bone density; Waist to Hip ratio.

1 INTRODUCTION

Stress refers to a psycho physiological response of a living organism to a perspective challenge, change or threat. Occupational stress (job stress) is a psychosocial disorder which is an impact of the interaction between the worker and his work environment on the worker themselves. If left unidentified it can cause serious physical and physiological illness to the individual, which affects both the individual and the organization. Occupational stress may occur due to stress factors at the individual level, or at the organization level or at the interface of the two. Industrial workers with perceived psychological stress can have impaired job performance and increased risk of stress-related illness such as hypertension, risk factors of coronary heart disease, depression, sleep disorders etc. It is also likely that stress-induced elevations of cortisol may contribute to a tendency to overeat, which in turn contributes to obesity. Obesity itself is associated with disturbances in Hypothalamic-Pituitary-Adrenal axis (HPA) function, leading to cortisol dysregulation. Obesity is associated with a state of chronic low grade inflammation which is associated with various systemic diseases. The relationship between the obesity and bone density in industrial workers with occupational stress has not been well documented. Hence, this pilot study was conducted to observe the relationship, if any between the occupational stress, obesity and bone mass in industrial workers under Employees' State Insurance Scheme. E-mail : buddhatozen@yahoo.co.in

Results: The Body Mass Index & Waist to Hip ratio correlated with osteopenia. The Obese persons had increasing osteopenia along with increased lipid profiles. Overweight persons had slight osteopenia along with normal lipid profile. Persons who were perceived to be under stress had poor control of Blood Sugar inspite of being treated with oral hypoglycaemic agents. Most of the subjects had high scores on perceived stress scale, high incidence of substance abuse, lack of knowledge regarding healthy diet & healthy lifestyle.

2 II.

3 MATERIALS AND METHODS

4 S Author

A cross-sectional study was conducted on all patients who attended the ESICMC & PGIMSR Model Hospital Health Camps (n = 105 out of total 300) aging from 18 years to 70 years of either sex. Patients with chronic diseases, infections, chronic medications, psychiatric disorders, cancers etc were excluded from the study. The data collection was done for around 8 hours so as to cover all those in the day shift. Perceived Stress Scale (4 point) accepted worldwide for assessment of Stress was given in the form of Clinical Questionnaire to the subjects which was formulated and adapted as suitable for the industrial workers. The questionnaires were constructed in English and later translated to the native language (Kannada) with the help of Expert. The researcher and the administration agreed prior that each participant would spend no more than ten minutes on the questionnaire. The time was considered critical as the absence of participant from work would mean production delay. The idea of research was clearly explained to the participants in a 3, 4

General Physical Examination along with measurement of anthropometric parameters including Body Mass Index (BMI), Waist to hip (W/H) ratio, Blood pressure & Systemic examination (cardiovascular, respiratory, gastrointestinal, endocrine & neurological system) were carried out on all subjects. Subjects stood barefoot during all anthropometric assessments. Waist circumference was measured by a tape measure at the midpoint between the upper iliac crest and lower costal margin in the midaxillary line. W/H ratio was correlated with Asian Standards. (Women 88-90, Men 100-102). BMI was categorized into three types: Normal (18.5-24.9), overweight (25-29.9) & Obese (30-39.9). 5,6 Blood samples from the patients were subjected to biochemical & Pathological investigations like blood sugar, lipid profile & Haemogram by the Laboratory Personnel under the supervision of the Faculty.

Dual Energy X-Ray Absorptiometry (DEXA) Bone Scan was done to assess the Bone Density. Bone Density was expressed as T-Score (T-score is a comparison of a person's bone density with that of a healthy 30-year-old of the same sex). A T-score of -2.5 or lower qualifies as osteoporosis. A T-score of -1.0 to -2.5 signifies osteopenia, meaning below-normal bone density without full osteoporosis. 7 Comparison of the parameters obtained were done using ONE WAY ANOVA test. Correlations between the anthropometric measurements, biochemical data, Perceived stress scale & the degree of osteopenia (based on bone density index) were done using Pearson's correlation co-efficient. All statistical tests were conducted at 5% level of significance using SPSS software & online statistical tools.

5 III.

6 RESULTS

Based on the BMI, the subjects were grouped into Normal, Overweight & Obese persons. Perceived stress was more in obese group compared to overweight & normal group. Although osteopenia was increasing with increasing BMI, the increments were insignificant. Mean Arterial Pressure was increased in overweight & obese persons. Overweight persons had slight osteopenia along with normal lipid profile. The Obese persons had increasing osteopenia along with increased lipid profiles. Persons who were perceived to be under stress as per the Perceived Stress Scale Questionnaire had poor control of Blood Sugar with increasing BMI, inspite of being treated with oral hypoglycaemic agents. (TABLE 1 Volume XI Issue IV Version I common gathering. The participants were allowed to interact with the researcher and clarify the concepts explained. Following this the participants who were willing to participate were asked to sign the informed consent forms. The subjects filled the questionnaire with the help of the researcher regarding understanding of questions. The 10-item scale assesses feelings and thoughts during the last month. The PSS measures the degree to which situations in one's life are perceived as stressful. It is more strongly related to life event impact scores as opposed to the number of stressful events, thereby representing one's appraisal of the events as being stressful. The participants were asked to rate their feelings on a scale of 0-4, with 0 indicating never and 4 indicating very often. The PSS scores range from 0 to 40, with higher scores indicating higher levels of stress.

(* -p<0.05 -significant) (* -p<0.05 -significant)

IV.

7 DISCUSSION

According to the WHO report, "Raising Awareness of Stress at Work in Developing Countries" in 2007, occupational stress is one of the most common forms of stress in developing countries as the socioeconomic status, social inequalities and overpopulation forces employees to work based on job availability without a choice. Occupational stress (job stress) is a psychosocial disorder which is an impact of the interaction between the worker and his work environment on the worker themselves. Studies have shown that blue collar workers are highly vulnerable to occupational stress, both in developed and developing In our study, most of the subjects had increased score of perceived Job stress which correlated with increasing body mass. The hypothalamic-pituitary-adrenal (HPA) axis plays a central role in the regulation of energy metabolism through the actions of the

101 glucocorticoids. Stress may contribute to HPA axis dysregulation, tendency to overeat, which in turn contributes
102 to a cascade starting with obesity and ending with type 2 diabetes and CVD. 9 describe the neuroendocrine
103 abnormalities associated with visceral obesity and report a decreased cortisol variability in relation to increased
104 abdominal obesity. 10,11 Although we cannot establish causal pathways from this cross-sectional analysis, it is
105 possible that higher stress levels caused altered HPA axis function, which in turn contributed to increases in
106 BMI. But, the bidirectional nature of the relationship between BMI and stress, genetically mediated adaptations
107 which might cause obesity and perceived stress as a consequence of social stigma associated with obesity cannot
108 be ruled out. A key component of obesity is intra-abdominal accumulation of fat, which is responsible for a
109 great portion of the increased CVD risk associated with obesity. 12 Obese group in our study showed increasing
110 osteopenia with increasing W/H ratio & BMI. In obesity, adipose tissue is infiltrated with an increased amount
111 of macrophages, which are an important source of inflammatory cytokines. Obese humans express higher levels
112 of TNF-a in adipose tissue than do lean individuals. Adipose tissue also produces other proinflammatory factors
113 including interleukin-6 (IL-6) and C-reactive protein (CRP). Obesity has also been implicated in the development
114 or progression of musculoskeletal diseases such as osteoarthritis, a common inflammatory bone disease. Obesity
115 may decrease bone formation (osteoblastogenesis) while increasing adipogenesis because adipocyte and osteoblasts
116 are derived from a common multi-potential mesenchymal stem cell. Obesity may increase bone resorption through
117 upregulating proinflammatory cytokines such as IL-6 and TNF alpha. These proinflammatory cytokines are
118 capable of stimulating osteoclast activity through the regulation of the RANKL/RANK/OPG pathway. Obesity
119 is associated with significant increase in serum leptin and decrease in adiponectin. The action of leptin on bone
120 appears to be complex and both positive and negative effects have been reported. 13,14,15 V.

121 8 LIMITATIONS OF THE STUDY

122 The results from this study might be limited to a localised region of Bangalore and cannot be directly compared
123 with results from other parts of India considering the diversity in culture and other factors within the country.
124 Another key limitation of this study is the number of study subjects. All statistical results are significant but
125 certainly have a wide confidence interval. This clearly indicates a small sample size. Also, we have not measured
126 any laboratory parameters indicative of stress like salivary cortisol, dehydro-epi-androsterone etc due to time
127 constraints. To address these limitations, we are proposing a follow-up study of this sample to determine the
128 prospective relationships among stress, obesity, and HPA axis dysfunction.

129 9 VI.

130 10 CONCLUSION

131 Psychological stress appears to influence the body mass and osteopenia though the causal relationships among
perceived stress, obesity, and HPA function remain to be elucidated. Identification ^{1 2}

)
In Normal & overweight subjects, stress was
correlating negatively with obesity & increasing
osteopenia. In obese group, stress was correlating
201 positively with increasing W/H ratio & osteopenia. Also, W/H ratio correlated
positively with increasing
202 osteopenia in obese persons. Most of these correlations were statistically
insignificant. (TABLE 2) Many subjects who had high scores on perceived
stress scale had high incidence of substance
abuse, lack of knowledge regarding healthy diet &
healthy lifestyle.

Figure 1:

132

¹© 2011 Global Journals Inc. (US)

²December Volume XI Issue IV Version I

1

Dallman et al.

[Note: © 2011 Global Journals Inc. (US) Study found a higher incidence of obesity in people reporting higher levels of job stress. The majority of studies examining stress-HPA axis relationships have either focused on job stress alone or treated body mass index (BMI) as a confounding variable and controlled for it rather than examining its independent relationship to stress and HPA axis function. In a review of the findings from several of their studies, Bjorntorp and Rosmond]

Figure 2: Table 1 :

Figure 3:

-
- 133 [Cohen et al. ()] 'A global measure of perceived stress'. S Cohen , T Kamarck , R Mermelstein . *J Health Soc*
134 *Behav* 1983. 24 (4) p. .
- 135 [Robling et al. ()] 'Biomechanical and molecular regulation of bone remodeling'. A G Robling , A B Castillo , C
136 H Turner . *Annual review of biomedical engineering* 2006. 8 p. .
- 137 [Janssen et al. ()] 'Body mass index and waist circumference independently contribute to the prediction of
138 nonabdominal, abdominal subcutaneous, and visceral fat'. I Janssen , S B Heymsfield , D B Allison , D
139 P Kotler , R Ross . *Am J Clin Nutr* 2002. 75 (4) p. .
- 140 [Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: Executive summary Am .
141 'Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults:
142 Executive summary'. *Am J Clin Nutr* 1998. 68 (4) p. . (Expert panel on the identification, evaluation, and
143 treatment of overweight in adults)
- 144 [Cao et al. ()] 'Expression of RANKL and OPG correlates with age-related bone loss in male C57BL/6 mice'. J
145 Cao , L Venton , T Sakata , B P Halloran . *J Bone Miner Res* 2003. 18 (2) p. .
- 146 [Hamrick et al. ()] *Leptin deficiency produces contrasting phenotypes in bones of the limb and spine*, M W Hamrick
147 , C Pennington , D Newton , D Xie , C Isles . 2004. *Bone*. 34 p. .
- 148 [Wellen and Hotamisligil ()] 'Obesity-induced inflammatory changes in adipose tissue'. K E Wellen , G S
149 Hotamisligil . *The Journal of clinical investigation* 2003. 112 (12) p. .
- 150 [Dallman et al. ()] 'Occupational stress management programmes: a practical overview of published effect
151 studies'. M F Dallman , S E La Fleur , N C Pecoraro , F Gomez , H Houshyar , S F Akana , H
152 References Références Referencias Vander Hek , H N Plomp . *Occup Med (Lond)* 1997. 47 (3) p. . (Minireview:
153 Glucocorticoids-Food intake, abdominal obesity,)
- 154 [Cohen and Williamson (ed.) ()] *Perceived stress in a probability sample of the United States*, S Cohen , G
155 M Williamson . Spacapan S, Oskamp S (ed.) 1998. Beverly Hills, CA: Sage Publications. p. . (The social
156 psychology of health)
- 157 [Mohan et al. ()] 'Prevalence of job strain among Indian foundry shop floor workers'. G M Mohan , S Elangovan
158 , P S Prasad , P R Krishna , A K Mokkaapati . *Work* 2008.
- 159 [Brunner et al. ()] 'Prospective effect of job strain on general and central obesity in the Whitehall II Study'. E
160 J Brunner , T Chandola , M G Marmot . *Am J Epidemiol* 2007. 165 (7) p. .
- 161 [Mcewen ()] 'Protective and damaging effects of stress mediators'. B S Mcewen . *N Engl J Med* 1998. 338 (3) p. .
- 162 [Hsu et al. ()] 'Relation of body composition, fat mass, and serum lipids to osteoporotic fractures and bone
163 mineral density in Chinese men and women'. Y H Hsu , S A Venners , H A Terwedow , Y Feng , T Niu , Z
164 Li . *Am J Clin Nutr* 2006. 83 (1) p. .
- 165 [Primm ()] 'What Workplace Stress Research is Telling Technical Communicators'. D Primm . *Technical*
166 *Communication* 2005. 52 (4) p. .