

# 1 Effect of Cigarette Smoking on Blood Lipids -A Study in 2 Belgaum, Northern Karnataka, India

3 basvaraj devaranavadgi<sup>1</sup>

4 <sup>1</sup> Subbaiah Institute of Medical Sciences, Shimoga, Karnataka, India.

5 Received: 2 June 2012 Accepted: 26 June 2012 Published: 7 July 2012

6

---

## 7 **Abstract**

8 Cigarette smoking is an important and independent risk factor of atherosclerosis, coronary  
9 artery disease and peripheral vascular disorders. Apart from active smokers, passive-smokers  
10 are also prone for the development of smoking related disorders. Smoking adversely affects the  
11 concentration of the plasma lipids and lipoprotein levels. The lipid profile was measured from  
12 100 selected smokers and nonsmokers and the study shows that as the intensity and duration  
13 of smoking increases a significant increase in the levels of very low density lipoprotein  
14 -cholesterol, low density lipoprotein -cholesterol, triglyceride and total cholesterol are noted in  
15 almost all groups of cigarette smokers as compared to nonsmokers. Simultaneously a  
16 significant reduction in the level of High density lipoprotein-cholesterol is observed in cigarette  
17 smokers as the intensity and duration is increased. These findings add another health  
18 enhancing benefit by the cessation of smoking.

19

---

20 **Index terms**— Cigarette smoking, Triglyceride, low density lipoprotein cholesterol, very low density  
21 lipoprotein cholesterol, High Density lipoprotein cholesterol

## 22 **1 Introduction**

23 Lipids play an important role virtually in all aspects of biological life. Some of these roles include serving as  
24 hormones or hormone precursors, helping in digestion, providing energy, storage function and metabolic fuels;  
25 acting as functional and structural compounds in biomembranes and forming insulation to allow nerve conduction  
26 or to prevent heat loss (1).

27 Cigarette smoking is an important and independent risk factor for atherosclerosis, coronary artery disease  
28 and peripheral vascular disorders. (2). There is a dose response relationship between the number of cigarettes  
29 smoked per day and cardiovascular morbidity and mortality (3). Long delay between smoking and onset of  
30 smoking related diseases resulted in the ignorance of ill effects of smoking (4). On an average smoker lose more  
31 than a day of their life span for every week of smoking. Smoking kills more than one in three regular smokers (5).  
32 In India consumers not paid much attention to the tobacco smoking related diseases. (6). Qualities of Indian  
33 cigarettes are far away from western standards (7). India is the 3 rd largest producer and exporter of tobacco in  
34 the world. About 550 million kgs of tobacco is grown in 4.2 lakh hectares of land and 250 million kgs of tobacco  
35 is released for local consumption (8). In India 337 million people above 10 years of age consume tobacco. Every  
36 year 1 million people die prematurely due to tobacco smoking related diseases (9).

37 The mechanism by which smoking increases the cardiovascular diseases are unclear. Recently it has been  
38 suggested that smoking adversely affects the concentration of plasma lipids and lipoprotein levels. However  
39 studies to date have revealed incomplete, inconclusive or conflicting results about the association of smoking  
40 on the plasma lipids and lipoproteins (10). It has been estimated that 1% increase in plasma concentration is  
41 associated with a 2.7% increase in risk (11).

42 As tobacco is grown more in northern Karnataka and also due to paucity of work done in this part the present  
43 study was undertaken. The present study provides a detailed profile of the plasma lipid and lipoprotein levels  
44 depending on duration and intensity of smoking.

## 5 DISCUSSION

---

### 45 2 II.

### 46 3 Materials and Methods

47 The present study composed of 100 selected age and sex matched smokers and non-smokers between the age group  
48 of 20 to 60 years. All the subjects were consuming vegetarian diet and few of them were taking non-vegetarian  
49 diet occasionally, and belonging to different walks of the community. The subjects were volunteer participants  
50 in the study and gave informed consent.

51 All subjects were evaluated and selected by detailed medical history, physical examination, systemic  
52 examination and routine investigations to rule out any underlying diseases. Subjects having diseases, which  
53 are known to influence the blood lipids or patients on lipid lowering drugs or a diet restriction for any reason  
54 and persons chewing tobacco, ex-smokers, obese persons, alcoholics and having risk factors like hypertension,  
55 diabetes mellitus were excluded from the present study. Each patient gave informed consent and the study was  
56 approved by ethical and research committee of J.N. Medical College, Belgaum.

57 The present study comprises of 2 groups Group I-Non smokers (Control) n=25 Group II -Cigarette smokers  
58 n = 75 Cigarette smokers (Group -II) were divided into 3 subgroups depending upon duration and intensity of  
59 smoking. Each group comprises about 25 volunteers Group II A -Mild smokers (n =25) (Duration -1 to 5 years,  
60 smoking 10-15 cigarettes / day) Group II B -Moderate smokers (n=25) (Duration 6 to 10 years, smoking 16-20  
61 cigarettes / day) Group II C -Heavy smokers (n= 25) (Duration -more than 10 years, smoking >20 cigarettes/day)

62 In order to ensure accurate and reproducible results overnight 12-14 hours fasting blood samples were collected  
63 from these subjects.

64 Serum was separated by centrifugation at 3600 rpm for six minutes. The clear serum samples were employed  
65 for the estimation of total cholesterol (12), Triglycerides (13) and HDL-Cholesterol (14). The levels of LDL  
66 cholesterol and VLDL cholesterol were calculated by using Friedewalds formula (15). LDL Cholesterol (mg%) =  
67 Total Cholesterol -(HDL cholesterol + TG/5) VLDL Cholesterol (mg%) = TG/5

68 The significance level of different parameters between the study groups were carried out using students "t"  
69 test.

70 III.

### 71 4 Results

72 The present study was composed of 25 healthy non-smokers as controls and 75 active smokers between the age  
73 group of 20 to 60 years as the test group. Depending upon the duration and intensity of smoking, cigarette  
74 smokers were divided into 3 sub groups of mild, moderate and heavy smokers, as stated above.

75 The results of the present study are given in table-1 and graph-1. Table-1 gives the levels of total cholesterol,  
76 TG, HDL-C, LDL-C and VLDL-C in normal (Group-I), in mild smokers (Group-II A), in moderate smokers  
77 (Group -II B) and in heavy smokers (Group -II C). Graph-I depicts the comparison of the parameters in different  
78 test groups (Group-IIA, II B and IIC).

79 As it is evident from the table-1 and graph-1, the intensity and duration of smoking shows a significant increase  
80 in levels of cholesterol, triglyceride, LDL-C, VLDL-C in almost all the groups of cigarette smokers as compared  
81 to non smokers. Simultaneously a significant reduction in level of HDL-C is observed in cigarette smokers as  
82 compared to non-smokers and a parallel increase in these parameters with the increase in intensity and duration  
83 of smoking. IV.

### 84 5 Discussion

85 Cigarette smokers have a high risk of coronary heart disease than nonsmokers. Several possible explanations  
86 have been offered for this association altered blood coagulation, impaired integrity of the arterial walls, changes  
87 in the blood lipid and lipoprotein concentration.

88 Smoking promotes CHD and atherosclerosis. This may be due to nicotine in cigarette smoke causes an increase  
89 in myocardial oxygen requirement by increasing the use of free fatty acid and also smoking by an unknown  
90 mechanism lowers the antiatherogenic factor HDL-C, remains a significant independent predictor of coronary  
91 artery disease.

92 In our study the mean value of serum total cholesterol in cigarette smokers is significantly higher ( $P < 0.001$ ) as  
93 compared to nonsmokers (refer Table -1). It is observed that cholesterol levels are raised in all groups of cigarette  
94 smokers but the risk is more in heavy smokers.

95 Analyzing the results with regard to the duration of smoking it is observed that on the whole there is a  
96 significant increase in the level of serum cholesterol with regard to an increase in duration and intensity of  
97 cigarette smoking.

98 The mean values of serum LDL-C and VLDL-C are observed to be significantly high ( $P < 0.001$ ) in all groups  
99 of cigarette smokers (refer Table -1). The present study also showed a significant increase ( $P < 0.001$ ) in serum  
100 triglycerides in cigarette smokers as compared to non-smokers (refer Table -1

---

101 **6 .)**

102 The P values obtained with regards to all fractions of serum lipid profile are found to be highly significant in  
103 smokers who smoked more number of cigarettes as compared to nonsmokers.

104 The same characteristics are analyzed with regard to duration of smoking again a significant increase in VLDL-C,  
105 LDL-C, triglyceride and total cholesterol is noted in almost all groups of cigarette smokers as compared to  
106 nonsmokers. On the whole, a significant reduction in the level of HDL-C is observed in cigarette smokers smoking  
107 for longer duration.

108 There are contradicting and varying results regarding total cholesterol, TG, LDL-C and VLDL-C in smokers.  
109 An increase in the total cholesterol, TG, LDL-C and VLDL-C and a significant decrease in HDL-C found in the  
110 present study in smokers as compared to nonsmokers agrees with earlier reports (16)(17)(18)(19)(20). Further  
111 parallel increase is seen in these parameters in mild to heavy smokers (refer table-I). The rise in blood lipid levels  
112 in smokers may be through catecholamine and adenyl cyclase axis induced tissue lipolysis as suggested in chart  
113 -1. Majos O. D. et al. (21) in their study reported that there is significant decrease in HDL-C, but there is  
114 no change in total cholesterol and triglycerides in cigarette smokers as compared to non-smokers. The above  
115 findings, except for decrease in HDL-C are contradictory to our findings.

116 **7 155**

117 Chart -1 : Chart showing a possible mechanism by which nicotine absorbed from cigarette smoke may elevate  
118 plasma lipids and lipoproteins.

119 **8 Cigarette**

120 **9 Conclusion**

121 Our study clearly shows a strong relationship between elevation of serum lipids and cigarette smoking. It also  
122 emphasizes that the changes in the serum lipids tends to be high with the increase in duration and intensity of  
123 smoking. The risk of increase in serum cholesterol with an increase in LDL-C and decrease in HDL-C assume a  
124 great significance since this has been the pattern associated with CHD.

125 The low level of HDL-C in cigarette smokers and the increased exposure of the vascular endothelium to  
126 potentially atherogenic lipoproteins as a consequence of impaired clearance of triglyceride rich lipoproteins may  
provide a mechanism whereby smoking predisposes to greater risk of developing atherosclerotic plaques and CHD.




Figure 1:

127 1 2  
128

---

<sup>1</sup>© 2012 Global Journals Inc. (US) © 2012 Global Journals Inc. (US) 2

<sup>2</sup>© 2012 Global Journals Inc. (US) © 2012 Global Journals Inc. (US)

## 9 CONCLUSION

---

1

intensity of smoking.

| PARAMETERS                 | CIGARETTE SMOKERS                |                           |                          |                         |
|----------------------------|----------------------------------|---------------------------|--------------------------|-------------------------|
|                            | NONSMOKERS<br>GROUP -I<br>(n=25) | GROUP -II A<br>(n=25)     | GROUP -II B<br>(n=25)    | GROUP -II C<br>(n=25)   |
| Total Cholesterol<br>(mg%) | 155.28<br>24.09                  | 196.38 ±18.54<br>P<0.001  | 202.78<br>20.56          | 214.13<br>26.98         |
| Triglycerides<br>(mg%)     | 121.20<br>32.70                  | 167.78 ± 25.41<br>P<0.001 | 171.57± 32.42<br>P<0.001 | 191.16<br>35.45         |
| HDL-Cholesterol<br>(mg%)   | 46.90<br>6.71                    | 38.63 ± 3.06<br>P<0.001   | 35.46 ± 3.50<br>P<0.001  | 31.64 ± 2.56<br>P<0.001 |
| LDL-Cholesterol<br>(mg%)   | 84.08<br>24.42                   | 124.01 ± 18.12<br>P<0.001 | 133.50<br>21.76          | 143.87<br>24.32         |
| VLDL-Cholesterol<br>(mg%)  | 24.24<br>6.54                    | 33.78 ± 4.91<br>P<0.001   | 34.35 ± 6.53<br>P<0.001  | 38.62 ± 7.34<br>P<0.001 |

Values are expressed as Mean ± SD. All P values are in comparison with nonsmokers.

Figure 2: Table 1 :

---

129 [ Global Journals Inc. (US) ()] , *Global Journals Inc. (US)* 2012.

130 [Law et al. ()] 'By how much and how quickly does reduction in serum cholesterol concentration lowers the risk  
131 of ischemic heart disease'. M R Law , S G Wald , Thompson . *BMJ* 1994. 308 p. .

132 [Lopez Virella ()] 'Cholesterol determination in high density lipoproteins separated by three different method'.  
133 Lopez Virella . *Clin Chem* 1977. 23 p. 882.

134 [Muscat and Harris ()] 'Cigarette smoking and plasma cholesterol'. J E Muscat , R E Harris . *Am Heart J* 1991.  
135 121 p. .

136 [Wendy ()] 'Cigarette smoking and serum lipid and lipoprotein concentrations an analysis of published data'. Y  
137 C Wendy . *BMJ* 1989. 298 p. .

138 [Vaidya ()] 'Cigarettes and deforestation'. S G Vaidya . *Tobacco control* 1995. 1 (2) p. .

139 [Tilwani ()] 'Effect of smoking on lipid profile'. R Tilwani . *JAPI* 1997. 45 (7) p. .

140 [Sinha ()] 'Effect of smoking on lipid profile in the young'. A K Sinha . *JAPI* 1995. 43 (3) p. .

141 [Friedewald ()] 'Estimation of the concentration of low-density lipoprotein cholesterol in plasma without use of  
142 the preparative ultracentrifuge'. W T Friedewald . *Clin Chem* 1972. 18 p. .

143 [Pollock ()] 'Forty years on a war to recognize and win.-How the tobacco industry has survived the revelations  
144 on smoking and health'. David Pollock . *British Medical Bulletin* 1996. 52 (1) p. .

145 [Majos ()] 'Lipid effects of smoking'. O D Majos . *Am Heart J* 1988. 115 p. .

146 [Rastogi ()] 'Lipid profile in smokers'. R Rastogi . *JAPI* 1989. 37 (12) p. .

147 [Lipid profile in smokers -A clinical study JIACM ()] 'Lipid profile in smokers -A clinical study'. *JIACM* 2002.  
148 3 (1) p. .

149 [Adedeji and Etukudo ()] 'Lipid profile of cigarette smokers in Calabar municipality'. D A Adedeji , M H Etukudo  
150 . *Pakistan Journal of nutrition* 2006. 5 (3) p. .

151 [Cheryl ()] 'Plasma lipid and lipoprotein profiles of cigarette smokers from randomly selected families, enhancement-  
152 ment of hyperlipidemia and depression of high density lipoprotein'. S B Cheryl . *Am. J. Cardiol* 1983. 52 p.  
153 .

154 [Bucolo and David ()] 'Quantitative determination of serum triglycerides by use of enzymes'. G Bucolo , H David  
155 . *Clin Chem* 1973. 19 p. 475.

156 [Carr and Dreker ()] 'Simplified rapid technique for extraction and determination of serum cholesterol without  
157 saponification'. J J Carr , J J Dreker . *Clin Chem* 1956. 2 p. .

158 [Moxhan ()] 'Smoking'. Johan Moxhan . *Medicine* 1995. 23 (2) p. .

159 [Shyam et al. ()] 'The health economic and social costs of smoking'. C Shyam , P V Prabhakar , V S Singhal .  
160 XXXIX: 93. *Swasth Hind* 1995.

161 [Urmil et al. ()] 'Tobacco and smoking'. A C Urmil , P A Somaliya , V R Gupta . XXXIX: 98. *Swasth Hind* 1995.

162 [Macky and Crofton ()] 'Tobacco and the developing world'. Judith Macky , John Crofton . *British Medical  
163 Bulletin* 1996. 52 (1) p. .

164 [Sanghvi ()] *Tobacco use and cancer in India*" *World smoking and health*, L D Sanghvi . 1980. 5 p. .