

Study Several Biochemical Parameters into Patient's with Hepatitis B Virus

Health and Medical Technical College, Baghdad, Iraq.

Received: 6 February 2013 Accepted: 4 March 2013 Published: 15 March 2013

Abstract

Objective: Hepatitis B virus infections is widely seen all through the world. Disorder in the antioxidant system and oxidative stress may play a role in the pathogenesis of chronic liver diseases. The aim of this study was to estimate the oxidant/antioxidant status in patients with HBV positive group, patients with HBV negative and compared to control group. **Material and Methods:** Thirty six patients with HBV positive group, Thirty six patients with HBV negative and compared to thirty control group were included in this study. Glutathione, superoxide dismutase (SOD) activities, malondialdehyde (MDA), Vit C, Vit E and albumin levels were measured in all patients and control group spectrophotometrically. **Results:** SOD activity, GSH, VitC, Vit E and albumin were significantly lower in patients with HBV positive group when compared to patients with HBV negative and control group. However, MDA levels was increased in each patient group as compared to the control group. **Conclusion:** The current study supposed that deficiency of antioxidant barrier may cause oxidative stress in patients with HBV, and may be antioxidant treatment should be useful for these patients.

Index terms— hbv, antioxidant enzymes, malondialdehyde, vit c and vit e.

1 Introduction

epatitis B is inflammation of the liver, which can be caused by viruses, medications, or toxic agents, the infection is usually characterized by the presence of hepatitis B surface antigen [1]. Other markers are used to determine if the virus is active and replicating when it can cause serious liver damage. During the course of HBV, clearance of hepatitis B early antigen (HBeAg) represents a key event; because it implies that the host is no longer immunotolerant and enters a low replication phase [2]. Age at onset of infection is an important factor affecting the outcome HBV infection. A major world health problem is hyper endemic in South-East Asia and sub-Saharan Africa, being a major cause of morbidity and mortality [3].

Antioxidants are a molecule which can safely interact with free radicals and terminate the chain reaction before vital molecules are damaged. Oxidative damage has been reported to be involved in several hepatic diseases [4].

Antioxidant systems neutralizing the harmful effects of the endogenous Reactive Oxygen Species (ROS) products [5]. Under certain conditions, the oxidative or anti-oxidative balance shifts towards the oxidative status as a result of increase in ROS and/or impairment in antioxidant mechanism [6,7].

Superoxide dismutase can be also considered a member of antioxidants mechanisms of cell since it catalyzes transformation of highly reactive superoxide anion to the less potent hydrogen peroxide. It plays an important role in protection of cells against oxygen toxicity [8].

Glutathione is involved in important cell function including vitamin C metabolism, chelating of copper ions and biotransformation of foreign substances and intermediate oxygen metabolites. GSH is synthesized mainly in liver, it is the main of intracellular defense against free radical and electrophilic xenobiotic of hepatocytes [9].

Vitamin E is generally accepted to be lipid-soluble antioxidant in human. The water-soluble antioxidant; this vitamin is not synthesized in human. It also exhibits a number of important physiological activities that are not related to its antioxidant properties, its function are an electron donor for different enzymes in the cells,

vitamin E was reported to protected hepatocytes against toxic injury [4]. Many studies recoded elevation in the levels of ALT and AST in HBV patients as a result of body's immune response and damage of hepatocytes due to the infection [10].

The aim of the present study is to determine the role of oxidative stress on hepatic damage in patients with hepatitis B virus HBV infection and correlation the MDA, GSH, SOD, Vitamin C, E albumin and uric acid levels with liver function in sera of patients with HBV infection.

2 II.

3 Materialand Methods

The sampling procedure was done in 36 patients (29.53 ± 6.81 years) with HBV positive and 34 patients (31.33 ± 5.52) years with HBV negative. None of these patients received antioxidant medicines or foods. Patients were chosen from the patients referred to the Medical City in Iraq. Patients were compared with 30 healthy control subjects were included (mean age 32.5 ± 5.00). All patients were subjected to a detailed history taking, thorough clinical examination, and laboratory investigations including liver function test, , in addition to lipid peroxidation (the level of lipid peroxidation expressed as malondialdehyde(MDA)), uric acid, Glutathione , vitamin E, vitamin C ,superoxide dismutase [SOD]and albumin had been measured in patients with hepatitis B-virus . Blood samples were obtained from the patients and control group, Five ml were collected from each subject by vein puncture, centrifuged at 3000 rpm for 5 min after allowing the blood to clot at room temperature. The serum GPT, GOT, total serum bilirubin, direct serum bilirubin, Uric acid and Albumin levels were measured by spectrophotometric methods supplied by Giese Diagnostic. Plasma malondialdehyde [MDA] was determined according to the modified method of Satoh [11]. Glutathione was estimated by the method of Beutler's method [12]. Superoxide dismutase was determined according to the method of Misra HP and Fridovich I [13]. Ascorbic acid levels were estimated by the method of Tietz [14]. Vitamin E levels were determined according to a modified of Hashim and Schuttringer [15].

All statistical analyses in studies were performed using SPSS version 17.0 for Windows (Statistical Package for Social Science, Inc., Chicago, IL, USA). Descriptive analysis was used to show the mean and standard deviation of variables. The significance of difference between mean values was estimated by Student-Test. The probability $P < 0.05$ = significant, $P > 0.05$ = non-significant. Correlation analysis was used to test the linear relationship between parameters. ANOVA test was used to show the differences between variables of differentiated groups.

4 III.

5 Results and Discussion

The mean and standard deviation of GPT, GOT, TSB (total, direct and indirect bilirubin) were showed significant increased in their concentration in patients with HBV positive when compared with patient with negative HBV and also with control , as shown in Table 1. Antioxidants play an central role in shielding the body from an oxidative insult by superoxide anion radicals peroxides and hydroxyl radicals and. The sensitive balance between the pro-and anti-oxidant forces in the body appears to be very crucial in determining the state of health, wellbeing and longevity [16]. Hepatitis B is one of the diseases that might cause oxidative stress in the affected subject leading to reduction of the antioxidants of the body, SOD in hepatic diseases may be cause free radical formation [17]. Reduction of antioxidant defense of the liver contribute to the role of oxygen radical formation promote the pathological process in the liver [18]. Several of the ROS which have helpful physiological functions are produced incessantly in the individual organism, other than they may be intimidating for normal cell function and reliability when produced in excess. Consequently, aerobic organisms developed protection mechanisms, such as and SOD against the harmful effects of ROS. Several studies have produced confirmation that a good correlation exists between type and severity of disease and antioxidant level in blood, such as cardiovascular diseases, neurological diseases [19]. Over 90% of GSH inflow in systemic circulation is accounted for by the influx of this peptide from the liver [20].

Reduced blood GSH levels have been reported for patients with liver disease of both alcoholic and nonalcoholic etiology [20,21]. The current study is in agreement with the above results. The primary cause accounting for the decreased blood GSH level in patients with liver diseases is a decreased production in and decreased inflow from the liver [20].

The results in the present study showed that there were statistically significant lower levels of serum SOD, GSH, Vit E, Vit C and albumin among patients with HBV positive group than those of control cases. Since a significant negative correlation between serum GOT, GPT and TSB with SOD, GSH , Vit E, Vit C among the patients, the current study suggest that the association of plasma SOD , Vit E, Vit C and alb with GPT and GOT may improve the biochemical assessment of liver damage.

Detection of the increase of MDA levels which is a product of lipid peroxidation in all patient groups indicates that the oxidative stress is increased in HBV infection. Several study reported elevated MDA levels in patients with chronic hepatitis B and C [22,23]. These findings agree with our findings of a significant elevation of MDA levels in HBV infected patients. Moreover, The result showed a significant positive correlation between serum MDA with GOT, GPT and total bilirubin. One hypothesis has been showed that hydroxyl radicals may react by

either hydroxylation or hydrogen abstraction, setting off free-radical chain reactions that subsequently increases MDA concentration [24]. In summary, the present results agree with other studies that have shown increased MDA level and changes in activities of GSH [25]. These findings indicate that the glutathione antioxidant system is imbalanced in hepatocellular damage, and they support the hypothesis that oxidative stress plays an important role in the development of these liver diseases.

In conclusion, serum MDA, GSH and SOD measurements are useful in monitoring hepatocellular damage in patients with HBV positive. Also the present study considered that deficiency of antioxidant barrier may cause oxidative stress in patients with HBV and, so antioxidant treatment should be useful for these patients.

1

Characteristic	Patient with HBV	Patients Control with HBV
	positive [n=36]	negative [n=30]
GPT [Iu/ml]	48.80±9.31	32.30±3.98
	a,b	a
GOT [Iu/ml]	50.33±8.89	30.80±4.58
	a,b	a
TSB [mg/dl]	2.55±0.25	1.50±0.16
	a,b	a
Direct S.B [mg/dl]	1.52±0.90	0.81±0.07
	a,b	c
Indirect S.B [mg/dl]	1.08±0.08	0.73±0.06
	a,b	c

a p< 0.001 compared to control group
b p<0.001 compared to patients with HBV negative group
c p< 0.01 compared to control group

Table 2 showed mean and standard deviation of serum, MDA, GSH, SOD, vitamin E, vitamin C, Albumin and uric acid, showed significant difference between patients groups [Patients with HBV positive group, Patients with HBV negative group] and control Group. Serum SOD activity, GSH, vitamin E, vitamin C, Albumin and uric acid were significantly decreased in Patients with HBV positive group when compared to Patients with HBV negative group and control group shown in Table 2.

Figure 1: Table 1 :

2

() F

Characteristic Patients with HBV

	positive [n=36]
MDA	6.25±1.32 a,b
GSH	496±121 a,b
SOD	1.10±0.29 a,b
Vit E	0.88±0.25 a,d
Vit C	0.98±0.30 a,d
Alb	2.79±0.14 a,b
Uric acid	5.11±0.13 e,f

a p< 0.001 compared to control group b p<0.01 compared to group2 c p<0.01 compared to control d p<0

Figure 2: Table 2 :

3

Characteristic		GOT		GPT		TSB	
		r	p	r	p	r	p
MDA	0.81		0.01	0.77	0.01	0.69	0.01
GSH	-	0.75	0.01	-	0.80 0.01	-	0.75 0.01
SOD	-0.75		0.01	-0.69	0.01	-0.78	0.01
Vit E	-0.68		0.01	-0.71	0.01	-0.68	0.01
Wit C	-0.72		0.01	-0.77	0.01	-0.75	0.01
Alb	-0.69		0.01	-0.70	0.01	-0.66	0.01
Uric Acid	-0.08		N.S	0.04	N.S	0.06	N.S
Age	0.89		N.S	0.05	N.S	0.04	N.S

Figure 3: Table 3 :

[Cengiz and Fusun] , B Cengiz , F B Fusun .
[Mehmet] , H Mehmet .
[Matiushin et al. ()] 'Activity of blood antioxidant enzymes in chronic liver damage'. B N Matiushin , A S Ioginov , G N Iakimchuk . *Vopr Med Khim* 1996. 41 (4) p. .
[Harold ()] *Antioxidant: new research. 1 st ed Nova science publisher USA*, V Harold . 2006.
[Aruoma ()] 'Characterization of drugs as antioxidant prophylactics. Free Radic'. O Aruoma . *Biol. Med* 1996. 20 p. .
[2. 2. Chang, M.H. ()] 'Chronic hepatitis virus infection in children'. *J. Gastroenterol. Hepatol* 2. 2. Chang, M.H. (ed.) 1999. 13 p. . (Hepatol.)
[El-Ghaffar et al. ()] 'Effect of interferon- γ therapy on oxidative stress in hepatitis C virus infection'. Abd El-Ghaffar , Y Foud , H H Eid , A . *Arab journal of laboratory medicine* 1999. 25 (3) p. .
[Halliwell ()] 'Free radicals, antioxidants, and human disease: curiosity, cause, or consequence?'. B Halliwell . *Lancet* 1994. 344 p. .
[Bianchi et al. ()] 'Glutathione kinetics in normal man and in patients with liver cirrhosis'. G Bianchi , E Bugianesi , M Ronchi , A Fabbri , M Zoli , G Marchesini . *J Hepatol* 1997. 26 p. .
[Halliwell and Gutteridge ()] B Halliwell , J M Gutteridge . *Free Radicals in Biology and Medicine*. 3, 1999. (rd edition)
[Beutler et al. ()] 'Improved method for the determination of blood GSH'. E Beutler , O Duron , B M Kelly . *J. Lab. Clin. Med* 1963. 61 p. .
[Tietz (ed.) ()] *In; Text book of clinical chemistry*, N W Tietz . N W Tietz, W B Saunders company (ed.) 1986. Philadelphia, London, Toronto. p. .
[Hakim and Ozcan ()] 'Increased oxidative stress associated with the severity of the liver disease in various forms of hepatitis B virus infection'. C Hakim , E Ozcan . *Infectious Diseases* 2005. 5 p. 95.
[Yagi ()] 'Lipid peroxides in hepatic, gastrointestinal and pancreatic diseases'. K Yagi . *Adv-Exp-Med-Biol* 1994. 366 p. .
[Aziz et al. ()] 'Metabolic Investigation in Hepatitis B Patients with or Without Diabetes mellitus to Determine the Impact of Anti-Oxidant Activity WorlAppli Sci'. Abdul Aziz , M Seemab , S Ahmeds;Bikhar;Devrajani , G; A; Memon , Q Ali , S Waqas . *J* 2011. 15 (6) p. .
[Fattovich ()] 'Natural history of hepatitis B Brith'. G Fattovich . *J. Hpatol* 2003. 39 p. .
[Koruk ()] 'Oxidative stress and enzymatic antioxidant status in patients with nonalcoholic steatohepatitis'. M Koruk . *Annals of clinical and laboratory science* 2004. 34 (1) p. .
[P?nar et al. ()] *Oxidative Stress in Patients with Chronic Hepatitis B and C*, C P?nar , K Ergul , K Omer , A Murat . 2011. *Balkan Med J*. 28 p. .
[Sen ()] 'Oxygen toxicity and antioxidants: State of the Art IndianPhysiol'. C K Sen . *Pharmacol* 1995. 39 (3) p. .
[Weinbum and Lyerla (ed.) ()] *Prevalence and control of infection with Hepatitis viruses*, Weinbum , R Lyerla . Margolid H.S (ed.) 2003. UK PP 5: Ed Academic Press.
[Hashim and Schuttringer ()] 'Rapid determination of tocopherol in Marco-and micro quantities of plasma, results obtained in various nutrition and metabolic studies'. S A Hashim , G R Schuttringer . *Am. J. Clin. Nutr* 1966. 19 (2) p. .
[Oewi?tek and Juszczuk ()] 'Reduced glutathione concentration in erythrocytes of patients with acute and chronic viral hepatitis'. K Oewi?tek , J Juszczuk . *J Viral Hepatitis* 1997. 4 p. .
[Satoh ()] 'Serum lipid peroxide in cerebrovasculardisorders determined by new colorimetric method'. K Satoh . *Clin.Chim. Acta* 1978. 90 p. .
[Stefan et al. ()] 'The effects of super oxide dismutase on H ₂ O ₂ formation. Free Radic'. I Stefan , S I Liochev , I Fridovich . *Biol. Med* 2007. 42 p. .
[Lebensztejn ()] 'The role of free oxygen radicals in children with chronic viral hepatitis'. D M Lebensztejn . *B. RocznikiAkademiiMedycznej w Bialymstoku* 1995. 40 (3) p. .
[Misra and Fridovich ()] 'The role of superoxide anion in the auto oxidation of epinephrine and a simple assay for superoxide dismutase'. H P Misra , I Fridovich . *J BiolChem* 1972. 247 p. .
[Hassan ()] 'Time course of antioxidant enzyme activities in liver transplant recipients'. L Hassan . *Transplantation proceedings* 2005. 37 (9) p. .