

Chondromyxoid Fibroma of the Lateral Malleolus -En-Block Excision and Steinmann Pin Stabilisation of the Ankle

C. Fletcher

Received: 6 December 2017 Accepted: 5 January 2018 Published: 15 January 2018

Abstract

We report a case of a 25 year old male with a chondromyxoid fibroma located in the lateral malleolus. This was therefore a case of a rare tumour in an unusual location. The tumour was treated via an en-block excision and the ankle was stabilised with a Steinmann pin. The original plan was that the Steinmann pin was to provide temporary stabilisation while awaiting definitive histology. Ankle arthrodesis was supposed to be performed at a later date if the histology confirmed a benign tumour. Delayed wound healing resulted in the Steinmann pin being left in situ for 12 weeks. Two years post Steinmann pin removal; there was no clinical or radiological evidence of tumour recurrence or ankle instability. The patient was satisfied with his function and a secondary procedure was deemed unnecessary. The surgical options for benign distal fibular tumours including chondromyxoid fibroma will be discussed.

Index terms— chondromyxoid fibroma, lateral malleolus, excision, reconstruction.

1 I. Introduction

Chondromyxoid fibroma is a rare, slow-growing tumour accounting for less than 1% of all primary bone tumours. It tends to undergo regional

2 II. Case Presentation

A 25-year-old unemployed male presented with complaints of a hard, painful mass on the lateral right ankle, which was gradually increasing in size over one weight bearing. The intensity of the pain was progressively worsening over the previous four weeks, and this was associated with mild limping. There was no history of trauma.

Core needle biopsy suggested a chondromyxoid fibroma. Preoperatively a staged procedure was planned because of the following factors: 1) The rarity of the diagnosis of chondromyxoid fibroma of the lateral malleolus, 2) The soft tissue invasion and 3) Chondromyxoid fibroma may be confused with malignancies histologically. The plan was for definitive histology to be obtained prior to performing a definitive procedure. Stage one consisted of an enblock excision and Steinmann pin stabilisation of the ankle. Stage two would have been definitive ankle arthrodesis using large fragment cancellous screws, if the post excision histology confirmed a benign tumour. Via a lateral approach (see fig. 3), the fibula was resected 3 cm proximal to the gross tumour (see fig. 4). The distal resection was just distal to the fibular tip. The lateral ligamentous complex was not readily discernible due to scar tissue. There was cortical irregularity of the tibia (see fig. 5) adjacent to the tumour. It was uncertain as to whether or not this was due to a simple mass effect of the distal fibular tumour. Grossly, a complete en-block resection was achieved (see fig. ??). The histological diagnosis remained unchanged post excision. Initial postoperative radiographs demonstrated a reduced ankle as well as complete tumour resection (see fig. ?? and fig. 8). The wound took twelve weeks to heal and the Steinmann pin was left in situ over that duration. At the two year follow up assessment; his ankle and subtalar joints remained stiff however he did not have any complaints in terms of functional limitation. C year. The pain was worse at night and after prolonged enlargement towards local tissues. According to the world health organization; it is a benign tumour with areas which are lobulated and has zones which are round or fusiform separated by abundant intercellular myxoid, chondroid and fibrinoid

3 III. DISCUSSION

44 material as well as varying forms of multinucleated cells (1). The fibrotic areas are as a result of repair of
45 degenerated areas and the myxomatous zones are as a result of necrosis of the chondroid tissue (1). The tumour
46 is more common in males and it is most commonly located in the proximal tibia, distal femur, pelvis and foot
47 (2). Histologically, it is not uncommon to be confused with more aggressive tumours. Chondromyxoid fibromas
48 when treated by excision do not tend to recur and there are no reports of metastasis (2).

49 On examination, there was a 9 by 5 cm hard mass with hyper pigmented overlying skin (see fig. 1). The mass
50 was nonpulsatile, nontender, non mobile, and was not fluctuant or warm. There were no ulcers, no evidence of
51 surrounding oedema and no neurovascular deficits. The range of motion of the ankle and subtalar joints was
52 limited.

53 Radiographs revealed an oval, expansile, lobulated cystic lesion. There was obvious soft tissue invasion laterally
54 and medially, there was a thin peripheral sclerotic rim (see fig. 2).

55 His ankle remained stable and pain free. There was no clinical or radiological evidence of tumour recurrence.

56 3 III. Discussion

57 Chondromyxoid fibroma (CMF) is a benign, locally aggressive bone tumour which has lobulated areas of spindle-
58 shaped or stellate cells with significant amounts of myxoid or chondroid intercellular material. A literature review
59 of distal fibular tumours only consists of case reports and case series regardless of the diagnosis (3). Sharma
60 et al. reported on the 40-year experience of CMF in the foot and ankle area from a bone registry. There
61 were 10 cases with a mean patient age of 19 years and none of those tumours were located in the fibula (4).
62 Persian et al. reported on all the tumours in the distal fibula in their hospital and none of the tumours were
63 CMF. The differential diagnosis seen in that report included angioma, fibroma, lung adenocarcinoma metastasis,
64 aneurismal bone cyst, schwannoma, osteochondroma and multifocal mesenchymal neoplasia. Giant Cell Tumour
65 and chondrosarcoma (??) are other differentials.

66 Due to the rarity of distal fibular tumours, and the contribution of the distal fibula in the biomechanics of the
67 ankle, the treatment remains controversial (3). Post resection of the distal fibula, numerous restraints are lost and
68 this has formed the basis for ankle reconstruction to avoid ankle instability (3). The ankle is a complex formed
69 by the lower fibular end, distal tibia, and talus along with complexed ligamentous supports. It is surrounded by
70 a thin capsule which is weak anteriorly and posterior, however there are numerous static and dynamic stabilisers
71 laterally and medially. The lateral malleolar tip is distal and posterior to the medial malleolar tip which limits
72 eversion while simultaneously acting as a lateral buttress (6).

73 When the core needle biopsy suggested a CMF, it was recognised that this rare diagnosis may be confused with
74 a more common malignant diagnosis such as a chondrosarcoma. A malignant diagnosis was also a consideration
75 because of the cortical breach of Berenstein-Weyel et al. found only 3 cases of CMF in the lateral malleolus in
76 their literature review. The first case was a 28-year-old patient who had intralesional curettage, cementation and
77 intramedullary Kirshner wiring (8). The second case was a 25-year-old female with a recurrent CMF. The initial
78 procedure was curettage without grafting. This was revised two years later with an en-block resection of the
79 distal 12cm of the fibula and allograft reconstruction (9). The allograft was fixed onto the native fibula with a
80 plate and screw construct along with a syndesmotic screw. The third case was a 27-year-old female whose tumour
81 did not have a cortical breach and was therefore amenable to an intralesional excision followed by phenolisation,
82 cementation and Steinmann pin stabilisation in the fibula (6). The index case however required en-block excision
83 because unlike the first and third cases above, there was lateral cortical breach with soft tissue invasion.

84 Berenstein-Weyel et al reported on a case of CMF which was successfully solely treated with radiofrequency
85 ablation in an 11-year-old patient with a 4cm lesion. Regardless of the specific diagnosis, surgical treatment
86 usually has good outcome in all benign cases involving the distal fibula (3).

87 The more commonly used reconstructive options post resection of the distal fibula in the adult population
88 includes: 1) Ipsilateral fibular transplantation (??) 2) Non-anatomic soft tissue reconstruction using peroneus
89 brevis (10) 3) Osteoarticular allograft (11) 4) Ankle arthrodesis (3)

90 Nadkarni et al. reported on reversing the proximal fibula 180° and then incorporated the head of fibula into
91 the ankle mortise. The reversed portion was fixed to the remaining fibula using a plate and screw construct
92 plus a syndesmotic screw. The lateral followed by a protocol involving non weight bearing for three months and
93 full weight bearing at six months. This was part of the reason why the Steinmann pin was retained for twelve
94 weeks in addition to the fact that the wound took the same duration to completely heal. Kumar et al. described
95 transecting the peroneus brevis tendon at the myotendinous junction then suturing this to the anterior talofibular
96 and calcaneofibular ligaments before tenodesis was done to the distal tibia. This was followed by six weeks cast
97 immobilisation. Aside from the delayed wound healing, this was not contemplated as a secondary procedure as
98 the lateral ligamentous complex was compromised due to the presence of poor quality scar tissue. Soft tissue
99 reconstruction has been found to be less stable than bony procedures (5), however it avoids the reduced function
100 as well as non-union which are both potentially associated with arthrodesis (10). The reduced function from
101 ankle arthrodesis is due to an alteration of the gait pattern while completely restricting the range of motion.
102 Soft tissue reconstruction also avoids the non-union risk associated with autologous grafting or allograft, post
103 excision. Additionally, there is also a lower infection risk due to a lack of implants involved in the procedure (4).

104 Jamshidi et al. reported union in all four cases where allograft was fixed to the native bone with a plate
105 the fibula and the irregularity of the tibial cortex. The stabilisation with a Steinmann pin was meant to be

106 temporary, until the definitive diagnosis was made post resection. The delayed wound healing prevented the
107 surgical team from undergoing a secondary procedure. The Steinmann pin was thus unintentionally definitive
108 fixation. ligamentous complex was also sutured. This was and screw construct with a separate syndesmotic
109 screw. Average union time was twelve months. One case had an asymptomatic breakage of the syndesmotic
110 screw and another had a 15 degree late valgus deformity. The patients all had favourable ankle scores including
111 favourable ankle range of motion (11). In our setting, there is no local bone bank and the cost of procuring the
112 allograft would have been prohibitive.

113 We believe that this procedure may be considered in any patient with a medical condition which predisposes
114 to poor wound healing or if a patient has any significant risk factor which impairs wound healing. Another
115 potential group of patients to consider are low demand patients in whom reduced range of motion of the
116 ankle will not significantly impair their ability to perform their activities of daily living. Patients who are
117 in an environment with limited resources including implants and equipment may also benefit. The procedure is
118 technically straight forward and reproducible with minimal cost. Our unemployed patient was unable to afford
119 any implants hence fixation options were significantly limited. The period of non weight bearing is similar to
120 the much more technically demanding procedures. There is an avoidance of a nonunion risk which may be a
121 complication of both arthrodesis, and plate and screw fixation of the bone graft (whether allograft or autogenous
122 grafting) and the native bone. There is also the avoidance of dissection in the common personal nerve territory
123 (5) or an incongruent fibular head articulation after the 180°^r eversal procedure utilised by Nadkarni.

124 Currently, the procedure of choice for distal fibular benign tumours is determined by the surgeon's experience
125 and preference, as opposed to the nature and tumour extent, which reflects the lack of guidelines and high-quality
126 evidence in the current literature (3).



Figure 1: Figure 1 :



2

Figure 2: Figure 2 :

3 III. DISCUSSION



3

Figure 3: Figure 3 :



4


Figure 4: Figure 4 :

3 III. DISCUSSION



5

Figure 5: Figure 5 :



67

Figure 6: Figure 6 :Figure 7 :



3 III. DISCUSSION

127 .1 Declaration of interest

128 The author has no conflict of interest relevant to this article.

129 .2 Funding

130 This research did not receive any specific grant from funding agencies in the public, commercial, or notfor-profit
131 sectors, and no material support of any kind was received.

132 [Kumar et al. ()] , A Kumar , M J Naidu , B Mandela . 2015.

133 [Aneurismal Bone Cyst of Distal Fibula Treated With En-Bloc Excision and Ligament us Ankle Stabilisation: A Rare Case Repo
134 *Aneurismal Bone Cyst of Distal Fibula Treated With En-Bloc Excision and Ligament us Ankle Stabilisation:*
135 *A Rare Case Report*, 14 p. .

136 [Gateless et al. ()] 'Benign bone tumours' S Gateless , R Wilkins , E U Conrad 2 Nd . *Inkster Course Lect* 1996.
137 45 p. .

138 [Berenstein-Weyel et al. ()] 'Chondromyxoid fibro of the distal fibula treated by percutaneous radiofrequency
139 ablation' J T Berenstein-Weyel , E Label , D Katz , Y Applaud , A Geyser . *Journal of Orthopaedic Surgery*
140 2017. 25 (2) p. .

141 [Sharma et al. ()] 'Chondromyxoid fibro of the foot and ankle: 40 years' Scottish bone tumour registry
142 experience' I H Sharma , M J Jane , R Reid . *International Orthopaedics (SICOT)* 2006. 30 p. .

143 [Giorgio et al. ()] 'Chondromyxoid fibro of the lateral alveolus: a case report' Di Giorgio , L Touloupakis , G
144 Mastantuono , M Vitaly , F Imperator , L . *Journal of Orthopaedic Surgery* 2011. 19 (2) p. .

145 [Fotiadis et al. ()] 'Chondromyxoid fibro: a rare tumour with an unusual location' E Fotiadis , P Haritopoulos
146 , E Samoladas . *Arch Northup Trauma Surg* 2008. 128 p. .

147 [Lersundi et al. ()] 'Chondromyxoid fibro: a rarely encountered and puzzling tumour' A Lersundi , H J Mankin
148 , A Morikis , F J Cornice . *Clan Northup Relate Res* 2005. 439 p. .

149 [Persian et al. ()] 'Clinical management and surgical treatment of distal fibular tumours: a case series and review
150 of the literature' C Persian , E Marlette , M S Spinally , C Grace , C Fabricant , N Manfully , G Macabre .
151 *International Orthopaedics (SICOT)* 2012. 36 p. .

152 [Sirveaux et al. ()] 'Distal fibula reconstruction using a frozen allograft: a case report' F Sirveaux , O Roche ,
153 P Hutten . *Rev Chirr Northup Reparative Appear Mot* 2004. 90 p. .

154 [Nadkarni et al. ()] 'Giant Cell Tumour of distal fibula managed by en block resection and reconstruction with
155 ipsilateral proximal fibula' S Nadkarni , A S Pundit , R V Nair . *Journal of Orthopaedic Case Reports* 2015.
156 5 (1) p. .

157 [Jamshidi et al. ()] 'Reconstruction of distal fibula with osteoarticular allograft after tumour resection' K
158 Jamshidi , F N Masher , Z Mascara . *Foot and Ankle Surgery* 2013. (19) p. .

159 [References Références Referencias] *References Références Referencias*,