

# 1 Analysis of RBC Antibody Screening in a Hospital Population 2 Over a Two Years Period

3 Dr. Molly Joseph

4 Received: 14 December 2017 Accepted: 31 December 2017 Published: 15 January 2018

5

---

## 6 **Abstract**

7 RBC antibody screening plays an essential role in the pre-transfusion testing of the blood or  
8 blood products before blood transfusion in the recipients as well as in antenatal screening to  
9 prevent Rh-incompatibility. The antibody screening tests performed in a clinical  
10 laboratory/blood bank are designed to detect the presence of these unexpected antibodies  
11 especially all antibodies in the serum. Methods routinely used for detection of these  
12 antibodies are the Coomb's test (antihuman globulin test), both direct and indirect types.

13

---

14 **Index terms—**

## 15 **1 I. Material & Methods**

16 total of 378 patients and 2050 donors were included in this study period of two years (July 2016 to June 2018) in  
17 the Department of Pathology, St. Stephens Hospital, New Delhi, India. All the hospital population (patients as  
18 well as donors) blood samples were included in the study. The antibody screening tests performed were Indirect  
19 Coomb's tests, Direct Coomb's tests and Auto control. The method of screening used was gel card technology.

## 20 **2 II. Statistical Analysis**

21 Qualitative variables are expressed as frequencies / percentages and compared between groups using Chi-square  
22 / Fisher's Exact Test. Quantitative variables are written regarding mean  $\pm$  sd and compared using Unpaired  
23 t-test / Mann-Whitney Test. A p-value  $< 0.05$  is considered statistically significant. The data is tabulated in  
24 MS Excel and analysis performed using Statistical Package for Social Sciences (SPSS) version 16.0 software.

## 25 **3 a) Study Design**

## 26 **4 Cross-sectional study b) Sample Size Determination**

27 The formula used for sample size estimation was  $d^2 n = Z^2 P(1-P)$

## 28 **5 III. Result and Discussion**

29 Our study included 378 patients and 2050 donors. The age of patients ranged from new born to 80 years with  
30 a mean age of 28.64 years. The maximum Author ? ? ? ? ? DNB, Junior Resident and Department of  
31 Pathology, St. Stephen's Hospital, Delhi. e-mails: jagritiyadav91@gmail.com, richa\_jindal2003@yahoo.com,  
32 hemagoyal88@gmail.com, kuldeep\_0787@yahoo.com, mollymaria54@yahoo.co.in number of cases were in the age  
33 group of 21-30 years (58.20 %), followed by 31-40 years (19.84 %) and two cases were in the age group of 71-  
34 80 years (0.53 %). Female predominance was seen with a male to female ratio of 1:5.6. The most common  
35 causative factors in our study were the previous history of transfusion, females presented with pregnancy either  
36 primigravida or multigravida and Rh-negative blood group. Blood group B was the most frequent blood group  
37 followed by blood group O. The present study has an alloimmunization rate of 3.44% which is comparable to  
38 all the other studies mentioned above. A study by Nikam et al. 7 had least alloimmunization rate of 0.74  
39 %. Choudhary et al. had the highest alloimmunization rate of 9.8%. 3 The percentage of alloimmunization  
40 in all the abovementioned studies fall somewhere between <1% to 10%. According to published data rates

## 7 IV. CONCLUSION

---

41 of alloimmunization in random patients vary from 0 to 3 observational studies in random patients, who most  
42 often receive an incidental transfusion, and pregnant women estimated the prevalence between <1-3%. 12 This  
43 incidence increases in multi-transfused patients and transfusion-dependent patients. The reported prevalence of  
44 alloimmunization in multi-transfused patients in India is comparatively low varying from approximately 3% to  
45 10%. 3,9,13

### 46 6 a) Analysis of Autocontrol in Patients

47 Out of 378 cases, two patients showed auto control positivity.

48 Out of 378 patients, AC was positive only in two patients, and those were females (0.53%). Among those two  
49 patients, one was Rh-positive and the another one was Rh-negative. History of previous transfusion (1.7%) is  
50 the most common cause of auto antibodies followed by pregnancy (0.43%). Similarly, in our study, one patient  
51 had history of transfusion and other was a pregnant female had auto control positive rate. It was observed in  
52 our study that antibodies are more commonly found in the age group between 20-40 years.

53 There is a paucity of literature on the detection of antibodies by auto control. In our study, only ten donors  
54 had antibodies in their blood (0.49%). Makroo RN et al. drawn similar results in his study in which out of 82153  
55 donors 227(0.27%) had antibodies in their blood. 17 The study by Pahuja S et al. had a total of 7756 donors out  
56 of which 4(0.05%) donors had antibodies in their blood. 16 Garg N et al. had a similar result of 0.09% antibodies  
57 in 47450 donors. 18 The blood with antibodies were discarded and not transfused to patients.

## 58 7 IV. Conclusion

59 Clinically significant antibodies were frequently detected in our patients and donors population. Alloimmunization  
60 in Rh D positive women was low as compared to Rh D negative women. The previous history of transfusion  
61 was an important cause for the development of antibodies. Males were more than females in donor population  
62 showed that males are more active in donating the blood.

63 We recommended that ? Antibody screening must be done both in patients and donors to find the irregular  
64 antibodies. ? Antibody screening should be done in pregnant females to prevent Rh incompatibility or HDFN.  
65 ? Antibody screening should be done in donors to detect the presence of alloantibodies and is an important  
66 to provide compatible blood products and to avoid transfusion reactions. ? Multi-transfused patients have a  
67 high probability of developing alloantibodies, so extended screening is recommended in the patients to prevent  
hemolytic transfusion reactions. <sup>1</sup>

1

Total Patients n = 378 Gender Distribution Gender n Male 57 Female 321 Total 378 Age Group (Years) Age

? 10 11 -20	29 16
21 -30 31 -40 41 -50 51 -60 61 -70 71 -80	220
Total	75 14
	13 9
	2 378

Mean ± sd ABO Blood Group 28.64 Blood Group n A 88 B 138 AB 42

O	110
Total	378
Rh D Distribution	
Rh D Positive	352
Rh D Negative	26

[Note: In our study ICT was positive in patients with mean age of 30.24 years, whereas DCT was positive in patients with mean age of 37.04 years and AC was positive with a mean age of 30.59 years. A]

Figure 1: Table 1 :

68

---

<sup>1</sup>Analysis of RBC Antibody Screening in a Hospital Population Over a Two Years Period

---

2

Author	Year	Total Patients	All Immunised Patients (ICT Positive Cases)	The Rate of Alloimmunization (%)
Sirchia et al. 1	1985	1432	74	5.2
Chow et al. 2	1994	436	26	6
Choudhary et al. 3	1999	81	8	9.8
Ansari et al. 4	2007	80	3	3.75
Roopam et al. 5	2009	96	5	5.21
Pahuja et al. 6	2010	211	8	3.79
Nikam et al. 7	2011	74	1	0.74
Usman et al. 8	2011	800	30	3.75
Sood et al. 9	2013	306	13	4.24
Makroo RN et al. 10	2014	49,077	403	0.82
Present study	2018	378	13	3.44

Figure 2: Table 2 :

3

Author	Year	Total Patients	All Immunised Patients (DCT Positive Cases)	The Rate of Alloimmunization (%)
Nakamura Y et al. 14	1984	421	14	3.3
Pahuja S et al. 6	2010	211	-	0.47
Valsami S et al. 15	2015	2695	70	2.59
Present study	2018	378	15	4.23

DCT is the cornerstone of the diagnosis of hemolytic disease of the newborn (HDN). In our study, out of 378 patients, DCT was positive in 16 patients. The positivity rate of DCT in our study was 4.23%. Pahuja S et al. drawn similar results by DCT and had a 0.47% rate of immunization. 6

Figure 3: Table 3 :

4

Study	Year	Total Doners	All Doners (ICT Positive)	The Rate of Alloimmunization (%)
Pahuja S et al. 16	2013	7756	4	0.05
Garg N et al. 18	2014	47450	46	0.09
Makroo RN et al. 17	2018	82153	227	0.27
Present study	2018	2050	10	0.49

Figure 4: Table 4 :

69 [Roopam et al. ()] *A prospective study for detection and identification of red cell alloantibodies in multiply*  
70 *transfused thalassemia major patients: 34 th National Congress of Indian Society of Blood Transfusion*  
71 *and Immune Haematol*, J Roopam , J Perkins , J T Susan , N A Choudhury . 2009. p. .

72 [Pahuja et al. (2010)] 'Alloimmunization and red cell autoimmunization in multi-transfused thalassemics of  
73 Indian origin'. S Pahuja , M Pujani , S K Gupta , J Chandra , M Jain . *Hematology* 2010 Jun 1. (3) p.  
74 .

75 [Makroo et al. (2014)] 'Antibody screening and identification in the general patient population at a tertiary care  
76 hospital in New Delhi'. R N Makroo , A Bhatia , N L Rosamma . *India. Indian J. Med Res* 2014 Sep: 140.  
77 (3) p. .

78 [Ansari and Moshtaghian ()] 'Assessment of frequency of alloimmunization and erythrocyte autoimmunization  
79 in transfusion dependent thalassemia patients'. S Ansari , P V Moshtaghian . *Acta MedicaIranica* 2008. 46  
80 (2) p. .

81 [Lamba et al. ()] 'Clinically Significant Minor Blood Group Antigens amongst North Indian Donor Population'.  
82 D S Lamba , R Kaur , S Basu . *Adv Hematol* 2013. 2013. p. 215454.

83 [Sood et al. ()] 'Detection of alloimmunization to ensure safer transfusion practice'. R Sood , R N Makroo , V  
84 Riana , N L Rosamma . *Asian J. Transfu Sci* 2013 Jul: 7. (2) p. 135.

85 [Usman et al. (2011)] 'Frequency of red cell alloimmunization among patients with transfusion dependent beta  
86 thalassemia in Pakistan'. M Usman , M O Saira , M Moinuddin , S Ahmad , R Perveen , S Usman . *Int. J.*  
87 *Hematol Oncol* 2011 Jan: 1: 27. (4) p. .

88 [Valsami et al. (2015)] 'Importance of direct antiglobulin test (DAT) in cord blood: causes of DAT (+) in a  
89 cohort study'. S Valsami , M Politou , T Boutsikou , D Briana , M Papatesta , A Malamitsi -Puchner .  
90 *Pediatr Neonatol* 2015 Aug 1. 56 (4) p. .

91 [Garg et al. (2014)] 'Prevalence of irregular red blood cell antibodies among healthy blood donors in Delhi  
92 population'. N Garg , T Sharma , B Singh . *Transfus Apher Sci* 2014 Jun 1. 50 (3) p. .

93 [Makroo et al. (2018)] 'Prevalence of irregular red cell antibody in healthy blood donors attending a tertiary  
94 care hospital in North India'. R N Makroo , S Rajput , S Agarwal , M Chowdhry , B Prakash , P Karna .  
95 *Asian J Transfus Sci* 2018 Jan: 12. (1) p. .

96 [Nikam et al. ()] 'Prevalence of red cell allo-immunization in repeatedly transfused patients with B-thalassemia  
97 in'. S A Nikam , S B Dama , S A Saraf , C J Jawale , R V Kirdak , R P Chondekar . *Organized by Department*  
98 *of Zoology, D.B.F. Dayanand College of Arts and Science* 2011. p. . (Bio-Resources for Bio-Industries and  
99 Economic Zoology)

100 [Schonewill and Leiden ()] *Red blood cell alloantibodies after transfusion*, H Schonewill , Leiden . 2008. University  
101 Press.

102 [Sircchia et al. (1985)] 'Red cell alloantibodies in thalassemia major'. G Sircchia , A Zanella , A Parravicini , P  
103 Rebulla , F Morelatti , G Maseri . *Transfusion* 1985 Mar 4. 25 (2) p. .

104 [Shukla and Chaudhary ()] 'Red cell alloimmunization in multi-transfused chronic renal failure patients under-  
105 going hemodialysis'. J S Shukla , R K Chaudhary . *Indian J Pathol Microbiol* 1999. 42 p. .

106 [Pandey et al. (2014)] 'Red cell alloimmunization in transfused patients: A silent epidemic revisited'. H Pandey  
107 , S S Das , R Chaudhary . *Asian J. Transfus Sci* 2014 Jul. 8 (2) p. .

108 [Chow et al. ()] 'Red cells, HLA and platelet antibody formation in patients with multiple transfusions'. M P  
109 Chow , H Y Hu , J Y Lyou , J S Lin , C H Yung , A Lee , T D Lee . *Actahaematologica* 1994. 92 (2) p. .

110 [Pahuja et al. (2012)] 'Screening of blood donors for erythrocyte alloantibodies'. S Pahuja , S Kushwaha , N Sethi  
111 , M Pujani , M Jain . *Hematology* 2012 Sep 1. (5) p. .

112 [Nakamura et al. (1984)] *Significance of positive direct antiglobulin test for cord blood -in administration of anti-*  
113 *d-immunoglobulin for postpartum immunoprophylaxis*. *Nihon Sanka Fujinka Gakkaizasshi*, Y Nakamura , I  
114 Sada , S Tanaka , Y Nomura , S Shinagawa . 1984 Apr: 36. p. .