



GLOBAL JOURNAL OF MEDICAL RESEARCH: C

GYNECOLOGY AND OBSTETRICS

Volume 18 Issue 2 Version 1.0 Year 2018

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Non Invasive Follicular Thyroid Neoplasm with Papillary Like Nuclear Features (NIFTP): A Case Report

By Dr. Shweta, Dr. Richa Goyal, Dr. Preeti Joseph John & Dr. Yogesh Gauba

Oncquest Lab Ltd

Abstract- World health organization (WHO) 2017 fourth edition has introduced entity Non Invasive follicular thyroid neoplasm with papillary like nuclear features (NIFTP), which reflects indolent course of this tumor. NIFTP has an established diagnostic criteria including papillary like nuclear features and entire tumor capsule submission to exclude invasion. NIFTP shows RAS type mutation expressed by other follicular lesions. This nomenclature change with strict and reproducible diagnostic criteria will help to better guide the treatment and reduce the burden of over treatment.

Keywords: *non invasive follicular thyroid neoplasm with papillary like nuclear features (niftp), thyroid, invasion, capsule, world health organization (who), papillary thyroid carcinoma.*

GJMR-C Classification: NLNC Code: WK 200



NONINVASIVEFOLICULARTHYROIDNEOPLASMWITHPAPILLARYLIKENUCLEARFEATURESNIFTPACASEREPOR

Strictly as per the compliance and regulations of:



RESEARCH | DIVERSITY | ETHICS

Non Invasive Follicular Thyroid Neoplasm with Papillary like Nuclear Features (NIFTP): A Case Report

Dr. Shweta ^a, Dr. Richa Goyal ^a, Dr. Preeti Joseph John ^p & Dr. Yogesh Gauba ^ω

Abstract- World health organization (WHO) 2017 fourth edition has introduced entity Non Invasive follicular thyroid neoplasm with papillary like nuclear features (NIFTP), which reflects the indolent course of this tumor. The diagnostic criteria for NIFTP includes papillary like nuclear features and complete tumor capsule submission to exclude invasion. NIFTP shows RAS type mutation expressed by other follicular lesions. This strict nomenclature change and reproducible diagnostic criteria will help to guide the management in better way and to reduce the burden of overtreatment.

Keywords: *non invasive follicular thyroid neoplasm with papillary like nuclear features (NIFTP), thyroid, invasion, capsule, world health organization (WHO), papillary thyroid carcinoma.*

I. INTRODUCTION

The fourth edition 2017 of the WHO book on the classification of the tumors of endocrine organs includes new entity 'Non Invasive follicular neoplasm with papillary like nuclear features' (NIFTP). ^[1] This was done additionally to characterize the clinical behavior of this lesion, thus further helping to predict prognosis and manage the patients appropriately. Hence overall helping to prevent overtreatment and decrease the cancer diagnosis burden. ^[2] The WHO has coded unspecified, borderline or uncertain behavior for NIFTP. Hence it is a neoplasm, but not cancer. NIFTP presents mostly similar to other thyroid neoplasms by detection of a nodule on routine examination or incidentally on imaging. Histopathology shows solid, well-circumscribed or encapsulated nodule. Diagnosis is made by strict inclusion and exclusion criteria following submission of the entire capsule for histology and carefully examining the nuclear and architectural features, (Table-1).^[3]

II. CASE REPORT

Forty years old female presented with the complaint of neck swelling from the past few months. Her ultrasonography neck was suggestive of multinodular goiter. Her thyroid function tests were normal. Fine needle aspiration cytology was suspicious of papillary carcinoma. She underwent total thyroidectomy at Mohandai Oswal cancer hospital. The specimen was received for histopathological examination at Oncquest laboratories, situated in Mohandai Oswal cancer hospital. Specimen measured 4.3x4x2cm. The left lobe measured 4x2x2cm. On cut section, it showed a well-circumscribed nodule measuring 1.7x1.7x1.3cm, which was paler than the surrounding thyroid parenchyma. Grossly areas of hemorrhage or necrosis were absent. The right lobe and isthmus showed unremarkable morphology. Sections of the nodule with capsule were embedded. Sections from the left lobe showed encapsulated neoplasm containing medium-sized follicles lined by cuboidal cells displaying nuclear clearing, overlapping and grooving in few cells. An occasional intranuclear inclusion was noted. However, psammoma bodies, tumor necrosis, and mitosis were absent. Papillary architecture (<1%) was noted. Re-grossing was done. Entire nodule and capsule were submitted. Sections showed no capsular or vascular invasion. Hence the final diagnosis of Non Invasive follicular thyroid neoplasm with papillary like nuclear features (NIFTP) was given.

III. DISCUSSION

WHO 2017 fourth edition stated NIFTP as a neoplasm of unspecified, borderline or uncertain behavior. Also, the word cancer was therefore omitted. ^[1] This new terminology reflects key histopathological features of this lesion that is lack of invasion, follicular growth pattern and nuclear features of papillary thyroid carcinoma. These tumors do not show molecular alterations associated with classical papillary thyroid carcinoma, such as BRAF V600E mutations. However, these demonstrate RAS and other mutations associated with follicular pattern thyroid tumors. ^[4] The treatment for NIFTP is simple lobectomy, near total or total thyroidectomy. No further surgery is needed. No postoperative radioactive treatment is given.

Author a: Junior Consultant, Oncquest Lab Ltd.
e-mail: shweta150289@gmail.com

Author a: Lab Head & Consultant Pathologist, Oncquest Lab Ltd.
e-mail: richa.goyal@oncquest.net

Author p: Consultant Pathologist, Oncquest Lab Ltd.
e-mail: dr_priti77@yahoo.com

Author ω: Onco Surgeon, Department of Surgical Oncology, Mohandai Oswal Hospital, Ludhiana, Punjab, India.
e-mail: dryogeshgauba@rediffmail.com



The major advantage of NIFT for cytology is the reduction of SM category (Suspicious for malignancy Bethesda category V) diagnosis. In follicular patterned tumors it is very much dependent on good quality smears and the cytopathologist expertise. The main disadvantage is the false negative diagnosis in the SFN/FN (Follicular Neoplasm or Suspicious for a Follicular Neoplasm Bethesda category IV) leading to the need of a second surgery for total thyroidectomy in cases of the infiltrative follicular variant of papillary thyroid carcinoma (I-FVPTC).^[5]

IV. CONCLUSION

With this approach in mind, the ability to differentiate NIFT from classical papillary thyroid carcinoma will facilitate the conservative surgical management of the patients without radiotherapy or prophylactic central lymph node staging and with more studies adding to the evidence of good prognosis of these tumors.^[3,6] Therefore this nomenclature change will reduce mental burden, overtreatment, financial burden, and other cancer diagnosis related consequences.

Table 1: Diagnostic Criteria for NIFT [3]


| |
|---|
| 1. Encapsulation or clear demarcation * ^a . |
| 2. Follicular growth pattern * ^b with : |
| • <1% Papillae. |
| • No psammoma bodies. |
| • <30% Solid/trabecular/insular growth pattern. |
| 3. Nuclear score 2-3. |
| 4. No vascular or capsular invasion * ^c . |
| 5. No tumor necrosis. |
| 6. No high mitotic activity * ^d . |
| * ^a - Thick, thin, or partial capsule or well circumscribed with a clear demarcation from adjacent thyroid tissue. |
| * ^b - Including microfollicular, normofollicular, or macrofollicular architecture with abundant colloid. |
| * ^c - Requires adequate microscopic examination of the tumor capsule interface. |
| * ^d - High mitotic activity defined as at least three mitoses per 10 high-power fields (X400). |

Legends:

Table 1: Diagnostic Criteria for NIFT [3].

Photograph 1:

- Gross photograph of left lobe showing a well-circumscribed nodule paler than the surrounding thyroid parenchyma.
- Well encapsulated lesion with adjacent normal thyroid parenchyma. (X40).
- Lesion arranged in follicles (X100).
- Medium sized follicles lined by cuboidal cells displaying nuclear clearing and overlapping (X400).



Conflicts of Interest

The report has not been presented or submitted elsewhere fully or in part.

Supports and Acknowledgement

None

Registration Number of Clinical Trial

NA.

REFERENCES RÉFÉRENCES REFERENCIAS

- Lloyd R. V., Osamura R. Y., Klöppel G. & Rosai J. 2017 WHO Classification of Tumours of Endocrine Organs WHO / IARC Classification of Tumours. Lyon, France: IARC Publications.
- Thompson L. D. Ninety-four cases of encapsulated follicular variant of papillary thyroid carcinoma: a name change to Noninvasive Follicular Thyroid Neoplasm with Papillary - like Nuclear Features would help prevent overtreatment. Modern Pathology. 2016; 29 (7): 698-707. doi: 10.1038/modpathol.2016.65.
- Nikiforov Y. E., Seethala R. R., Tallini G., et al. Nomenclature revision for encapsulated follicular variant of papillary thyroid carcinoma. JAMA Oncology. 2016; 2 (8): 1023-1029. doi: 10.1001/jamoaoncol.2016.0386.
- Rivera M., Ricarte-Filho J., Knauf J., et al. Molecular genotyping of papillary thyroid carcinoma follicular variant according to its histological subtypes (encapsulated Vs infiltrative) reveals distinct BRAF and RAS mutation patterns. Modern Pathology. 2010; 23 (9): 1191-1200. doi: 10.1038/modpathol.2010.112.
- Amendoeira I., Maia T., Sobrinho-Simões M. Non-invasive follicular thyroid neoplasm with papillary - like nuclear features (NIFT): impact on the reclassification of thyroid nodules. Endocr Relat Cancer. 2018; 25 (4): 247-58.
- Jug R., Jiang X. Noninvasive follicular thyroid neoplasm with papillary-like nuclear features: an evidence-based nomenclature change. Patholog Res Int. 2017; 2017: 1057252.