Volume 19 Issue 5
Originally issued as G1



Global Journal of Medical Research: G
Veterinary Science and Veterinary Medicine
Volume 19 Issue 1 Version 1.0 Year 2019
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Carcass Characteristics, Hematology, Serum Chemistry, and Enzymes in Broiler Chickens Fed Maggot Meal as a Protein Substitute for Fishmeal

By Mbiba H. F., Etchu K. A. & Ndamukong K.
Institute of Agricultural Research for Development (IRAD)

Abstract- Conventional protein sources used in poultry farming are extensively competed for, by livestock and humans leading to high prices and reduced returns. Focus on better utilization of available alternative feed resources with little or no negative impacts on the health of broilers and consumers is useful. The objective of this research was to assess the performance of carcass characteristics, hematology, serum chemistry, and enzymes in broiler chickens fed maggot meal as a protein substitute for fishmeal. 225 Tropical Broc day old chicks brooded for two weeks and fed the control diet, were distributed in a completely randomized block design with five treatments and three replicates each consisting of the starter and finisher phases and the experiment conducted for eight weeks. Diets were compounded with maggot meal (MM) replacing FM at 0%, 25%, 50%, 75% and 100%.

Keywords: *maggot meal, hematology, serum chemistry, enzymes, carcass characteristics.*

GJMR-G Classification: *NLMC Code: QW 70*



Strictly as per the compliance and regulations of:



© 2019. Mbiba H. F., Etchu K. A. & Ndamukong K. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Carcass Characteristics, Hematology, Serum Chemistry, and Enzymes in Broiler Chickens Fed Maggot Meal as a Protein Substitute for Fishmeal

Mbiba H. F. [α], Etchu K. A. [σ] & Ndamukong K. [ρ]

Abstract- Conventional protein sources used in poultry farming are extensively competed for, by livestock and humans leading to high prices and reduced returns. Focus on better utilization of available alternative feed resources with little or no negative impacts on the health of broilers and consumers is useful. The objective of this research was to assess the performance of carcass characteristics, hematology, serum chemistry, and enzymes in broiler chickens fed maggot meal as a protein substitute for fishmeal. 225 Tropical Broc day old chicks brooded for two weeks and fed the control diet, were distributed in a completely randomized block design with five treatments and three replicates each consisting of the starter and finisher phases and the experiment conducted for eight weeks. Diets were compounded with maggot meal (MM) replacing FM at 0%, 25%, 50%, 75% and 100%. On the last day of week 8, 30 birds, 2 from each replicate, were randomly selected and weighed, each bird slaughtered and allowed to bleed for 2 minutes while blood samples collected from one bird per replicate were put into 2 tubes (one with EDTA and the other without) for studies of Hb, WBCS and RBCs, total protein, albumin, and globulin, AST and ALT. Dressed, eviscerated, carcass parts, liver, and gizzard, weights were taken. Then averages from each replicate statistically analyzed for any significant differences. Results showed better carcass characteristics and lower amounts of WBCs in the treatment groups with maggot meal, stable values of RBCs and Hb, no defined trend in variations of total protein, globulin, and enzymes studied between the various treatments. Given that broilers with the best carcass characteristics performance were those with 100% maggot meal inclusion and that physiological parameters were not deviated from normal values in birds fed experimental diets, it can be concluded from this study that maggot meal can completely replace fish meal at 5% in broiler feed for better carcass characteristics, and a stable physiological profile.

Keywords: maggot meal, hematology, serum chemistry, enzymes, carcass characteristics.

Author α: Institute of Agricultural Research for Development (IRAD), Bambui, PMB 80, Bamenda, Cameroon.
e-mail: hasan5basri@gmail.com
Author σ: Institute of Agricultural Research for Development (IRAD), Nkolbisson, Yaounde, Cameroon. e-mail: etchu74@gmail.com
Author ρ: Department of Animal Sciences, University of Buea, Box 63, Buea-Cameroon. e-mail: kjnndamukong@yahoo.com

I. Introduction

Broiler production is one of the main areas in animal farming that involves quite a large section of the population either skilled or unskilled. One important advantage in this is the fact that poultry meat is very rich in unsaturated fatty acids as against saturated fatty acids. Both turkey and chicken have about 30% saturated fatty acids, 43% monounsaturated fatty acids, and 22% polyunsaturated fatty acids. The ratio is a clear indication that poultry meat may stand a better position as a more healthful alternative for red meat (Encyclopaedia Britannica, 2014). Notwithstanding there are some challenges in areas of nutrition, health, and management (Awoniyi, 2004). Feeding alone accounts for 60-70% of the total production cost in modern poultry production systems (Smith, 1990; Church, 1991; Wilson and Beyer, 2000). Conventional protein sources used are extensively competed for, by other livestock and humans (Gadzirayi *et al*., 2012), thereby leading to very high prices and reduced returns. Any attempt to improve commercial poultry production and increase its efficiency, therefore needs to focus on better utilization of alternative but available feed resources. Knowledge of nutritional characteristics of these feeds and optimal levels of inclusion in rations and optimum combination of ingredients is useful (Kamalzadeh *et al*., 2008). Competition for conventional protein sources has prompted researchers to embark on research for alternatives like MM that are cheaply available and comparable to FM. The crude protein content of maggots is high (39-63%) (Aniebo and Owen, 2010) and akin to that of fish meal (Veldkamp *et al*., 2012).

Even though MM may reduce competition between man and other livestock, there is very high need to investigate its health implications in broiler and humans as well the effects on performance of broilers (Awoniyi, 2004). A study of the carcass characteristics, blood indices and enzymes should correlate the benefits of MM as a protein source in poultry feed vis-à-vis the physiological status of the birds. The study of these

© 2019 Global Journals

physiological parameters will serve to bridge the gap as work done in Cameroon on this aspect is scanty.

II. Materials and Methods

a) Study area

The study was carried out at Muyuka Agro-Industrial Farm situated between latitudes $4^0$16″ and $4^0$23″N and longitudes $9^0$21″ and $9^0$28″E in the fourth agro-ecological zone of Cameroon (AEZ IV). Muyuka on the windward side of mount Cameroon, experiences very high temperatures ranging from 25°C during the rainy season (March to September) to about 30°C in the dry season (October to March). The climate is typical of the equatorial type. The monthly rainfall ranges between 9.2mm to 374.1mm, the lowest realized in January while the heaviest is in August.

b) Birds

Two hundred and twenty-five Tropical Broc day-old broilers used for the experiment fed on the control diet (Table 1) during the brooding period of two weeks. Coal pots provided heat supplementation and prophylactic measures employed. Daylight served as the main lighting source during the day and electric current at night; lanterns served as the illuminating source in cases of power outage.

c) Experimental design

The trial used a completely randomized block design (CRBD) in which two weeks old chicks were randomly allocated to 5 treatments, each containing 45 birds; and each treatment had three replicates with 15 birds each. Diets formulated using maggot meal (MM) substituted fishmeal (FM) at graded levels; 0%, 25%, 50%, 75% and 100% at both the starter and finisher phases. Mineral and vitamin premixes customary to poultry production, oyster shell, salt, and bone served as complements (Tables 1 and 2).

Birds lived on deep litter while enjoying natural ventilation, Feed, and water *ad libitum*. Broiler starter, 23% crude protein, sustained for four weeks followed by broiler finisher, 19% crude protein from the end of the 4^{th} week till slaughter (8^{th} week).

d) Ration formulation for experimental diets

The feed ingredients used in this study included: Yellow maize, fishmeal, maggot meal, groundnut cake, kernel cake, soybean cake, wheat bran, and premixes. Proximate analysis gave crude protein and ME values for maggot meal while the nutrient master plan (*livestock feeds*) provided those for other ingredients. Levels of inclusion of protein (maggot meal and fishmeal) and energy sources to meet the protein and energy requirements were manipulated using Pearson's Square method and the various percentages calculated as indicated in Tables 1 and 2 with the help of a nutrient master plan which gave the protein and energy contents of each ingredient.

After all the ingredients were measured and put together at the same spot on the cardboard paper, they were mixed with the hand, making at least three complete turns to ensure proper mixing, and then put into bags with treatment labels.

Table 1: Percentage inclusion levels and chemical composition of experimental diets for broiler starter (weeks 1-4)

Ingredients	Treatment Composition				
	Control (T_0, 0%)	T_1 (25%)	T_2 (50%)	T_3 (75%)	T_4 (100%)
Maize	48.00	48.00	48.00	48.00	48.00
Soya bean cake(SBC)	12.50	12.50	12.50	12.50	12.50
Fishmeal (FM)	5.00	3.75	2.50	1.25	0.00
Maggot meal (MM)	0.00	1.25	2.50	3.75	5.00
Groundnut cake (GNC)	17.00	17.00	17.00	17.00	17.00
Palm kernel cake (PKC)	5.00	5.00	5.00	5.00	5.00
Wheat bran (WB)	10.50	10.50	10.50	10.50	10.50
Premix	0.75	0.75	0.75	0.75	0.75
Bone meal (BM)	0.50	0.50	0.50	0.50	0.50
Salt	0.25	0.25	0.25	0.25	0.25
Oyster shell	0.50	0.50	0.50	0.50	0.50
Total	100.00	100.00	100.00	100.00	100.00
Crude Protein (%CP)	23.20	22.99	22.79	23.58	22.37
Metabolizable energy (ME) Kcal/Kg	2882.72	2888.25	2893.78	2899.31	2904.85

© 2019 Global Journals

Table 2: Percentage inclusion levels and chemical composition of experimental diets for broiler finisher weeks 5-8

Ingredients	Treatment Composition				
	Control (T_0 0%)	T_1 (25%)	T_2 (50%)	T_3 (75%)	T_4 (100%)
Maize	48.00	48.00	48.00	48.00	48.00
Soya bean cake(SBC)	9.00	9.00	9.00	9.00	9.00
Fishmeal (FM)	5.00	3.75	2.50	1.25	0.00
Maggot meal (MM)	0.00	1.25	2.50	3.75	5.00
Groundnut cake (GNC)	10.00	10.00	10.00	10.00	10.00
Palm kernel cake (PKC)	15.00	15.00	15.00	15.00	15.00
Wheat bran (WB)	10.00	10.00	10.00	10.00	10.00
Premix	0.75	0.75	0.75	0.75	0.75
Bone meal (BM)	1.00	1.00	1.00	1.00	1.00
Salt	0.25	0.25	0.25	0.25	0.25
Oyster shell	1.00	1.00	1.00	1.00	1.00
Total	100.00	100.00	100.00	100.00	100.00
Crude Protein (%CP)	19.53	19.36	19.12	18.91	18.70
Metabolizable energy (ME) Kcal/Kg	2811.57	2817.10	2822.63	2828.16	2833.70

e) Carcass characteristics

Determination of carcass characteristics took place on the last day of the 8th week. After recording live weights of 30 randomly selected birds, two from each replicate, we slaughtered each bird and allowed to bleed for about two minutes before putting in hot water for almost a minute to soften the skin for easy plucking. Dressed weight represented the bulk after removal of the shanks, crop, entrails and other organs. The carcass parts consisted of head, neck, wings, breast, back, thigh and drumstick.

We discarded entrails and weighed eviscerated birds, livers, gizzards, and carcass parts. Then averages from each replicate statistically analyzed for any significant differences between treatments.

f) Studies of hematological parameters, serum chemistry, and enzymes

Studies of hematological parameters (hemoglobin, white and red blood cells), serum chemistry (total protein, albumin, and globulin), and enzymes (aspartate amino-transaminase and alanine amino-transaminase) were carried out at the end of the experiment. Blood was collected at the time of slaughter for carcass analysis into 30 tubes from 15 birds; fifteen tubes had the anticoagulant ethylene diamine tetra-acetate (EDTA) to prevent clotting, and the rest of the cylinders had no anticoagulant. Two hoses (one with EDTA and the other without the anticoagulant) were used to collect blood from one bird per replicate.

g) Method of data processing and analysis

Data were organized in Microsoft Office Excel Version 2010 and analyzed using SPSS 17.0 (SPSS Inc, 2008). Data screened for exploration using Kolmogorov Smirnov and Shapiro Wilk tests revealed that the data departed from the normal distribution. The non-parametric test, notably Kruskal Wallis test, was then used to compare groups for significant differences (Nana, 2012) and the Dunnett T_3 test used for paired comparisons. We took measurements of central tendencies and dispersion, presented the data using statistical tables and charts, and discussed at the 95% CL (Alpha=0.05).

III. Results and Discussion

a) The chemical composition of experimental diets

Table 3 reveals a drop in the CP value of analyzed components for all the treatments, except for the 100% MM. There was a reasonable increase in ME values, especially in 75% and 100% MM.

Table 5 shows a drop in the crude protein content of the control and 25% MM and a slight increase in the rest of the analyzed feed composition compared to the calculated feed composition. There was a considerable increase in ME values of the evaluated components compared to the premeditated constituents across the treatments.

The differences observed in the calculated and the analyzed compositions for both the starter and finisher diets may have been due to variations between the tabulated nutrient content values used in the calculations and the actual nutrient contents of the ingredients used in the experiment. Doing proximate analysis for all components before formulating and compounding the various feeds for the trial keeps this situation in check. Increase in ME with increasing maggot meal in the diets may be explained by the high fats in the maggots which release a lot of energy when oxidized (Adeniji, 2007).

© 2019 Global Journals

Table 3: Percentage inclusion levels and chemical composition of experimental diets at the starter phase (weeks 1-4)

Ingredients	Control (T_0 0%)	T_1 (25%)	T_2 (50%)	T_3 (75%)	T_4 (100%)
Maize	48.00	48.00	48.00	48.00	48.00
Soya beans cake (SBC)	12.50	12.50	12.50	12.50	12.50
Fishmeal (FM)	5.00	3.75	2.50	1.25	0.00
Maggot meal (MM)	0.00	1.25	2.50	3.75	5.00
Groundnut cake (GNC)	17.00	17.00	17.00	17.00	17.00
Palm kernel cake (PKC)	5.00	5.00	5.00	5.00	5.00
Wheat bran (WB)	10.50	10.50	10.50	10.50	10.50
Premix	0.75	0.75	0.75	0.75	0.75
Bone meal (BM)	0.50	0.50	0.50	0.50	0.50
Salt	0.25	0.25	0.25	0.25	0.25
Oyster shell	0.50	0.50	0.50	0.50	0.50
Total	100.00	100.00	100.00	100.00	100.00
Calculated composition					
Crude Protein (%CP)	23.20	22.99	22.79	23.58	22.37
Metabolizable energy (ME) Kcal/Kg	2882.72	2888.25	2893.78	2899.31	2904.85
Analyzed composition					
Crude Protein (%CP)	20.4	18.9	20.4	20.7	22.4
Metabolizable energy (ME) Kcal/Kg	3114.9	3121.7	3121.5	3359.8	3205.5

Table 4: Percentage inclusion levels and chemical composition of experimental diets at the finisher phase (weeks 5-8)

Ingredients	Control (T_1 0%)	T_2 (25%)	T_3 (50%)	T_4 (75%)	T_5 (100%)
Maize	48.00	48.00	48.00	48.00	48.00
Soya beans cake (SBC)	9.00	9.00	9.00	9.00	9.00
Fishmeal (FM)	5.00	3.75	2.50	1.25	0.00
Maggot meal (MM)	0.00	1.25	2.50	3.75	5.00
Groundnut cake (GNC)	10.00	10.00	10.00	10.00	10.00
Palm kernel cake (PKC)	15.00	15.00	15.00	15.00	15.00
Wheat bran (WB)	10.00	10.00	10.00	10.00	10.00
Premix	0.75	0.75	0.75	0.75	0.75
Bone meal (BM)	0.50	0.50	0.50	0.50	0.50
Salt	0.25	0.25	0.25	0.25	0.25
Oyster shell	0.50	0.50	0.50	0.50	0.50
Total	100.00	100.00	100.00	100.00	100.00
Calculated composition					
Crude Protein (%CP)	19.53	19.36	19.12	18.91	18.70
Metabolizable energy (ME) Kcal/Kg	2811.57	2817.10	2822.63	2828.16	2833.70
Analyzed composition					
Crude Protein (%CP)	17.1	16.0	20.4	19.0	20.0
Metabolizable energy (ME) Kcal/Kg	3253.4	3166.6	3015.7	3418.1	3296.6

b) Carcass analysis

Table 5 indicates the effects of graded level inclusion of MM in broiler diets on carcass characteristics. Live weight, dressed weight, eviscerated weight, carcass characteristics and organs were all significantly different ($P<0.001$) between treatments except for the liver which was not significantly different ($P>0.05$). Generally, the weight of carcass parts increased from T0 to T4. This increase in weight agrees with the findings of Hwangbo *et al*. (2009). The general increase in bulk of the carcass parts with increase inclusion of MM in the diets may have been due to the live weight which also increased with increased levels of maggot inclusion. Agbede and Aletor (2003) found no significant change in all carcass characteristics and organs except for the relative weights of the neck and heart which were significantly higher in diets containing 7.24% of gliricidia leaf protein concentrate in place of FM. Awoniyi *et al.* (2003) instead reported that equal replacement of FM with MM in broiler chicken diet had

© 2019 Global Journals

no significant effect on the relative length, breadth or weight of muscles of key economic importance in chickens. His report is similar to that of Quinton (2011). Amal *et al*., (2013), Meseret *et al*., (2012) and Bello *et al*., (2012) found no significant ($p>0.05$) differences between all treatments groups in Live weight, eviscerated weight of carcass cuts, dressing percentage, edible inner organs (liver, gizzard, and heart).

The weight of the liver was not significantly different between the control and experimental diets, although higher ($P>0.05$) in treatment groups than in non-experimental diet. Hwangbo *et al*. (2009) and Okah and Onwujiariri (2012) also found no significance ($P>0.05$) in weight of the liver amongst treatments. This indifference in bulk of the liver may have been an indication that there was no infection in the maggot meal that could cause undesirable effects on the nutrition of broilers as indicated by Hwangbo *et al*. (2009) and Okah and Onwujiariri (2012). These results differed from those of Teguia *et al*. (2002) who obtained proportional increases in weights of the liver and gizzard from the control through treatment groups and linked it to toxicity. Live and dressed weights were significantly higher ($P<0.001$) in treatment groups than in control. T_4 was, in turn, higher ($P<0.001$) than the rest of the treatment groups which didn't differ significantly ($P>0.05$) between themselves in live and dressed weights. Eviscerated weight in T_4 was significantly ($P<0.001$) higher than in the rest of the treatments which were, in turn, greater than the control though not significantly ($P>0.05$) different. The above findings agree with those of Gadzirayi *et al*., (2012) using mature *Moringa oleifera* leaf meal as a protein supplement to soybean meal but goes contrary to the outcome of Yisa *et al*., (2013) who stated that complete withdrawal of FM implied poorer development of the meat yielding parts.

Table 5: Comparison of carcass parts between treatments

Carcass parts	Average Weight (Mean ± SE) g					
	T_0 (N=24)	T_1 (N=24)	T_2 (N=24)	T_3 (N=24)	T_4 (N=24)	Kruskal Wallis Test
Live weight (g)	1466.667±13.03	1616.667±13.003^{ab}	1616.667±4.915^{ac}	1666.667±34.403^{bc}	1900.000±22.522	X^2=77.517 P<001
Dressed weight (g)	1366.667±13.03	1483.333±17.720^{ab}	1483.333±4.915^{ac}	1516.667±35..441^{bc}	1800.000±22.522	X^2=69.471 P<001
Eviscerated weight (g)	1166.667±21.43^{abc}	1216.667±13.003^{ade}	1233.333±4.915^{bdf}	1266.667±38.385^{cef}	1416.667±13.003	X^2=48.809 P<001
Head (g)	41.840±0.236^{a}	43.867±0.455^{bc}	43.913±0.210^{bd}	45.257±1.322^{acde}	47.167±0.615^{e}	X^2=34.850 P<001
Neck (g)	74.870±0.845^{a}	79.912±0.805^{bc}	84.153±0.299^{d}	84.000±03.37^{abde}	80.787±0.943^{ce}	X^2=28.617 P<001
Wing (g)	126.077±2.321	136.137±2.022^{a}	146.297±0.232^{b}	142.663±3.145^{ab}	167.890±1.997	X^2=73.667 P<001
Breast (g)	252.380±4.603^{a}	260.150±7.789^{abc}	280.483±3.282^{bd}	286.823±6.730^{cd}	364.863±4.096	X^2=67.717 P<001
Back (g)	218.307±5.862^{abc}	220.352±3.941^{ade}	214.233±1.086^{bdf}	208.947±8.047^{cef}	240.160±2.341	X^2=20.117 P<001
Drumstick (g)	156.213±2.658	175.232±3.426^{ab}	172.470±0.321^{ac}	173.907±4.142^{bc}	213.957±2.982	X^2=65.450 P<001
Thigh (g)	170.117±2.789^{abc}	174.413±3.178^{ade}	173.470±0.908^{bdf}	187.540±7.708^{cefg}	197.750±2.706^{g}	X^2=29.750 P<001
Liver (g)	38.587±1.310^{abcd}	41.147±0.368^{aefg}	41.737±0.296^{behi}	39.483±1.409^{cfhj}	41.643±0.459^{dgij}	X^2=10.483 P=.033
Gizzard (g)	44.847±0.185^{a}	47.523±.556b^{c}	43.090±0.696^{a}	49.005±1.121^{bd}	47.4467±.284^{cd}	X^2=31.167 P<001

$^{a, b, c, d, e, f, g, h, I, j}$ Dunnett T_3: Paired comparison between treatments and within weeks; pairs with the same letter are not significantly different at the 0.05 Level.

c) Hematology, serum chemistry, and enzyme studies

The results presented in Table 6 reveal that there was a significant difference ($P<0.001$) between treatments in the hematological parameters, serum chemistry, and enzymes studied, except for albumin which was the same throughout the control and treatment groups. T_2 recorded the lowest total protein and globulin and T_3 the highest. ALT was significantly lower ($P<0.001$) in control compared to the rest of the treatment groups and topmost in T_2, but not significantly different ($P>0.05$) from the rest of the treatment groups.

© 2019 Global Journals

Table 6: Comparison of hematology, serum chemistry, and enzyme parameters between treatments

Parameters	Average (Mean ± SE)					
	T0 (N=6)	T1 (N=6)	T2 (N=6)	T3 (N=6)	T4 (N=6)	Kruskal Wallis Test
Total protein (g/dl)	34.667±0.428^{abc}	34.000±1.228^{adef}	30.667±1.322^{bdg}	36.333±1.095^{ce}	32.667±0.354^{fg}	X^2=11.241 P=0.024
Albumin (g/dl)	1.000±0.000	1.000±0.000	1.000±0.000	1.000±0.000	1.000±0.000	X^2=0.000 P=1.000
ALT (U/L)	5.333±0.260	9.000±0.590^{abc}	10.333±0.260^{a}	8.333±0.098^{b}	9.333±0.098^{c}	X^2=67.419 P<001
AST (U/L)	211.333±1.722^{ab}	222.333±2.871^{c}	185.000±4.283	219.333±3.589^{acd}	211.000±0.742^{bd}	X^2=49.885 P<001
Globulin (g/dl)	34.333±0.491^{ab}	33.000±1.228^{acde}	29.667±1.322^{cf}	35.333±1.095^{bd}	31.000±0.450^{ef}	X^2=15.509 P<001
Hb (g/dl)	13.000±0.170	11.667±0.260^{abc}	11.000±0.170^{ade}	11.667±0.260^{bdf}	11.667±0.260^{cef}	X^2=31.657 P<001
RBC/L (×10^6)	2.237±3.490×10^{4abcd}	1.400±4.500×10^{4aefg}	2.000±5.898(×10^{4behi}	1.967±9829.464^{cfhj}	1.767±9829.464^{digj}	X^2=81.987 P<001
WBC/L	1300.000±17.025^{ab}	1200.000±34.050^{acd}	800.000±128.537^{ce}	950.000±118.260^{bde}	1666.667±26.006	X^2=54.345 P<001

$^{a, b, c, d, e, f, g, h, I, j}$ Dunnett T_3: Paired comparison between treatments and within weeks; pairs with the same letter are not significantly different at the 0.05 Level.

ALT= Alanine Amino-Transaminase, AST= Aspartate Amino-Transaminase, RBC= Red Blood Cell, WBC= White Blood Cell, Hb= Hemoglobin.

RBC and WBC were low in T_1 and T_2 and highest in T_0 and T_4 respectively. Generally, the values of WBC presented in this study (0.800×103-1.667×103/L) are very low compared to those (2.35×106-3.39×106/L) reported by Nworgu *et al.* (2007) using fluted pumpkin leaf extract. The low WBC in this report is an indication of the absence of infections in these birds, supporting the fact that MM caused no health hazards. RBCs reported by Nworgu *et al.* (2007) were higher than those in this study, the highest (2.237×106/L) being the control, even though not significantly different from the rest of the treatment groups. The range of RBCs obtained in this study falls within the range for broilers (1.5 to 4.5 x 106 cells/µL) as stated by Martinho (2012). Hemoglobin was higher (P<0.001) in control than in the treatment groups which did not differ (P>0.05) amongst themselves, but generally it was still higher than the value (8.22g/dl) obtained by Orawan and Aengwanich (2007) for broiler chickens. However, these values fell within the normal range of 7-13g/dl (Bello *et al*., 2012). Total protein values in this experiment were high surely as a result of higher levels of globulin. Total protein range (3.10-3.70g/dl) and globulin (1.10-1.40) reported by Nworgu *et al.* (2007) are far lower than those obtained in this study (30.667-34.667g/dl and 29.667-34.333g/dl respectively). AST was lowest in T_2 and highest in T_1. The values of alanine transaminase (ALT) reported in this study (5.333-10.333U/L) were very low compared to 23.00-24.84U/L obtained by Nworgu *et al.* (2007); meanwhile, aspartate transaminase (AST) was exceedingly higher (185.000-222.333U/L) than what Nworgu *et al.* (2007) obtained (17.000-21.110U/L). The exceptionally high AST in this report could be the result of oxidative stress.

IV. Conclusion

Conclusions derived from this study follows thus: Better carcass characteristics in treatments with maggot meal is an indication that maggot meal is not inferior to fishmeal. Lower amounts of white blood cells in the treatment groups with maggot meal than in control indicate the absence of any infection in the system attributable to MM. The stable values of RBCs and Hb were an indication that the birds were not suffering from anemia, thereby indicating that MM did not upset the physiological status of the birds. The values of total protein, globulin, and enzymes studied did not show any defined trend in their variations between the various treatments. Given that broilers with the best carcass characteristics performance were those with 100% maggot meal inclusion and that physiological parameters were not deviated from normal values in birds fed experimental diets, it can be concluded from this study that maggot meal can entirely replace fish meal at 5% in broiler feed for better carcass characteristics, and a stable physiological profile.

Acknowledgment

We acknowledge the great support from Mr. ZHU LYING who granted us free access to his poultry farm (Muyuka Agro-Industrial Farm) and materials to carry out this research from the beginning till the end, and the moral support from his staffs.

© 2019 Global Journals

References Références Referencias

1. Adeniji, A. A. (2007).Effect of replacing groundnut cake with maggot meal in the diet of broilers. *International Journal of Poultry Science,* 6 (11): 822-825.
2. Agbede J. O. and Aletor V. A. (2003). Evaluation of fish meal replaced with leaf protein concentrate from glycricidia in diets for broiler-chicks: effect on performance, muscle growth, haematology and serum metabolites. *International Journal of Poultry Science,* 2 (4): 242-250.
3. Amal O. A., Mukhtar A. M., Mohamed K. A. and Ahlam A. H. (2013). Use of Halfa Bar Essential Oil (HBO) as a Natural Growth Promoter in Broiler Nutrition. *International Journal of Poultry Science* 12 (1): 15-18.
4. Aniebo, A. O. and Owen, O. J. (2010). Effects of age and method of drying on the proximate composition of housefly larvae (*Muscadomestica)* meal (HFLM). *Pakistan Journal of Nutrition,* 9 (5): 485-487.
5. Awoniyi, T. A. M., Aletor, V. A. and Aina, J. M. (2003). Performance of broiler - chickens fed on maggot meal in place of fishmeal. *International Journal of Poultry Science,* 2 (4): 271-274.
6. Awoniyi, T. A. M., Adetuyi, F. C. (2004). Microbiological investigation of maggot meal, stored for use as livestock feed component. *Journal of Food, Agriculture and Environment,* 2 (3 and 4): 104-106.
7. Bello M. K., Oyawoye O. E. and Bogoro E. S. (2012). Performance, carcass characteristics and blood composition of broilers fed varying levels of palm kernel meal (*elaiseguinensis*) supplemented with different levels of fishmeal. *International Journal of Poultry Science,* 11 (1): 73-77.
8. Church, D. C. (1991). *Livestock Feeds and Feeding (3rd Ed.)*. Prentice Hall. Incorporation cliffs, New Jersey. USA, 465 pp.
9. Encyclopaedia Britannica. (2014). Poultry processing.
10. Gadzirayi, C. T., Masamha, B., Mupangwa, J. F and Washaya, S. (2012).Performance of broiler chickens fed on mature *Moringaoleifera* leaf meal as a protein supplement to soybean meal. *International Journal of Poultry Science,* 11 (1): 5-10.
11. Hwangbo, J., Hong, E. C., Jang, A., Kang, H. K., Oh, J. S., Kim, B. W. and Park, B. S. (2009). Utilization of housefly maggots, a feed supplement in the production of broiler chickens. *Journal of Environmental Biology*, 30 (4): 609-614.
12. Kamalzadeh, A., Rajabbeygi, M. and Kiasat, A. (2008). Livestock production systems and trends in livestock industry in Iran. *Journal of Agricultural and Social Science,* 4: 183-188.
13. Martinho F. (2012).Blood transfusion in birds. *Revista Lusófona de Ciência e Medicina Veterinária*, 5: 1-30.
14. Meseret G., Berhan T. and Tadelle D. (2012). Effects of Replacing Peanut Seed Cake with Brewery Dried Yeast on Laying Performance, Egg Quality and Carcass Characteristics of Rhode Island Red Chicken. *International Journal of Poultry Science* 11 (1): 65-72.
15. Nworgu, F. C., Ogunbero, S. A. and Sotesi K. S. (2007). Performance and some blood chemistry indices of broiler chicken served fluted pumpkin leaves extract supplement. *American-Eurasian Journal of Agriculture.*
16. Okah, U. and Onwujiariri, E. B. (2012). Performance of finisher broiler chickens fed maggot meal as a replacement for fish meal. *Journal of Agricultural Technology*, 8 (2): 471-477.
17. Orawan, C. and aengwanich, W. (2007). Blood cell characteristics, Haematological values and average daily gained weight of Thai indigenous, crossbred and broiler chickens. *Pakistan Journal of Biological Sciences*, 10 (2): 302-309.
18. Quinton P. (2011). Comparison of the carcass characteristics of broilers chicks grown on a diet containing either *Muscadomestica* larvae meal, fish meal or soya bean meal as the main protein source (Master's thesis).*Stellenbosch University, Faculty of Agri Sciences, Stellenbosch, South Africa.*
19. Smith, A .J. (1990). *Poultry*. *The Tropical Agriculturalist*. Tropical Centre for Agriculture and Rural Co-operation, Macmillan Education Ltd, London, United Kingdom.
20. Téguia A., Mpoame M. & OkourouMba J. A. (2002). The Production Performance of Broiler Birds as Affected by the Replacement of Fish Meal by Maggot Meal in the Starter and Finisher Diets. *TROPICULTURA*, 20(4): 187-192.
21. Veldkamp, T., van Duinkerken, G., van Huis, A., lakemond, C., Ottevanger, M. M., Bosch, E. G. and van Boekel, M. A. J. S. (2012). Insects as a sustainable feed ingredient in pig and poultry diets - a feasibility study. Wageningen UR Livestock Research and Central Veterinary Institute of Wageningen UR (University & Research centre). Wageningen.
22. Wilson, K. J. and Beyer, R. S. (2000). Poultry Nutrition Information for the Small Flock. Kansas State University Agricultural Experiment Station and Cooperative Extension Service. http://www.oznet.ksu.edu.
23. Yisa A. G., Edache J. A., Udokainyang A. D. and Iloama C. N. (2013). Growth Performance and Carcass Yield of Broiler Finishers Fed Diets Having Partially or Wholly Withdrawn Fish Meal. *International Journal of Poultry Science* 12 (2): 117-120.

© 2019 Global Journals