

Artificial Intelligence formulated this projection for compatibility purposes from the original article published at Global Journals. However, this technology is currently in beta. *Therefore, kindly ignore odd layouts, missed formulae, text, tables, or figures.*

CrossRef DOI of original article:

1	Result of a Questionnaire Survey of 61 Females on their
2	Satisfaction with Fish, Calcium and Protein Intake
3	Naomi Katayama ¹ and Shoko Kondo ²
4	¹ Nagoya Womens University
5	Received: 1 January 1970 Accepted: 1 January 1970 Published: 1 January 1970
6	

7 Abstract

⁸ Since the calcium intake has not reached the target amount of 600 800 mg per day in the

⁹ Japanese diet, calcium intake is recommended in the daily diet. And in recent years, Japan

¹⁰ has been promoting protein intake for the elderly to prevent sarcopenia and frailty. Japan also

recommended to take EPA and DHA to maintain smooth blood vessels and memory

¹² maintenance. Therefore, as a first step to understanding the actual situation, this study

conducted a questionnaire survey on the information of fish rich in EPA and DHA and the

¹⁴ intake of Ca and protein.Sixty-one participants were given a self-anubnustered questionnaire

¹⁵ regarding fish, calcium and protein intake. The questionnaire method is self-administered, and

¹⁶ the questions are about age, height, weight, and desired purchase price of fish. We asked the

¹⁷ following questions, frequency of eating fish, frequency of eating meat, frequency of eating

18 eggs, and frequency of eating bean products.

19

20 Index terms—questionnaire surveys, consumption of fish, calcium intake, protein intake.

²¹ 1 Introduction

n Japan, the Ministry of Health, Labor and Welfare publishes dietary standards for Japanese every five years.
According to the Japanese Dietary Intake Standards issued by the Ministry of Health, Labor and Welfare in
2020, the daily protein intake for adults is 60g for males and 50g for females.

Per capita consumption of seafood in Japan continues decline. According to the Japanese "Food Supply and Demand Chart," the consumption of edible seafood per person per year, peak intake of fish per person is 40.2 kg in 2001, but peak intake of fish per person is 24. 6 kg in 2016 which is 1.1 kg less than the previous year. This fish intake is about the same level as in the late 1930s.In recent years, in Japan, protein intake is starting to decrease. This is thought to be due to factors such as the aging of the population. Calcium intake in Japan

30 is much less than the recommended amount of 800~700mg for males and 650~600mg for females. The recent

calcium intake in Japan is 470~550mg for males and 400~500mg for females. That is nearly 250mg less than the recommended amount.

Data from 1995 to 2015 show that Japanese protein intake is declining. Total protein may be sufficient, but animal proteins are too few, and it can be said that the Japanese lack "good quality protein."That means that the amino acid score does not exceed 100.

The 61 females (18-28) who participated in a health class were asked about satisfaction with their intake of fish, calcium, and protein intake.

38 **2** II.

³⁹ **3** Materials and Methods

⁴⁰ Participants were 61 females who were briefed about the study and signed a consent form. The questions consisted ⁴¹ of the following seven items. 1) Do you like fish to eat? 2) Do you know a commercial fish that can eat up to 42 the bones? 3) Would you like to eat a fish that can be eaten up to the bones of a commercial product? 4) Do 43 you feel that you are deficient in calcium intake daily? 5) Do you want to take calcium positively? 6) Do you 44 feel that you are deficient in protein intake daily? 7) Do you want to take protein positively? Participants self-

administered responses to a sevenitem questionnaire. In addition, participants were asked about their frequency
 of consumption of fish, meat, eggs, and beans using a self-administered questionnaire. The participants also

answered the desired purchase price of the fish whose bones are edible.

48 **4** III.

⁴⁹ Results 61 participants aged 18 to 28 years (see Table 1.), and the average age \pm standard deviation was 21.1 \pm 2.0. Weight was 52.9 \pm 7.1, and height was 158.4 \pm 5.9 (see Table ??

⁵¹ **5** .).

Table 3 shows the results of a seven-item selfadministered questionnaire conducted on the participants. Among the participants, 83.6% answered that they like to fish. The participants of 75.4% responded that they know a commercial fish that can eat up to the bones. Among the participants, 82.0% answered that they would like to eat a commercial product of fish whose bones are edible. In addition, 68.9% of the participants answered that they felt calcium deficiency in their daily lives. And 95.1% of the participants insisted on positive calcium intake. Similarly, 32.8% of the participants felt that they were not getting enough protein in their daily lives. The participants of 90.2% responded that they want to take protein positively.

Participants filled out a self-administered questionnaire about the frequency of consumption of fish, meat,
eggs, and beans (see Table 4.). Many participants responded to the question as follows. They ate fish 1-2 times
a week, meat 3-4 times a week, eggs 3-4 times a week, and beans 1-2 times a week.

In addition, participants answered the following questions: 6.6% of participants ate meat every day, 23.0% of participants ate eggs every day, and 18.0% of participants ate beans every day. However, none of the participants ate fish every day. Table 5 shows the results of a self-administered questionnaire survey of participants regarding

the desired purchase price of commercially available fish that even the bones can be eaten. As a result, among

the participants, the most requested price was 200 yen (about 1.5 US dollars). However, the reality is that the

67 price of commercially available fish with edible bones in 400 yen (about 3-4 US dollars). It is about twice the 68 purchase price requested by the participants.years

⁶⁹ 6 IV.

70 7 Discussions

71 From the results of a self-administered questionnaire survey conducted on 61 females, they did not feel a lack of protein, but felt a lack of calcium. However, 61 females wanted to consume both protein and calcium actively. 72 The participants knew that even the bones of the fish were edible and tried to eat them, but the purchase price 73 they wanted to buy them at a price far below the actual market price. Therefore, consumers consider fish that 74 can be eaten, even the bones, expensive. As the protein in daily life, it is possible to purchase eggs, beans, and 75 meat every day, but it is not easy to eat fish every day. Fish contain not only high-quality protein, but also large 76 amounts of EPA and DHA1), which are suitable for blood vessels 2-4) and the brain 5). In order to prevent 77 Alzheimer's disease 6,7), it is desirable to take EPA and DHA, which are abundant in fish oil. They are also 78 reported to be effective in preventing neuropathy 8) and leading to health promotion by numerous reports 9,10) 79 Eating fish for calcium ??1) and for protein 12,13) intake is recommended. However, the high price of fish 80 prevents consumers from purchasing fish daily. Even if it is challenging to buy raw fish, is it possible to increase 81 the consumption of fish whose bones are edible by storing it at room temperature in retort pouches and using 82 canned food?

canned food?
Encourage consumers to consume processed fish caught in season (such as canned fish that can store for a long
time and even the bones are edible) to increase protein and calcium intake.

86

V.

87 8 Conclusions

Concerning the consumption of calcium, which did not meet the Japanese dietary intake standards, and the 88 consumption of fish, which is declining a selfadministered questionnaire survey was conducted on 61 females. 89 As a result, the participants felt that their protein intake was sufficient, but felt that their calcium intake was 90 low. The participants answered that they wanted to ingest both protein and calcium actively. The participants 91 knew that there were fish available on the market that could be eaten, including the bones, from which calcium 92 and protein could be ingested simultaneously, but they did not know that the price was higher than the asking 93 price. Some participants ate eggs, meat, and beans daily, but none ate fish daily. In the future, we would like to 94 introduce low-price canned fish and retort-pouch foods to consumers, increase the consumption of fish, even the 95

⁹⁶ bones which can be eaten, and increase the intake of calcium and protein.

1

old	18	19	20	21	22	23	24	25	26
number of participants	3	$\overline{7}$	14	20	$\overline{7}$	2	3	2	2
Table 2. Basic information of 61 participants									
	Average age Average height					eight	Averag	ge wei	
Average \pm Standard diviation		21.	1 ± 2.0		158.4	1 ± 5.9	9		52.9
	Que	stion	items					Yes	No
Do you like fish to eat?								83.6	16.4
Do you know a commercial fish that can eat up to the bones?							75.4	24.6	
Would you like to eat a fish that can be eaten up to the bones of a commercial product?								82.0	18.0
Do you feel that you are deficient in calcium intake on a daily basis?								68.9	29.5
Do you want to take calcium positively?						95.1	3.3		
Do you feel that you are deficient in protein intake on a daily basis?							32.8	67.2	
Do you want to take protein positively?								90.2	9.8

Figure 1: Table 1 .

3

Year 2022 32

Figure 2: Table 3 .

 $\mathbf{4}$

Figure 3: Table 4 .

 $\mathbf{5}$

Questions	50 yen 100 yen 120 yen 150 yer	n 180) yen	200	yen	250 y	yen	280
Suggested purchase price (num- ber)	1	5	1	8	1	16	5	yen 2
Suggested purchase price (%) Questions	1.6 298 yen 300 yen 350 yen 400 ye	-	-	-	-	26.2 n 100	-	
Suggested purchase price (num- ber)	1	7	2	5	4	1	1	1
Suggested purchase price $(\%)$	1.6	11.	53.3	8.2	6.6	1.6	1.6	1.6

2

Figure 4: Table 5 .

8 CONCLUSIONS

97 .1 Acknowledgements

- 98 This study was supported by the Japanese Society of Taste Technology, 2021.
- 99 [Free], Free. Clinical Trial
- [Horrocks et al. (1999)], L A Horrocks, Y K Yeo, L A Horrocks. 10479465. *Pharmacol Res* 1999 Sep. 1999. 40
 (3) p. . (Pharmacol Res.. Review)
- 102 [Larsen et al. (2000)], T Larsen, Thilsted, Hansen. 10743499. Br. J. Nutr 2000 Feb. 83 (2) p. .
- 103 [Swanson et al. (2012)], D Swanson, R Block, S A Mousa, D Swanson. 2012 Jan.
- [Evans et al. (2012)], S J Evans, M Kamali, A R Prossin, G J Harrington, V L Ellingrod, M G Mcinnis, C
 F Burant, S J Evans. 10.1016/j.jpsychires.2012.07.016. J Psychiatr Res 2012 Nov. 2012 Aug. 46 (11) p. .
- 106 [Stark et al. (2016)], K D Stark, M E Van Elswyk, M R Higgins, C A Weatherford, N SalemJr, K D Stark
- 107 . 10.1016/j.plipres.2016.05.001. 27216485. Prog Lipid Res 2016 Jul. 2016 May 20. 2016. 63 p. . (Prog Lipid
 108 Res. Free article. Review)
- [Grimm et al. (2017)], Mow Grimm, D M Michaelson, T Hartmann, Mow Grimm. 10.1194/jlr.R076331.
 28528321. J Lipid Res 2017 Nov. 2017 May 20. 2017. 58 (11) p. . (J Lipid Res. Free PMC article. Review)
- III [Ajith and Ajith ()], T A Ajith, T A Ajith . 10.2174/1574884713666180807145648. 30084334. Curr Clin
 Pharmacol 2018. 2018. 13 (4) p. . (Curr Clin Pharmacol. Review)
- [Arellanes et al. (2020)], I C Arellanes, N Choe, V Solomon, X He, B Kavin, A E Martinez, N Kono, D
 P Buennagel, N Hazra, G Kim, D 'orazio, L M Mccleary, C Sagare, A Zlokovic, B V Hodis, H N
 Mack, W J Chui, H C Harrington, M G Braskie, M N Schneider, L S Yassine, H N Arellanes, IC.
- 116 10.1016/j.ebiom.2020.102883. 32690472. EBioMedicine 2020 Sep. 2020 Jul 17. 2020. 59 p. 102883. (Free PMC article. Clinical Trial)
- 118 [Bianchi et al. (2019)], V E Bianchi, P F Herrera, Laura R Bianchi, VE. 10.1080/1028415X.2019.1681088.
 119 31684843. Nutr Neurosci 2021 Oct. 2019 Nov 4. 2021. 24 (10) p. . (Nutr Neurosci.)
- [Cialdella-Kam et al. (2016)] 'A Mixed Flavonoid-Fish Oil Supplement Induces Immune-Enhancing and Anti Inflammatory Transcriptomic Changes in Adult Obese and Overweight Women-A Randomized Controlled
 Trial'. L Cialdella-Kam , D C Nieman , A M Knab , R A Shanely , M P Meaney , JinF , Sha W Ghosh , S
 Cialdella-Kam , D C Nieman , A M Knab , R A Shanely , M P Meaney , JinF , Sha W Ghosh , S
- Cialdella-Kam , L . 10.3390/nu8050277. 27187. Nutrients 2016 May 11. 2016. 8 (5) p. 277. (Nutrients)
- [A Recent Update on the Effects of Omega-3 Fatty Acids in Alzheimer'sDisease] A Recent Update on the Effects
 of Omega-3 Fatty Acids in Alzheimer'sDisease,
- [Shahidi et al. (2018)] 'Annu Rev Food Sci Technol'. F Shahidi , P Ambigaipalan , F Shahidi . 10.1146/annurev food-111317-095850. 29350557. Annu Rev Food Sci Technol 2018 Mar 25. 2018. 9 p. . (Review)
- 128 [Association of plasma ?-3 and ?-6 lipids with burden of disease measures in bipolar subjects] Association of 129 plasma ?-3 and ?-6 lipids with burden of disease measures in bipolar subjects,
- 130 [Brain delivery of supplemental docosahexaenoic acid (DHA): A randomized placebo-controlled clinical trial]
- Brain delivery of supplemental docosahexaenoic acid (DHA): A randomized placebo-controlled clinical trial,
- [Dietary Animal and Plant Protein and Human Bone Health: A Whole Food Approach Linda K. Massey. American Society for N
 'Dietary Animal and Plant Protein and Human Bone Health: A Whole Food Approach'. Linda K. Massey.
 American Society for Nutritonal Sciences 2003. p. .
- [Effect of nutrition on neurodegenerative diseases. A systematic review] Effect of nutrition on neurodegenerative
 diseases. A systematic review,
- 137 [Emerging Evidence for the Importance of Dietary Protein Source on Glucoregulatory Markers and Type 2 Diabetes: Different Ed
 'Emerging Evidence for the Importance of Dietary Protein Source on Glucoregulatory Markers and Type 2
- Diabetes: Different Effects of Dairy, Meat, Fish, Egg, and Plant Protein Foods. Kevin B. Comerford and
 Gonca Pasin'. Nutrients 2016. 8 p. .
- 141 [Free PMC article ()] 22884424. Free PMC article, 2012.
- 142 [Free PMC article. Review. 2. Global survey of the omega-3 fatty acids, docosahexaenoic acid and eicosapentaenoic acid in the bl
- 143 Free PMC article. Review. 2. Global survey of the omega-3 fatty acids, docosahexaenoic acid and 144 eicosapentaenoic acid in the bloodstream of healthy adults, 10.3945/an.111.000893. 22332096. 2012 Jan 5.
- 145 2012. 3 p. . (Adv Nutr)
- [Health benefits of docosahexaenoic acid (DHA)] Health benefits of docosahexaenoic acid (DHA),
- [Omega-3 fatty acids EPA and DHA: health benefits throughout life] Omega-3 fatty acids EPA and DHA: health
 benefits throughout life,
- 149 [Omega-3 fatty acids, lipids, and apoE lipidation in Alzheimer's disease: a rationale for multi-nutrient dementia prevention]
- 150 Omega-3 fatty acids, lipids, and apoE lipidation in Alzheimer's disease: a rationale for multi-nutrient dementia
- 151 prevention,
- 152 [Omega-3 Polyunsaturated Fatty Acids and Their Health Benefits] Omega-3 Polyunsaturated Fatty Acids and
- 153 Their Health Benefits,