

CrossRef DOI of original article:

Knowledge Evaluation of Accredited Social Health Activists (ASHAs) on Maternal and Child Health, in a Rural Area of Gomati District, Tripura

Dr. Sampa Mitra

Received: 1 January 1970 Accepted: 1 January 1970 Published: 1 January 1970

Abstract

Background: National Rural Health Mission (NRHM) provides a trained female community health worker i.e., Accredited Social Health Activist (ASHA), to every Indian village. An ASHA acts as a bridge between the rural people and the health service outlets. Objective: The objective is to understand the knowledge levels of ASHAs regarding various aspects of maternal and child health. Method: This cross-sectional study, conducted between April 2017 and July 2017, attempts to assess the knowledge levels of 232 ASHAs, working in Ompi CHC (Community Health Centre), Killa PHC (Primary Health Centre) and Atharabula PHC. These centres are located in Ompi RD (Rural Development) Block and Killa RD Block of Gomati District of Tripura. Knowledge level is assessed by performing binomial test at 5

Index terms— ASHA, Gomati District, knowledge level, maternal and child health, binomial test, crosssectional study.

1 Introduction

One of the key features of the National Rural Health Mission (NRHM) is to provide a trained female community health worker i.e., Accredited Social Health Activist (ASHA) to every Indian village (1). An ASHA acts as a bridge between the rural people and the health service outlets, and plays a central role in achieving the national health and population policy goals. An ASHA is expected to provide antenatal, natal and postnatal services to women, give primary medical care (using her kit), assist in the control of diseases with the help of proper information, education and surveillance, counsel people (especially women) on family planning, safe abortion, child immunization, Vitamin A supplementations, appropriate breastfeeding, birth spacing, sex discrimination, child marriage, girls' education, and care of the child (especially newborn), help in household surveys, collaborate with health functionaries working for the community disease control, create awareness on health and its determinants, support the people in utilizing the existing health services etc. (1).

The performance of ASHAs is, therefore, crucial for the success of NRHM.

The objective of this work is to assess the knowledge levels of the ASHAs, with regard to various aspects of maternal and child health, in a rural area of Gomati District, Tripura.

For understanding the knowledge levels of ASHAs, the binomial tests have been done at 5% level of significance.

Before undertaking this study, a brief literature survey has been conducted. However, no study dealing with knowledge evaluation of ASHAs of Gomati District of Tripura, on maternal and child health, has been found.

2 II.

3 Materials and Methods

It is a cross-sectional study, conducted between April 2017 and July 2017, on 232 ASHAs working in Ompi CHC (Community Health Centre), Killa PHC (Primary Health Centre) and Atharabula PHC. These centres (i.e., two PHCs and one CHC) are located in Ompi RD (Rural Development) Block and Killa RD Block; these two blocks

42 have been selected by the simple random sampling method from the eight RD blocks of the Gomati District of
43 Tripura.

44 All ASHAs working in the aforesaid centres, during the study period, who have given informed consent
45 regarding participation in this study, have been included; only those ASHAs who are not willing to participate
46 even after knowing the purpose of the study, have been excluded from the study. It may be noted here that,
47 at the very beginning, the purpose and the objective of the study have been clearly discussed O Knowledge
48 Evaluation of Accredited Social Health Activists (ASHAs), on Maternal and Child Health, in a Rural Area of
49 Gomati District, Tripura with the prospective participants, and they have been given the option to participate
50 or quit at any time.

51 Data have been collected with the help of a pre-designed, pre-tested and semi-structured questionnaire. ASHAs
52 have been visited on ASHA Varosa Divas of each centre, and interviewed face-to-face using the questionnaire. All
53 the collected data have been tabulated, and utilized to perform binomial tests (2), and the results of the tests
54 have been subsequently interpreted.

55 The formula (3,4) for calculating the p-value (p1), pertaining to the binomial test, is given in equation-1: $p1 = \frac{n!}{((n-X)! X!) p^X q^{n-X}}$
56

57 where, n=total number of ASHAs participating in the study=232; X=n/2=116; p=proportion of ASHAs who
58 have given correct answer, in response to a question; q=proportion of ASHAs who have given incorrect answer,
59 in response to the same question. If $p1 < 0.05$, for a question, then it can be inferred that a significant number of
60 ASHAs have correct knowledge with regard to the topic associated with the question; otherwise, the number of
61 ASHAs, having correct knowledge, is not significant.

62 The values of p1 are calculated for all the questions associated with a particular subject, and subsequently
63 averaged (using the formula for arithmetic mean) to get p1 min . If $p1 \text{ min} < 0.05$, for a subject, then it can
64 be inferred that a significant number of ASHAs have correct knowledge with regard to the particular subject;
65 otherwise, the number of ASHAs, having correct knowledge, is not significant.

66 If $p < 0.5$ (and consequently, $q > 0.5$), then the value of p1 may generate erroneous inference regarding
67 significance. Hence, in these cases (where $p < 0.5$ and $q > 0.5$), p and q are both taken as 0.5 only for the sake of
68 calculating p1.

69 Since the data have been collected for purely academic purpose, and the permission for data collection has
70 been obtained from the CMOH (Chief Medical Officer of Health) of the Gomati District, and also the MOICs
71 (Medical Officers in Charge) of the respective centres, there is no chance of any kind of exploitation involved in
72 this study.

73 The method used in this work, is shown in fig. -1.

74 4 Results

75 The knowledge levels of ASHAs with regard to antenatal care, postnatal care, breastfeeding, child health,
76 immunization and family planning, are shown respectively in tables-1-5. At the end of each table, the percentage
77 for the significant number of ASHAs has been calculated by averaging the percentages corresponding to the
78 number of correct answers, in the corresponding table.

79 The values of p1 in tables-1-5 show that a significant number of ASHAs have correct knowledge regarding the
80 topic associated with each of the questions, except question number-8 of table-1, question number-2 of table-2,
81 and question numbers-4 and 6 of table-5.

82 IV.

83 5 Discussions

84 From tables-1-5, the following inferences can be drawn: aspects of all of the aforesaid subjects (viz., antenatal
85 care, postnatal care, breastfeeding, child health, immunization and family planning). But, the situation is quite
86 far from the reality. Hence, there is no room for complacency, and immediate arrangements should be made for
87 proper and adequate training of the ASHAs, particularly with regard to the four aforesaid topics, pertaining to
88 each of which, a significant number of ASHAs do not have correct knowledge. Now, a brief review of the outcomes
89 of some of the other studies on ASHAs, may help in understanding the relevance of this work.? A significant

90 A study undertaken by ??arg et al. shows that in some villages of rural Haryana, majority of ASHAs knew
91 about assisting in immunization (100%), accompanying women for delivery (98%), and providing antenatal care
92 (96.10%) and family planning (96.40%) services, as a part of their duty (5). In the current study also, it has been
93 found that a significant number of ASHAs have thorough knowledge about antenatal care (78.9%), immunization
94 and family planning (66.5%).

95 Fathima et al. conducted a survey on ASHAs of some parts of Karnataka, and found that a significant number
96 of ASHAs (>80%) were involved in home visits, and counselling with regard to antenatal care, breastfeeding
97 and immunization, but only a small proportion (<25%) of ASHAs gave advice on the use of contraceptives (6).
98 These observations, except the one concerning contraceptives, are similar to those of the current study, where
99 it has been found that a considerable number of ASHAs have good knowledge about antenatal care (78.9%),
100 breastfeeding (88.2%), immunization and family planning (66.5%). However, in the present study, 55.6% of the

101 ASHAs know what a condom (nirodh) is (though only 39.7% of them know what an oral contraceptive pill (OCP)
102 is).

103 Guha et al. worked on the ASHAs of some villages of Maharashtra (Wardha), and observed that though
104 majority of the ASHAs undertook home visits and assisted in antenatal care and postnatal care, only a few of
105 them were aware of their roles in facilitating breastfeeding and the use of contraceptives (7). Regarding antenatal
106 care and postnatal care, their findings are similar to those of the present study which indicates good knowledge
107 among significant number of ASHAs about antenatal care (78.9%) and postnatal care (69.8%). But, unlike their
108 observations about breastfeeding and contraceptives, the present study indicates that a considerable number of
109 ASHAs are knowledgeable about breastfeeding (88.2%) and condom (nirodh) (55.6%). (However, according to
110 the current study, only 39.7% of the ASHAs know what an oral contraceptive pill (OCP) is.)

111 Taksande et al. studied the ASHAs of selected villages of Maharashtra (Wardha), and found that 48% of
112 them were aware of the antenatal care registration in clinic (8). This outcome is not quite in agreement with the
113 finding of the current study that 78.9% of the ASHAs have good knowledge about antenatal care.

114 Panda et al. undertook a study on ASHAs working in some parts of Odisha, and found that 100% of the
115 ASHAs helped in immunization, and 98% of them knew about family planning activities (9). Their observations
116 are similar to that of the current study that majority of the ASHAs (66.5%) have correct knowledge about
117 immunization and family planning.

118 Sugandha et al., after conducting a study on the ASHAs of Mysuru (Karnataka), noted that 51.5% of them
119 had average knowledge about antenatal care, 86.1% had good knowledge regarding postnatal care, and 90.5%
120 had thorough knowledge about contraception (10). With regard to postnatal care and contraception, there are
121 similarities between their outcomes and those of the current study, since the present study indicates correct
122 knowledge, among significant number of ASHAs, in the fields of postnatal care (69.8%), immunization and
123 family planning (66.5%). However, in contrast to their observation For table-5, p1 min =0.033. Hence, it may
124 be concluded that a significant number of ASHAs (66.5%) have correct knowledge regarding immunization and
125 family planning. about antenatal care, the current study shows that a considerable number of ASHAs (78.9%)
126 have good knowledge about antenatal care.

127 Chaurasiya et al., after analyzing relevant information regarding the services provided by the ASHAs, collected
128 from their beneficiaries (i.e., mothers having child up to 2 years of age), in some villages of a district of western
129 Uttar Pradesh, observed that ASHAs escorted 43.5% of the beneficiaries ?4 times to the hospital/health centre
130 for antenatal care visit, and performed ?6 home visits for postnatal care for 57.2% of the beneficiaries (11). These
131 observations indicate that majority of the ASHAs have good knowledge about postnatal care, but not antenatal
132 care. However, in the present study, significant number of ASHAs have been found to have thorough knowledge
133 regarding both antenatal care (78.9%) and postnatal care (69.8%).

134 Bhattacharya et al. worked on how the pregnant women utilized services of the ASHAs in some villages of
135 the West Tripura District of Tripura, and found that the ASHAs helped 76.69% of the women to have adequate
136 antenatal check-ups (12). This finding indicates that majority of the ASHAs have good knowledge about antenatal
137 care, and hence, is similar to the outcome of the current study that 78.9% of the ASHAs have thorough knowledge
138 about antenatal care.

139 V.

140 6 Conclusions

141 The work has attempted to assess the knowledge levels of the ASHAs, with regard to various aspects of maternal
142 and child health, in a rural area of Gomati District, Tripura. On the basis of the results of the binomial tests, it can
143 be concluded that a significant number of ASHAs, associated with Ompi CHC, Killa PHC and Atharabula PHC,
144 have correct knowledge regarding various aspects of the subjects of antenatal care, postnatal care, breastfeeding,
145 child health, immunization and family planning, except certain topics viz., heavy manual labour by women
146 during pregnancy, minimum number of postnatal check-ups, time-limit of abortion after pregnancy, and oral
147 contraceptive pill (OCP). However, since all ASHAs need to become aware of all the relevant details with respect
148 to the aforesaid subjects, there is an urgent need for proper and adequate training of the ASHAs, especially with
149 regard to the four abovementioned topics.

150 A more comprehensive idea about the ASHAs of Tripura could have been had, if data were collected from other
151 districts also. Besides, a more detailed questionnaire might have been helpful in judging the ASHAs' knowledge
152 levels regarding various aspects of maternal and child health, to a greater degree. If possible, these tasks may be
153 taken up in future.



Figure 1: Fig.- 1 :




Figure 2:

6 CONCLUSIONS

Serial Number	Question	Number of Correct Answers	Number of Incorrect Answers	p
1.	What is the optimal time for registration of pregnant women?	222 (95.7%)	10 (4.3%)	0
2.	What is the minimum number of ANC check-ups required?	220 (94.8%)	12 (5.2%)	0
3.	Is early pregnancy diagnosis to be done by testing urine, stool or blood?	181 (78.0%)	51 (22.0%)	1.2
4.	What is the purpose of Nischay kit?	198 (85.3%)	34 (14.7%)	1.3
5.	How many times should the pregnant women (primi) be given tetanus toxoid (TT) injection?	210 (90.5%)	22 (9.5%)	0
6.	What is the implication if after blood test, haemoglobin concentration level is found to be less than 7.0 gm/dl?	168 (72.4%)	64 (27.6%)	5.1
7.	Is there any necessity to give women mo food during pregnancy?	219 (94.4%)	13 (5.6%)	0
8.	Should women carry out heavy manual labour during pregnancy?	102 (44.0%)	130 (56.0%)	0
9.	What is the minimum period for which all pregnant women must take IFA (iron and folic acid) tablet (1 tablet daily) to prevent anaemia?	116 (50.0%) (for calculating p1)	116 (50.0%) (for calculating p1)	1.3
10.	What are the side effects of IFA tablet?	139 (59.9%)	93 (40.1%)	-3
11.	What are the potential danger signs during pregnancy which prompt immediate referral?	151 (65.1%)	81 (34.9%)	1.3
12.	What are the potential danger signs during pregnancy which prompt immediate referral?	196 (84.5%)	36 (15.5%)	2.3
13.	Is it the duty of an ASHA to escort the pregnant women to the hospital for institutional delivery?	229 (98.7%)	3 (1.3%)	0
14.	What should an ASHA do if, after home delivery, mother reports excessive vaginal bleeding or severe abdominal pain?	151 (65.1%)	81 (34.9%)	1.3
14.	What is the meaning of birth-preparedness?	178 (76.7%)	54 (23.3%)	1.3

For table-1, $p_1 \text{ min} = 0.008$. This implies that a significant number of ASHAs (78.9%) have correct knowledge to antenatal care

Serial Question Number	Question	Number of Correct Answers	Number of Incorrect Answers	p1
1.	What is the meaning of postnatal period?	138 (59.5%)	94 (40.5%)	1.4699X10 ⁻³
2.	What is the minimum number of postnatal check-ups required?	99 (42.7%)	133 (57.3%)	0.1046
3.	What are the common major complications which can happen during the postpartum period?	116 (50.0%) (for calculating p1)	116 (50.0%) (for calculating p1)	5.3545X10 ⁻³¹
4.	What should be the minimum spacing between two live child births?	213 (91.8%)	19 (8.2%)	0
5.	Which women are more at the risk of developing complications during delivery?	158 (68.1%)	74 (31.9%)	8.7765X10 ⁻¹⁴
6.	What is the time interval following delivery, after which the placenta usually comes out?	171 (73.7%)	61 (26.3%)	1.5836X10 ⁻¹⁴

Table-3: Knowledge of ASHAs regarding breastfeeding (n=232)

Serial Question Number	Question	Number of Correct Answers	Number of Incorrect Answers	p1
1.	Why is colostrum necessary for a newborn baby?	206 (88.8%)	26 (11.2%)	0
2.	When should a newborn be put to breast, after birth?	221 (95.3%)	11 (4.7%)	0
3.	At least how many times should a newborn be breastfed per day?	195 (84.1%)	37 (15.9%)	3.1401X10 ⁻³³
4.	What are the indications of inadequate breastfeeding to a baby?	170 (73.3%)	62 (26.7%)	4.8518X10 ⁻¹⁴
5.	What should be the duration of exclusive breastfeeding, according to WHO (World Health Organization)?	210 (90.5%)	22 (9.5%)	0
6.	What is the age at which, complementary food is necessary for a baby?	225 (97.0%)	7 (3.0%)	0

[Note: For]

Figure 4: Table - 2

6 CONCLUSIONS

-

Serial Number	Question	Number of Correct Answers	Number of Incorrect Answers	p1
1.	What is the minimum body weight of a normal baby at birth?	226 (97.4%)	6 (2.6%)	0
2.	When should a baby be considered to have low birthweight (LBW)?	128 (55.2%)	104 (44.8%)	0.02
3.	What are the signs of a high risk baby?	148 (63.8%)	84 (36.2%)	1.06 -5
4.	What are the signs/symptoms due to which, an ASHA should advise the mother/family to take a baby to the hospital immediately?	182 (78.4%)	50 (21.6%)	2.48 -21
5.	What are the signs/symptoms which indicate severe dehydration of a baby in case of diarrhoeal disease?	183 (78.9%)	49 (21.1%)	3.43 -22
6.	Should a baby be given bath immediately after birth?	203 (87.5%)	29 (12.5%)	0
7.	When can the body temperature of a newborn baby be considered as less than normal?	190 (81.9%)	42 (18.1%)	4.89 -28
8.	How many times should an ASHA undertake home visit, in case of institutional delivery?	186 (80.2%)	46 (19.8%)	1.43 -24
9.	How many times should an ASHA undertake home visit, in case of home delivery?	205 (88.4%)	27 (11.6%)	0

Serial Number	Question	Number of Correct Answers	Number of Incorrect Answers	p1
1.	What are the disease/diseases against which BCG (Bacillus Calmette-Guérin) vaccine provides protection?	225 (97.0%)	7 (3.0%)	0
2.	What are the disease/diseases against which DPT* vaccine provides protection?	192 (82.8%)	40 (17.2%)	4.69 -30
3.	What is the minimum interval between two doses of Vitamin A?	182 (78.4%)	50 (21.6%)	2.48 -21
4.	Up to what time interval after pregnancy, is abortion legal?	108 (46.6%)	124 (53.4%)	0.10
5.	What is a condom (nirodh)?	116 (50.0%) (for calculating p1)	116 (50.0%) (for calculating p1)	0.02
6.	What is an oral contraceptive pill (OCP)?	129 (55.6%)	103 (44.4%)	0.02
8.	What is an oral contraceptive pill (OCP)?	92 (39.7%)	140 (60.3%)	0.10

154 .1 Acknowledgements

155 The author is deeply indebted to Dr. Shantanu Jamatia, PGDPHM student (session: 2016-17), AIIH & PH,
156 Kolkata, for collecting the data and helping in the analysis.

157 Besides, the author is grateful to the ASHAs (who have participated in this study) associated with Ompi
158 CHC, Killa PHC and Atharabula PHC, for their cooperation in the data collection process.

159 .2 Source of Funding: Personal Source

160 .3 Conflict of Interest:

161 The author certifies that there is no conflict of interest, involved in this study.

162 [National Health Mission. About Accredited Social Health Activist] , [https://nhm.gov.in/index1.php?](https://nhm.gov.in/index1.php?lang=1&level=1&sublinkid=150&lid=226)
163 [lang=1&level=1&sublinkid=150&lid=226](https://nhm.gov.in/index1.php?lang=1&level=1&sublinkid=150&lid=226) *National Health Mission. About Accredited Social Health*
164 *Activist (ASHA)* . (Internet)

165 [Van Den Berg] , R G Van Den Berg . (Binomial Test -Simple Tutorial) (Internet)

166 [Mitra and Chakraborty ()] ‘A Study on the Socio-Demographic and the Clinical Profiles, Along with Some
167 Management Indicators, of the MDR TB Patients at the Domjur Block’. S Mitra , B Chakraborty . *J*
168 *Bacteriol Infect Dis* 2020. (Internet)

169 [Panda et al. ()] ‘A Study on the Work Profile of ASHA Workers in a District of Odisha in Eastern India’. M
170 Panda , S Nanda , R C Giri . *Int J Community Med Public Heal* 2019. 6 (2) p. .

171 [Garg et al. ()] ‘An Evaluation of ASHA Worker’s Awareness and Practice of Their Responsibilities in Rural
172 Haryana’. P K Garg , A Bhardwaj , A Singh , S K Ahluwalia . *Natl J Community Med* 2013. 4 (1) p. .

173 [Fathima et al. ()] ‘Assessment of ‘Accredited Social Health Activists’-A National Community Health Volunteer
174 Scheme in Karnataka State’. F N Fathima , M Raju , K S Varadharajan , A Krishnamurthy , S R Anathkumar
175 , P K Mony . *India. J Heal Popul Nutr* 2015. 33 (1) p. .

176 [Chaurasiya et al. ()] ‘Assessment of the Services of ASHA Workers on Antenatal and Postnatal Care in a District
177 of Western Uttar Pradesh’. S K Chaurasiya , N P Singh , S K Shukla , P K Bajpai , D J Mathew . *India. J*
178 *Fam Med Prim Care* 2020. 9 (7) p. .

179 [Hypothesis Testing with the Binomial Distribution] *Hypothesis Testing with the Binomial Distribution*, New-
180 castle University (Internet)

181 [Sugandha and Jagannath ()] ‘Knowledge of ASHA Workers About Maternal and Child Health Services in
182 Mysuru’. B K Sugandha , P Jagannath . *Public Heal Rev -Int J Public Heal Res* 2019. 6 (4) p. .

183 [Guha et al. ()] ‘Qualitative Assessment of Accredited Social Health Activists (ASHA) Regarding Their Roles
184 and Responsibilities and Factors Influencing Their Performance in Selected Villages of Wardha’. I Guha , A
185 V Raut , C H Maliye , A M Mehendale , B S Garg . *Int J Adv Med Heal Res* 2018. 5 (1) p. .

186 [Taksande et al. ()] ‘To Assess the Job-Related Difficulties and Dissatisfaction of Asha Workers in Selected Rural
187 Area’. V D Taksande , P A Ashtankar , C R Bansod , A V Bawane , P S Burchunde , D V Dudhe . *J Evol*
188 *Med Dent Sci* 2021. 10 (2) p. .

189 [Bhattacharya et al. ()] ‘Utilization of ASHA Services by the Pregnant Women of Rural Tripura, India’. H
190 Bhattacharya , N Luwang , M Sarkar , T Chakraborty , S Baidya . *Int J Res Med Sci* 2015. 3 (9) p.
191 .