



GLOBAL JOURNAL OF MEDICAL RESEARCH: K

INTERDISCIPLINARY

Volume 22 Issue 6 Version 1.0 Year 2022

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals

Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Salt Profile and Content in Foods Prepared in Restaurants and Bakeries: Analysis of the 2 Main Urban Centers in Cape Verde

By Edna Duarte Lopes, Júlio Monteiro Rodrigues, Maria da Luz Lima, Alzerina Maria Rocha Monteiro, Irina Maia Spencer, Janice Soares & Ailton Luís Lopes Ribeiro

Abstract- The excessive intake of sodium from salt is associated with the risk of cardiovascular diseases. According to the Noncommunicable Diseases Survey (DNS, N/P), more than 35% of the Cape Verdean population is considered hypertensive. There are no data on the salt content of foods prepared by restaurants and bakeries in Cape Verde. Most restaurants and bakeries use iodized salt (97%) of national origin (81%), from the islands of Sal (45.6%) and Maio (20.9%). Approximately 53.3% of the bread samples had salt contents greater than 1.4 g/100 g. More than half (59.3%) of ready-to-eat meals collected from restaurants in Praia and 42.4% in Mindelo had a salt content equal or more than 5.0 g/meal. Restaurants and bakeries in the cities of Praia and Mindelo use salt in a nonstandardized manner. The results of this study may contribute to the redesign of salt intake reduction strategies in Cape Verde.

Keywords: salt (NaCl) profile, salt (NaCl) content, restaurants and bakeries.

GJMR-K Classification: DDC Code: 647.9544361 LCC Code: TX907.5.F72



SALT PROFILE AND CONTENT IN FOODS PREPARED IN RESTAURANTS AND BAKERIES ANALYSIS OF THE 2 MAIN URBAN CENTERS IN CAPE VERDE

Strictly as per the compliance and regulations of:



RESEARCH | DIVERSITY | ETHICS

Salt Profile and Content in Foods Prepared in Restaurants and Bakeries: Analysis of the 2 Main Urban Centers in Cape Verde

Edna Duarte Lopes ^a, Júlio Monteiro Rodrigues ^a, Maria da Luz Lima ^b, Alzerina Maria Rocha Monteiro ^c, Irina Maia Spencer ^d, Janice Soares ^d & Ailton Luís Lopes Ribeiro ^x

Abstract- The excessive intake of sodium from salt is associated with the risk of cardiovascular diseases. According to the Noncommunicable Diseases Survey (DNS, N/P), more than 35% of the Cape Verdean population is considered hypertensive. There are no data on the salt content of foods prepared by restaurants and bakeries in Cape Verde. Most restaurants and bakeries use iodized salt (97%) of national origin (81%), from the islands of Sal (45.6%) and Maio (20.9%). Approximately 53.3% of the bread samples had salt contents greater than 1.4 g/100 g. More than half (59.3%) of ready-to-eat meals collected from restaurants in Praia and 42.4% in Mindelo had a salt content equal or more than 5.0 g/meal. Restaurants and bakeries in the cities of Praia and Mindelo use salt in a nonstandardized manner. The results of this study may contribute to the redesign of salt intake reduction strategies in Cape Verde.

Keywords: salt (NaCl) profile, salt (NaCl) content, restaurants and bakeries.

I. INTRODUCTION

Noncommunicable diseases (NCDs) are responsible for the death of 41 million people annually, equivalent to 71% of all deaths worldwide, with cardiovascular diseases being the main cause (17.9 million) (World Health Organization, Noncommunicable diseases, 2021). In turn, hypertension is the leading cause of cardiovascular disease and premature death worldwide (World Health Organization, Hypertension, 2021).

A medium-developed country, Cape Verde is located in the Atlantic Ocean, between the Equator and the Tropic of Cancer, between latitudes 14° 23' and 17° 12' north latitude and between 22° 40' and 25° 22' west longitude ("Portal informativo de Cabo Verde,"). The country imports the majority (>80%) of the food it consumes (ALMEIDA, 2022). In Cape Verde, due to changes in the demographic, epidemiological and nutritional profiles of the Cape Verdean population,

chronic NCDs are the leading cause of mortality in the country (MS, N/P). According to preliminary data from the 2019 Survey on Noncommunicable Diseases and Associated Risk Factors (IDNT II, acronym in Portuguese), 30.8% of the Cape Verdean population is hypertensive.

According to the World Health Organization (WHO), high blood pressure is responsible for at least 45% of deaths from heart disease and 51% of deaths from stroke (World Health Organization, Hypertension, 2021). The worldwide prevalence of high blood pressure (systolic and/or diastolic blood pressure $\geq 140/90$ mmHg) in adults (age ≥ 18 years) is estimated at 22% (Mills, Stefanescu, & He, 2020; World Health Organization, Hypertension, 2021). Data from the 2019 IDNT II indicate that cardiovascular diseases and cancer are leading causes of deaths in Cape Verde, highlighting poor eating habits among the most common risk factors (MS, N/P).

When poor eating habits are identified as one of the risk factors for circulatory system diseases and cancer, among other conditions, substances such as salt (Pure salt - sodium chloride, contains 39.34% sodium or 39.340 mg per 100g) and other additives such as preservatives used in canned foods, meat, fish, etc., are often mentioned. However, the identity and nature of these substances, as well as their effects on the body, are rarely explained (MOLOGNONI, 2019). In developed countries, up to three-quarters of the total salt intake comes from foods consumed outside the home and from processed foods, in contrast, the salt added to food, both at the table and during home preparation, generally represents a lower proportion of the total sodium intake (Coelho, 2021; Ruiz et al., 2020).

Sodium is an essential nutrient necessary for the normal functioning of cells. Its excessive intake is associated with adverse health conditions, particularly hypertension, which in turn is considered one of the main risk factors for some chronic diseases with high morbidity and mortality worldwide (National Academies of Sciences et al., 2019; World Health Organization, 2012; World Health Organization, 2020). According to Nascimento et al. (Nascimento, Gavron, Bowles, Chaves, & Bortolozo, 2017), sodium (Na) intake comes

Author a o p §: Department of Research, Science and Training, National Institute of Public Health, Chá de Areia C.P. nº 719, Praia - Cape Verde.

Author c: Praia Health Precinct Affiliation, Praia - Cape Verde.

Author d: National Directorate of Health, Praia - Cape Verde.

Author x: Department of Research, Science and Training, National Institute of Public Health, Chá de Areia C.P. nº 719, Praia - Cape Verde.
e-mail: ailton.ribeiro@insp.gov.cv



mainly from table salt, salt-based foods and industrialized products. Complications are related to sodium intake in amounts above that recommended by the WHO of a maximum of 2 g/day (World Health Organization, 2020), considering all its forms of presentation.

Although the 2019 IDNT II (MS, N/P) indicates that close to one-third of the Cape Verdean population is hypertensive, there are no data on the salt profile and content of foods prepared by restaurants and bakeries in Cape Verde. Thus, the main objective of this study was to evaluate the profile and content of salt in foods prepared in restaurants and bakeries in the cities of Praia and Mindelo in Cape Verde.

II. MATERIALS AND METHODS

a) Study design and population

This was an observational, cross-sectional study with a qualitative-quantitative approach conducted in restaurants and bakeries in the cities of Praia and Mindelo. These cities have the largest populations in the country (Instituto Nacional de Estatística de Cabo Verde, 2020). All restaurants and bakeries in the 2 cities, based on georeferencing data provided by the Independent Health Regulatory Entity (ERIS, acronym in Portuguese), Municipal Councils and the General Directorate of Tourism, were selected for the study, and all representatives who agreed to participate did so voluntarily. In cases where the georeferenced establishment was already closed, the data were collected from the nearest unreference establishment. Written informed consent was provided by the representative of each establishment prior to data collection.

Inclusion criteria were being on the official list provided by the competent entity or the existence of the establishment being known to field agents even though it was not on the official list, and the person responsible voluntarily accepting to participate in the study, by signing the free and informed consent form.

As exclusion criteria, the establishment is not on the official list and its existence is not known to field agents, or the person in charge does not accept to participate in the study, by signing the free and informed consent form. The heading levels should not be more than 4 levels. The fond of heading and subheadings should be 12 point normal.

b) Participant characteristics

A total of 155 managers of restaurants and bakeries in the cities of Praia and Mindelo participated in the study, most of whom (61.4%) were female. Most subjects were aged 25-44 years (58.2%). The mean age of the respondents was 32.41 ± 8.41 years. Regarding education level, 48.1% completed secondary education, 28.5% completed primary education, 15.2% completed undergraduate education, 2.5% completed vocational

training, 1.9% had a master's degree, and 1.3% were illiterate.

c) Data collection

i. Questionnaires and food samples

A protocol cover sheet containing the identification data of the establishment and 2 adapted self-report questionnaires (Appendix I and II) (Viegas CACL, 2013) were applied to record observations on the origin, transport, storage and use of salt used in the establishments under study.

The data were collected by 11 investigators under the supervision of 2 supervisors; the investigators were distributed between the cities of Praia (6 investigators; 1 supervisor) and Mindelo (5 investigators; 1 supervisor). The investigators underwent 1 week of training on techniques for questionnaire application and techniques for collecting, packaging and transporting food samples, on the ethical issues of the research, and on the analysis and interpretation of salt iodization data based on the instructions of the Iodized Salt Field Test Kit.

The food samples were collected in dry sterile polyethylene bags to avoid any possible contamination and were transported in thermal bags containing cold packs in accordance with the best practices for food transport (Associação de restauração e similares de Portugal, 2008). In the laboratory, the samples were stored at -20°C until analysis.

ii. Rapid salt iodization test

In each establishment, a small amount of salt (approximately one teaspoon) was collected on a white paper sheet, to which 1-2 drops of the test solution was added. Using a color table provided in the test kit, the result was recorded immediately under good lighting. For cases where no color was observed after 1 minute (suspicion of alkalinity in the salt sample), 5 drops of verification solution was added to a new salt sample, and 2 drops of the test solution was added to the sample (World Health Organization, 2007; Jooste PL & Strydom E, 2010).

16, 17). Last, using the color table, the result was recorded and transcribed to the questionnaire sheet.

d) Laboratory analysis

i. Sample preparation

Bread samples were ground in a blender immediately after their removal from the freezer, and the portion needed for salt content analysis was separated and set aside.

Samples of meals were slowly thawed, i.e., kept in a refrigerator at $\sim 4^{\circ}\text{C}$ for a period of 24 hours. Then, the samples were homogenized in a blender, and the necessary portion was removed for analysis.

ii. *Analysis of sodium chloride content*

The sodium chloride content was evaluated through titration by the Mohr method. After sample preparation, approximately 5 g of Portuguese bread/wheat bread and 10 g of meals were weighed in duplicate on an analytical inside 250-ml cups. Then, 100 ml of warm distilled water (50-55 °C) and 100 ml of boiling distilled water (~100 °C) were added to the cups containing bread and meal samples, respectively (Ward, R. E., & Carpenter, C. E., 2010).

The solution was vigorously homogenized (30 s) twice at an interval of 1 minute. After the solution had cooled to room temperature, the pH was adjusted to 6.5-10.0, and the solution was filtered through sterile gauze. Next, 15 ml of the filtered solution was transferred to a 250-ml Erlenmeyer flask, and 1.5 ml of 0.1 M potassium chromate indicator (K₂CrO₄) was added. The solution was then titrated with 0.1 M silver nitrate (AgNO₃) until the first appearance of a brownish red color; the process was continued for another 30 s, and the volume of titrant used was recorded.

iii. *Calculations*

The sodium chloride content in each replicate sample was calculated, and the mean, median, range and standard deviation were calculated.

e) *Statistical analysis*

The data were analyzed using the Statistical Package for the Social Sciences (SPSS, v. 26). The data were described using absolute and relative frequencies, and the chi-square test was used to compare the salt content in food prepared by the establishments in Praia and Mindelo. All tests were two-sided, and P values less than 0.05 were considered to be statistically significant.

f) *Ethics approval of research*

This study was approved by the National Committee on Ethics in Health Research (deliberation no. 33/2019, 31 May). All participants signed an informed consent form, which was filed under confidentiality at the National Institute of Public Health of Cape Verde. Written informed consent for participation was obtained from managers of restaurants and bakeries and their privacy and confidentiality were maintained. All personal identifiers were excluded, and data was kept confidential and used for the proposed study only.

i. *Guideline statement*

Resolution No. 33/2019

1. It was submitted, under the terms of article 9 of Decree-Law No. 26/2007, of 30 July, for the purposes of the National Committee on Ethics in Health Research (CNEPS, acronym in Portuguese) opinion in order to authorize the realization of the Project entitled "Salt profile and content in foods prepared in restaurants and bakeries: Analysis of the 2 main urban centers in Cape Verde", presented

by the National Institute of Public Health, INSP, with Dr Júlio Monteiro Rodrigues as the main researcher, Executive Director of the aforementioned Institute, which has a partnership with the Institute of Hygiene and Tropical Medicine, Lisbon, Portugal.

2. The research project team is comprised not only of technicians responsible for the INSP, but also of the Coordinator of the National Nutrition Program, the National Directorate of Health and technicians from the Praia Health Precinct, among others.
3. This is an observational, cross-sectional study with quasi-quantitative approaches.
4. The study will take place in the cities of Praia and Mindelo and will cover a representative sample of restaurants and bakeries, based on georeferencing, provided by the General Directorate of Tourism and/or the National Statistics Institute, INE.
5. Only duly registered restaurants and bakeries whose representatives accept to participate in the study will be considered.
6. As supporting documents, were presented, (i) the Letter addressed to CNEPS, (ii) the Research Project, (iii) the Schedule, (iv) the Indicative Budget, (v) the Information Sheets, (vi) the Terms of Free and Informed Consent, (vii) the Questionnaires to be applied and, (viii) the CV of the principal investigator?
7. CNEPS performed the document analysis of the research project during its 92nd Ordinary Meeting, held on April 25, 2019, and the project was pending additional information (see deliberation 28/2019).
8. On May 15, 2019, the research project with the corrections was submitted for reconsideration by the CNEPS.
9. CNEPS reassessed the research project during its 92nd Ordinary Meeting, held on May 30, 2019, having found that the aspects highlighted in the aforementioned determination were remedied, therefore, pursuant to article 11 of Decree-Law no. 26/2007, of 30 July, it decided to approve it.

III. RESULTS

a) *Salt used in restaurants*

A total of 125 restaurants were evaluated. Regarding the origin of the salt used during food preparation in the restaurants, 79% of restaurants used salt of national origin, 13.7% used salt of international origin, 4% used salt of both origins, and 3.2% used salt of an unknown origin.

Of the restaurants in which the salt used in food preparation was of national origin, 62.6% used salt from the island of Sal, and 25.3% used salt from the island of Maio; for 12.1%, the island of origin was unknown. Notably, the island of Santiago does not produce salt.

Of the restaurants in which the salt used in food preparation was of international origin, 95% used salt from Portugal, and 5% used salt from France.



Regarding the means used to transport salt, 61.1% of restaurants used a car, 18.2% used a car and boat, 7.1% used a car and plane, and 13.5% did not use any means of transport.

Most restaurants (58.4%) did not have a specific location for salt storage. The salt was stored in its original packaging (91.9%), protected from light (38.7%), exposed to heat in a nonventilated area (19.4%), stored in the presence of chemicals (9.7%), and exposed to moisture (6.5%).

Other forms of salt storage indicated by the respondents were in plastic buckets (30%), together with other nonperishable foods (25%), in flasks (15%), in stainless steel containers (10%), in plastic bowls (10%), inside a kitchen cabinet (5%) and in a bag inside a box (5%).

Among the restaurants that did not have a specific location for salt storage, in 79.6%, the salt was stored in the kitchen; in 12.2%, the salt was stored in the pantry; in 4.1%, the salt was stored on a kitchen shelf; and in 2%, the salt was stored on a shelf or near the stove.

Regarding the lighting where the salt was stored, the location was well lit in 30.6% and lit in 26.5% of restaurants, received natural light in 22.4%, low light in 12.2%, and no light in 4.1% of restaurants, and was exposed to light and heat in 2% of restaurants.

The salt storage location was ventilated in 44.7% of the restaurants, well ventilated in 23.4%, poorly ventilated in 12.8%, ventilated and exposed to heat in 6.4%, not ventilated in 4.3%, exposed to heat in 4.3% and exposed to natural ventilation in 2.1%.

In 29.2% of the restaurants, the salt was stored in its own packaging; in 37.5%, the salt was stored in plastic jars; in 12.5%, the salt was stored in a container with a lid; in 6.3%, the salt was stored in plastic bags; in 6.3%, the salt was stored in jars; in 4.2%, the salt was stored in a random container; and in 4.2%, the salt was stored in glass jars. In 82.1% of the restaurants, the salt was stored together with other products.

Most restaurants (83.3%) used a utensil for handling salt: spoon (69.8%), cup (4%), ladle (4%), stainless steel scoop (1.6%), spoon and jar (1.6%), milk pitcher (0.8%), jar (0.8%) and wooden spoon (0.8%).

b) Salt used in bakeries

A total of 30 bakeries were evaluated. Most of the salt used in the preparation of bread was of national origin, with 55.6% of bakeries using salt from the island of Sal and 37% using salt from the island of Maio; 7.4% reported not knowing the provenance.

The car was the most commonly used means for transporting salt (76.7%), followed by boat and car (20.0%) and plane and car (3.3%). Most bakeries purchased salt locally (76.7%) and stored it in a specific location (80%). Of these, 78.3% stored the salt in its original container, 47.8% stored the salt protected from light, 30.4% stored the salt in a location exposed to heat, 4.3% stored the salt in a nonventilated area, and 4.3% stored the salt in the presence of chemicals.

Of the 6 bakeries that did not have a specific location for salt storage, 50% stored it in the pantry, 33.3% stored it in the production area, and 16.7% stored it in a location with other products. In 66.7% of the bakeries, the salt was stored away from light; in 50%, the salt was stored in its own container; in 33.3%; the salt was exposed to heat; and in 16.7%; the salt was stored in an unventilated location.

A total of 96.4% of the bakeries used a utensil to handle the salt: cup (25.9%), ladle (25.9%), stainless steel scoop (14.8%), bowl (7.4%), shaker (7.4%), and either a spoon, scale plate, spatula, cake pan or jar (3.7% each).

c) Profile of the salt used in restaurants

Only 12% of the restaurant managers stated that they followed standard guidelines for the amount of salt to be used in food preparation.

Most (67.2%) restaurant managers believe that there are differences between different types of salt, whereas 30.4% think there are no differences and 2.4% do not know. Of those who reported the existence of differences between the types of salt, 59.5% referred to the quality, with emphasis on the presence of iodine (30%).

Table 1 shows the proportion of salt per meal used by restaurants in Praia and Mindelo. In both the soup and the main dish, most establishments used less than 1 g of salt.

Table 1: Amount of salt/meal used in restaurants

Salt	Restaurants			
	Soup		Main dish	
	n	%	n	%
< 1 g	75	59.5	65	51.6
1 to 2 g	25	19.8	40	31.7
2 to 4 g	7	5.6	14	11.1
4 to 6 g	1	0.8	2	1.6
> 6 g	-	-	5	4
Does not cook	17	13.5	-	-
Does not know	1	0.8	-	-
Total	126	100	126	100

Regarding the daily dose of salt recommended by the WHO, 48% of respondents were unaware of the existence of a WHO recommendation on the amount of salt that should be consumed daily per person; 18.4% were aware of such a recommendation. Only 37.3% of the respondents knew the relationship between salt and sodium.

d) Profile of the salt used in bakeries

Regarding the existence of a standard that regulates the amount of salt to be used in the preparation of bread, 83.3% of the bakeries reported that they followed a standard; for the remaining 16.7%, there was no compliance with any standard.

Iodized salt was the most used salt type (90%), followed by sea salt, fleur de sel and others, with 3.3% each.

Table 2: Amount of salt per kg of dough and per loaf

Salt	kg of dough		Loaf	
	n	%	n	%
< 1 g	4	14.3	21	75
1 to 2 g	1	3.6	5	17.9
2 to 4 g	1	3.6	-	-
4 to 6 g	1	3.6	-	-
> 6 g	21	75	2	7.1

The majority of the bakery managers (96.7%) believe that salt has an impact on the health of all individuals. Only 3.3% considered that it has an impact only on the health of individuals with some disease.

Most respondents (66.7%) were unaware of the daily dose of salt recommended by the WHO (5 g/day); 6.7% were aware of such a recommendation.

Regarding the relationship between salt and sodium, 36.7% of the respondents answered that salt contains sodium; 53.3% did not know the relationship between the 2.

e) Laboratory results

A total of 155 food samples were collected and analyzed, 30 of which were bread (Portuguese/wheat bread) and 125 of which were meals ready for consumption. Of these, 75 were collected in the city of Praia (16 bread samples and 59 meal samples), and 80 were collected in the city of Mindelo (14 bread samples and 66 meal samples).

Of the bread samples analyzed, 53.3% (n = 16) had salt contents above the limit recommended by the

Most respondents (69%) stated that there were differences between types of salt; the remaining 31% responded that no differences exist between different types. The most reported differences between the types of salt were quality (41.2%), followed by texture (17.6%). Among the respondents, 11.8% considered iodized salt the best, and differences were reported in terms of appearance, taste/aroma, presence of iodine, importance for health and salt content in the recipe (5.9% each).

The amount of salt per kg of dough was greater than 14 g/kg in 3 bakeries, and in 2 bakeries, the amount of salt was greater than 6 g per loaf (Table 2).

Portuguese standard (Decree Law no. 75/2009, of 12 August). This standard defines a maximum permitted salt content, i.e., 1.4 g per 100 g of baked bread (or 0.55 g of sodium per 100 g of bread). Of the samples with salt contents greater than 1.4 g/100 g, 43.75% (n = 7) were collected in the city of Praia, and 57.14% (n = 8) were collected in the city of Mindelo (Appendix I).

Approximately 59.3% (n=35) of the meal samples collected in the restaurants in Praia and 42.4% in Mindelo had a NaCl greater than 5.0 g/meal, with a median of 5.03 g (table 3). All of these samples had a common component (meat) that underwent some traditional preserving process (use of salt as a preservative).

In the present study, the amount of Na varied between 0.26 g and 6.89 g per meal (Appendix II), with a median of 2.01 g (table 3), indicating a high Na content based on the recommended daily intake of less than 2 g of Na established by WHO (World Health Organization, 2020).

Table 3: S Statistical analysis of the salt content in the analyzed ready-to-eat food samples

Statistical analysis	Results (g/100g)			NaCl (g)/meal	Na (g)/meal
	NaCl	Na	Cl		
Mean	1,34	0,54	0,81	5,44	2,18
Median	1,30	0,52	0,78	5,03	2,01
Minimum	0,50	0,20	0,30	0,66	0,26

Maximum	2,70	1,08	1,62	17,22	6,89
Range	2,20	0,88	1,32	16,56	6,62
SD	0,50	0,20	0,30	2,83	1,13

Bakeries and restaurants in Mindelo used a mean of 1.50 ± 0.3 g of salt/100 g of bread and 1.4 ± 0.5 g of salt/100 g of meal respectively. In Praia, the

mean salt content in bread was 1.35 ± 0.25 g /100g, and that in meals was 1.3 ± 0.5 g/100 g respectively (Table 3 and table 4).

Table 4: Statistical analysis of the salt content in the analyzed bread samples

Statistical analysis	Results (g/100g)		
	NaCl	Na	Cl
Mean	1,46	0,58	0,88
Median	1,405	0,562	0,843
Minimum	1,1	0,44	0,66
Maximum	2,3	0,92	1,38
Range	1,2	0,48	0,72
SD	0,27	0,11	0,16

There were no significant differences between Praia and Mindelo regarding the salt content used in the bakeries ($\chi^2 = 71,778$; $p = 0.419$) and in the restaurants ($\chi^2 = 367,488$; $p = 0.641$).

Most of the salt samples collected (97%) contained iodine.

IV. DISCUSSION

Most restaurants did not have a specific location for storing salt. It was stored in its own packaging, protected from light, exposed to heat, stored in a nonventilated place, stored in the presence of chemicals, and exposed to moisture. Several studies have shown that storing iodized salt in a dry or cold place protected from sunlight preserves its composition (World Health Organization, 2007; Goris, J. M., et al., 2018; Mekonnen, T. C., et al., 2018; Abebe, Z., Gebeye, E., & Tariku, A. 2017).

In this study, most salt samples contained iodine (97%), a result that is in accordance with the WHO recommendations ($\geq 90\%$) established to eliminate iodine deficiency disorders (World Health Organization, 2007).

The high percentage (97%) of use of iodized salt by the restaurants and bakeries may be due to the efforts of the Cape Verdean government to stimulate improvements in the iodine status in recent years. This rate is consistent with that found in a study conducted in Nigeria (95%) and is higher than that found in studies in Ghana (75.6%) and Senegal (10%) on the salt iodine content in the household diet and associated factors (Anteneh, Z. A., Engidayehu, M., & Abeje, G. 2017; Obssie, G. F., Ketema, K., & Tekalegn, Y. 2020).

In this study, 50% of the analyzed bread samples revealed a salt content greater than the limit

recommended by the Portuguese standard, which is 1.4 g/100 g of bread or 0.55 g of sodium/100 g of bread (Assembleia da República, 2009). Bread is one of the most consumed foods in the world, accounting for an average of 30% of daily salt intake (2, 3). It contributes a significant percentage of salt to the diet of the Cape Verdean population.

Of the samples with salt contents greater than or equal to 1.4 g/100 g, 57.8% (n = 11) were collected in the city of Mindelo, and 42.2% (n = 8) were collected in the city of Praia (Assembleia da República, 2009). Similar results were found in a study on the evaluation of salt content in bread regularly consumed in the Eastern Mediterranean region (Assembleia da República, 2009; Al Jawaldeh, A., & Al-Khamaiseh, M. 2018).

The results of the present study indicate that the salt content in meals prepared in restaurants in Cape Verde is high ($\bar{x} = 5.44$ g/meal) based on the WHO recommendation (World Health Organization, Salt reduction, 2020).

A higher salt content was observed in meals that contained meat, french fries, chorizo or shellfish. This may be due to some traditional preservation processes (use of salt as a preservative) (Ludwig, L. M., et al., 2021). Conversely, the samples consisting essentially of salad had low salt contents (Nascimento RFd, et al., 2017).

In the present study, the amount of Na ranged from 780 mg to 4,330 mg/meal, with a median of 1,900 mg/meal, which exceeds not only the recommendation for a meal (18, 27, 28) but also the WHO daily recommendation, which establishes a maximum intake of 2,000 mg of Na/day (World Health Organization, Salt reduction, 2020).

The present findings draw attention to the greater sodium intake. Approximately 59,3% (n=35) of the meal samples collected in the restaurants in Praia and 42,4% in Mindelo had a NaCl greater than 5.0 g per meal, with a median of 5.03 g, suggesting a much higher daily NaCl intake than the maximum limit (<5 g/day) recommended by the WHO (World Health Organization, Hypertension, 2021; World Health Organization, Salt reduction, 2020). An intake of less than 5 g of salt per day in adults helps to reduce blood pressure and the risk of cardiovascular disease, stroke and heart attack (World Health Organization, Salt reduction, 2020).

The results should be interpreted considering that the data were collected only in the 2 main population centers. Thus, it is possible that there are other bakeries and restaurants that were not taken into account.

V. CONCLUSIONS

In general, food managers/handlers in restaurants and bakeries in the cities of Praia and Mindelo are unaware of the origin of the salt they use, the way in which it is transported and stored before reaching establishments. The lack of guidelines regarding the proper use of salt for each food preparation, thus contributing to its indiscriminate use.

Our study revealed unprecedented data on the profile and content of salt in the 2 main urban centers of Cape Verde (Praia and Mindelo). Thus contributing to the redesign of salt intake reduction strategies in these cities and in the country. It is imperative and urgent that national legislation regulates the unit weight of bread and the amount of salt that ready-to-eat food, including bread, should contain.

ACKNOWLEDGMENTS

The authors would like to thank everyone who contributed to this study.

Conflict of Interest

No funding was obtained for this study. The authors declare that they have no financial and/or personal conflicts of interest in the design and implementation of this study.

REFERENCES RÉFÉRENCES REFERENCIAS

1. World Health Organization, Noncommunicable diseases, 2021. Available from: <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>.
2. Portal informativo de Cabo Verde, Available from: <http://www.caboverde-info.com/Identidade/Geografia/Artigos/Coordenadas-geograficas>.
3. ALMEIDA S, Todas as medidas do governo são no sentido de afastar o cenário da fome e mitigar a insegurança alimentar, Expresso das ilhas, 2022.
4. Mills, K. T., Stefanescu, A., & He, J. (2020). The global epidemiology of hypertension. *Nature reviews. Nephrology*, 16(4), 223–237. <https://doi.org/10.1038/s41581-019-0244-2>.
5. World Health Organization, Hypertension, 2021. Available from: <https://www.who.int/news-room/fact-sheets/detail/hypertension>.
6. Molognoni, L., Daguer, H., Motta, G. E., Merlo, T. C., & Lindner, J. D. (2019). Interactions of preservatives in meat processing: Formation of carcinogenic compounds, analytical methods, and inhibitory agents. *Food research international* (Ottawa, Ont.), 125, 108608. <https://doi.org/10.1016/j.foodres.2019.108608>. Available from: <https://pubmed.ncbi.nlm.nih.gov/31554117/>.
7. Coelho PFdC. A prática social do consumo insustentável de alimentos para viagem no Brasil e Alemanha, 2021. Available from: <https://repositorio.ufc.br/handle/riufc/57926>.
8. Ruiz, A., Lima, M. G., Medina, L., Pinto, R. L., Barros, M., & Filho, A. (2020). Can Meals Outside Homes Impact Sodium Intake?. *Current developments in nutrition*, 4(6), nzaa091. Available from: <https://doi.org/10.1093/cdn/nzaa091>.
9. The National Academies Collection: Reports funded by National Institutes of Health. In: Oria M, Harrison M, Stallings VA, editors. *Dietary Reference Intakes for Sodium and Potassium*, Washington, 2019. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538102/>.
10. World Health Organization, Salt reduction, 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/salt-reduction>.
11. World Health Organization, Guideline: sodium intake for adults and children, 2012. Available from: <https://www.who.int/publications/item/9789241504836>.
12. Nascimento RFd, Gavron AB, Bowles S, Chaves ES, Bortolozo EAFQ, (2017) Determination of the sodium and potassium levels in meals served in a university restaurant in the south of Brazil. *Braz. J. Food Technol.*, v. 20, e2016167. Available from: <http://dx.doi.org/10.1590/1981-6723.16716>
13. Instituto Nacional de Estatística de Cabo Verde, Anuário Estatístico – Cabo Verde 2018, 2020. Available from: <https://ine.cv/publicacoes/anuario-estatistico-cabo-verde-2018/>.
14. Viegas CACL. O sal na alimentação dos jovens: avaliação e percepção do sal das refeições. 2013. Available from: <http://hdl.handle.net/10362/13224>.
15. Associação de restauração e similares de Portugal, Código de boas práticas para o transporte de



alimentos. 2008. Available from: https://ahresp.com/app/uploads/2018/10/Codigo-CBPH_AHRESP.pdf.

16. World Health Organization, (2007); Assessment of iodine deficiency disorders and monitoring their elimination: a guide for programme managers, 3rd ed. World Health Organization. Available from: <https://apps.who.int/iris/handle/10665/43781>.
17. Jooste PL, Strydom E, (2010), Methods for determination of iodine in urine and salt. Best practice & research Clinical endocrinology & metabolism; 24(1): 77-88.
18. Ward, R. E., & Carpenter, C. E. (2010). Traditional methods for mineral analysis. In Food analysis (p. 207), 4th ed. Springer, New York.
19. Anteneh, Z. A., Engidayehu, M., & Abeje, G. (2017). Iodine content of dietary salt at household level and associated factors using iodometric titration methods in Dera District, Northwest Ethiopia. BMC nutrition, 3, 83. Available from: <https://doi.org/10.1186/s40795-017-0203-x>
20. Obssie, G. F., Ketema, K., & Tekalegn, Y. (2020). Availability of Adequately Iodized Dietary Salt and Associated Factors in a Town of Southeast Ethiopia: A Community-Based Cross-Sectional Survey. Journal of nutrition and metabolism, 2020, 1357192. Available from: <https://doi.org/10.1155/2020/1357192>.
21. Goris, J. M., Temple, V. J., Zomerdijk, N., & Codling, K. (2018). Iodine status of children and knowledge, attitude, practice of iodised salt use in a remote community in Kerema district, Gulf province, Papua New Guinea. PloS one, 13(11), e0197647. Available from: <https://doi.org/10.1371/journal.pone.0197647>
22. Mekonnen, T. C., Eshete, S., Wasihun, Y., Arefaynie, M., & Cherie, N. (2018). Availability of adequately iodized salt at household level in Dessie and Combolcha Towns, South Wollo, Ethiopia. BMC public health, 18(1), 1152. Available from: <https://doi.org/10.1186/s12889-018-6066-5>
23. Abebe, Z., Gebeye, E., & Tariku, A. (2017). Poor dietary diversity, wealth status and use of un-iodized salt are associated with goiter among school children: a cross-sectional study in Ethiopia. BMC public health, 17(1), 44. Available from: <https://doi.org/10.1186/s12889-016-3914-z>
24. Assembleia da República. Lei n.º 75/2009 de 12 de agosto Diário da República, 1.ª série, 2009 [5225 - 6]. Available from: <https://dre.pt/dre/detalhe/lei/75-2009-493513>.
25. Al Jawaldeh, A., & Al-Khamaiseh, M. (2018). Assessment of salt concentration in bread commonly consumed in the Eastern Mediterranean Region. Eastern Mediterranean health journal = La revue de santé de la Méditerranée orientale = al-Majallah al-sihhiyah li-sharq al-mutawassit, 24(1), 18-24.
26. Ludwig, L. M., Freitas, F. C. de, Chaves, M. A., Canan, C., & Corso, M. P. (2021). Estudo do processo de salga na produção de paleta suína salgada curada dessecada / Salting process study in obtaining a dried cured salted pork shoulder. Brazilian Journal of Development, 7(8), 85795-85806. Available from: <https://doi.org/10.34117/bjdv7n8-675>.
27. National Research Council (US) Subcommittee on the Tenth Edition of the Recommended Dietary Allowances. (1989). Recommended Dietary Allowances: 10th Edition. National Academies Press (US).
28. Feeding Guide for the Brazilian Population: Promoting the Health Food. (2008). – Brasília: Ministério da Saúde,. 210 p. Available from: https://bvsms.saude.gov.br/bvs/publicacoes/guia_alimentar_populacao_brasileira_2008.pdf.

Supplementary

Questionnaire applied to restaurants

A – SOCIODEMOGRAPHIC CHARACTERISTICS

1. Gender: F M 2. Age: _____ 3. Education level: _____

4. Profession/role

5. In your establishment, is there a standard that dictates the ideal amount of salt to be used in food preparation?

Yes No

6. Do you have any training in the field of Nutrition?

Yes No

6.1. If yes, please describe: _____

B – FOOD PREPARATION IN RESTAURANTS

7. What type of salt do you use to prepare food?

Coarse salt Refined salt Sea salt Fleur de sel Iodized salt

Industrialized Nonindustrialized Other _____

8. Do you consider there to be a difference between these types of salt? Yes No

8.1. If yes, please describe: _____

9. What is the origin of the salt you use in your establishment?
 National International
Island _____ Country _____

10. What was the means used to transport the salt to your establishment?
 Car Boat Airplane Other _____

11. Is there a specific location for salt storage?
 Yes No
11.1. If so, how is it stored?
 In its own packaging Protected from light Exposed to light Exposed to moisture
 Exposed to heat In the presence of chemical products In a nonventilated location
 Other _____

11.2. If not, specify the location and storage conditions:
Location: _____
 In its own packaging Protected from light Exposed to light Exposed to moisture
 Exposed to heat In the presence of chemical products In a nonventilated location
 Other _____

12. Do you use any utensil to handle the salt?
 Yes No
12.1. If so, which ones?
 Spoon Cup Other _____

13. What is the amount of salt you use, per person, to make a soup?
 <1 g (less than 1/2 teaspoon) 1 to 2 g (1/2 to 1 teaspoon) 2 to 4 g (1 to 2 teaspoons)
 4 to 6 g (2 to 3 teaspoons) >6 g (more than 1 tablespoon)

14. How much salt do you use, per person, to prepare a main dish?
 <1 g (less than 1/2 teaspoon) 1 to 2 g (1/2 to 1 teaspoon) 2 to 4 g (1 to 2 teaspoons)
 4 to 6 g (2 to 3 teaspoons) >6 g (more than 1 tablespoon)

15. Do you consider that salt intake has an impact on health?
 Yes, for all individuals
 Yes, but only for individuals with some disease
 No

16. What is the daily dose of salt recommended by the World Health Organization?
 12 g/day
 10 g/day
 5 g/day
 3 g/day
 Other _____

17. The following describes the relationship between salt and sodium:
 They are the same
 Salt contains sodium
 Sodium contains salt

18. If you were asked to reduce the amount of salt used in cooking, what measures would you use to flavor the food, compensating for the lack of salt (you can choose more than one option)?
 Use of aromatic herbs
 Use of spices
 Use of concentrates (bouillon cubes)
 Other _____

19. Do you consider it important to reduce salt intake?
 Yes No Yes, but only for individuals with diseases

20. If you tried to reduce salt in the preparation of the meals you cook, what do you think would be the consumers' reaction (you can choose more than one option)?
 Eat in another establishment/location
 Complain to those responsible for the meals



Complain in the complaints book
 Use a salt shaker, adding fine salt to the meals
 No reaction
 Another reaction _____

21. What are the diseases most associated with salt intake (you can choose more than one option)?

High cholesterol
 Heart disease
 Hypertension
 Kidney disease
 Obesity
 Other _____

Many thanks for your participation!

Questionnaire applied to bakeries

A – SOCIODEMOGRAPHIC CHARACTERISTICS

1. Gender: F M 2. Age: _____ 3. Education level: _____
 4. Profession/role _____
 5. In your establishment, is there a standard that dictates the ideal amount of salt to be used in food preparation?
 Yes No
 6. Do you have any training in the field of Nutrition?
 Yes No
 6.1. If yes, please describe: _____

B – BREAD PREPARATION IN BAKERIES

7. What type of salt do you use to make bread?
 Coarse salt Refined salt Sea salt Fleur de sel Iodized salt
 Industrialized Nonindustrialized Other _____
 8. Do you consider there to be a difference between these types of salt?
 Yes No
 8.1. If yes, please describe: _____
 9. What is the origin of the salt you use in your establishment?
 National International
 Island _____ Country _____
 10. What was the means used to transport the salt to your establishment?
 Car Boat Airplane Other _____
 11. Is there a specific location for salt storage?
 Yes No
 11.1. If so, how is it stored?
 In its own packaging Protected from light Exposed to light Exposed to moisture
 Exposed to heat In the presence of chemical products In a nonventilated location
 Other _____
 11.2. If not, specify the location and storage conditions:
 Location: _____
 In its own packaging Protected from light Exposed to light Exposed to moisture
 Exposed to heat In the presence of chemical products In a nonventilated location
 Other _____
 12. Do you use any utensil to handle the salt?
 Yes No
 12.1. If so, which ones? Spoon Cup Other _____
 13. What is the amount of salt you use, per kilo of dough?
 <1 g (less than 1/2 teaspoon) 1 to 2 g (1/2 to 1 teaspoon) 2 to 4 g (1 to 2 teaspoons)
 4 to 6 g (2 to 3 teaspoons) >6 g (more than 1 tablespoon) Other _____
 14. What is the amount of salt you use per loaf of bread?
 <1 g (less than 1/2 teaspoon) 1 to 2 g (1/2 to 1 teaspoon) 2 to 4 g (1 to 2 teaspoons) 4 to 6 g (2 to 3 teaspoons)
 >6 g (more than 1 tablespoon)

15. Do you consider that salt intake has an impact on health?

- Yes, for all individuals
- Yes, but only for individuals with some disease
- No

16. What is the daily dose of salt recommended by the World Health Organization?

- 12 g/day
- 10 g/day
- 5 g/day
- 3 g/day
- Other _____

17. The following describes the relationship between salt and sodium:

- They are the same
- Salt contains sodium
- Sodium contains salt

18. If you were asked to reduce the amount of salt used in bread-making, what measures would you use to flavor the bread, compensating for the lack of salt (you can choose more than one option)?

- Use of aromatic herbs
- Use of spices
- Use of concentrates (bouillon cubes)
- Other _____

19. Do you consider it important to reduce salt intake?

- Yes
- No
- Yes, but only for individuals with diseases

20. If you tried to reduce the salt in the preparation of the bread you produce, what do you think would be the consumers' reaction (you can choose more than one option)?

- Buy bread at another establishment/location
- Complain to those responsible for the establishment
- Complain in the complaints book
- Use of a salt shaker, adding fine salt to the bread
- No reaction
- Another reaction _____

21. What are the diseases most associated with salt intake (you can choose more than one option)?

- High cholesterol
- Heart disease
- Hypertension
- Kidney disease
- Obesity
- Other _____

Many thanks for your participation!

Amount of salt present in the analyzed bread samples

N	Provenance	Results (g/100g)		
		NaCl	Na	Cl
1	Praia	1,2	0,48	0,72
2	Praia	1,3	0,52	0,78
3	Praia	1,3	0,52	0,78
4	Praia	1,2	0,48	0,72
5	Praia	1,2	0,48	0,72
6	Praia	1,2	0,48	0,72
7	Praia	1,3	0,52	0,78



8	Praia	1,1	0,44	0,66
9	Praia	1,43	0,572	0,858
10	Praia	1,81	0,724	1,086
11	Praia	1,41	0,564	0,846
12	Praia	1,5	0,6	0,9
13	Praia	2,03	0,812	1,218
14	Praia	1,46	0,584	0,876
15	Praia	1,7	0,68	1,02
16	Praia	1,4	0,56	0,84
Mean		1,40	0,56	0,85
SD		0,25	0,10	0,15
17	Mindelo	1,6	0,64	0,96
18	Mindelo	1,2	0,48	0,72
19	Mindelo	1,53	0,612	0,918
20	Mindelo	1,4	0,56	0,84
21	Mindelo	1,5	0,6	0,9
22	Mindelo	2,3	0,92	1,38
23	Mindelo	1,2	0,48	0,72
24	Mindelo	1,4	0,56	0,84
25	Mindelo	1,6	0,64	0,96
26	Mindelo	1,4	0,56	0,84
27	Mindelo	1,9	0,76	1,14
28	Mindelo	1,2	0,48	0,72
29	Mindelo	1,6	0,64	0,96
30	Mindelo	1,5	0,6	0,9
Mean		1,52	0,61	0,91
SD		0,29	0,12	0,18

Amount of salt present in the analyzed ready-to-eat food samples

N	Provenance	Results (g/100g)			NaCl (g)/meal	Na (g)/meal
		NaCl	Na	Cl		
1	Praia	1,10	0,44	0,66	2,07	0,83
2	Praia	1,60	0,64	0,96	4,11	1,64
3	Praia	0,70	0,28	0,42	1,90	0,76
4	Praia	1,10	0,44	0,66	3,30	1,32
5	Praia	1,14	0,46	0,68	3,45	1,38
6	Praia	2,30	0,92	1,38	7,06	2,82
7	Praia	1,30	0,52	0,78	4,28	1,71
8	Praia	1,44	0,58	0,86	4,81	1,92
9	Praia	1,20	0,48	0,72	4,08	1,63
10	Praia	1,30	0,52	0,78	4,71	1,88
11	Praia	1,60	0,64	0,96	5,84	2,34
12	Praia	1,10	0,44	0,66	4,08	1,63
13	Praia	2,00	0,80	1,20	7,54	3,02
14	Praia	1,40	0,56	0,84	5,60	2,24

15	Praia	1,80	0,72	1,08	7,22	2,89
16	Praia	1,90	0,76	1,14	7,81	3,12
17	Praia	1,30	0,52	0,78	5,62	2,25
18	Praia	1,20	0,48	0,72	5,20	2,08
19	Praia	1,30	0,52	0,78	5,64	2,26
20	Praia	1,30	0,52	0,78	5,69	2,28
21	Praia	0,66	0,26	0,40	2,90	1,16
22	Praia	1,30	0,52	0,78	5,79	2,31
23	Praia	0,90	0,36	0,54	4,11	1,65
24	Praia	1,20	0,48	0,72	5,58	2,23
25	Praia	1,00	0,40	0,60	4,75	1,90
26	Praia	1,20	0,48	0,72	5,76	2,30
27	Praia	0,80	0,32	0,48	4,07	1,63
28	Praia	1,20	0,48	0,72	6,13	2,45
29	Praia	0,90	0,36	0,54	4,66	1,86
30	Praia	0,90	0,36	0,54	4,73	1,89
31	Praia	1,10	0,44	0,66	6,18	2,47
32	Praia	0,60	0,24	0,36	3,49	1,39
33	Praia	1,00	0,40	0,60	5,90	2,36
34	Praia	0,80	0,32	0,48	4,76	1,90
35	Praia	1,30	0,52	0,78	7,75	3,10
36	Praia	0,50	0,20	0,30	2,99	1,19
37	Praia	1,10	0,44	0,66	6,60	2,64
38	Praia	2,40	0,96	1,44	14,50	5,80
39	Praia	2,40	0,96	1,44	14,76	5,90
40	Praia	1,80	0,72	1,08	11,12	4,45
41	Praia	0,80	0,32	0,48	4,95	1,98
42	Praia	1,10	0,44	0,66	6,82	2,73
43	Praia	0,80	0,32	0,48	5,44	2,18
44	Praia	0,90	0,36	0,54	6,26	2,51
45	Praia	0,70	0,28	0,42	5,03	2,01
46	Praia	0,80	0,32	0,48	5,94	2,38
47	Praia	0,90	0,36	0,54	7,02	2,81
48	Praia	0,90	0,36	0,54	7,34	2,93
49	Praia	0,90	0,36	0,54	7,39	2,96
50	Praia	2,00	0,80	1,20	17,22	6,89
51	Praia	0,90	0,36	0,54	2,28	0,91
52	Praia	1,30	0,52	0,78	5,20	2,08
53	Praia	2,70	1,08	1,62	10,83	4,33

54	Praia	0,90	0,36	0,54	3,60	1,44
55	Praia	2,60	1,04	1,56	10,40	4,16
56	Praia	1,50	0,60	0,90	2,82	1,13
57	Praia	1,40	0,56	0,84	3,57	1,43
58	Praia	1,00	0,40	0,60	6,00	2,40
59	Praia	1,40	0,56	0,84	5,60	2,24
Mean		1,26508	0,5061	0,75898	5,93644	2,37424
SD		0,49818	0,19936	0,29882	2,91939	1,16795
60	Mindelo	1,50	0,60	0,90	6,00	2,40
61	Mindelo	1,10	0,44	0,66	4,40	1,76
62	Mindelo	1,00	0,40	0,60	4,00	1,60
63	Mindelo	0,90	0,36	0,54	2,70	1,08
64	Mindelo	1,60	0,64	0,96	4,80	1,92
65	Mindelo	1,40	0,56	0,84	5,60	2,24
66	Mindelo	1,20	0,48	0,72	4,80	1,92
67	Mindelo	2,10	0,84	1,26	8,40	3,36
68	Mindelo	1,20	0,48	0,72	4,80	1,92
69	Mindelo	1,50	0,60	0,90	6,00	2,40
70	Mindelo	1,70	0,68	1,02	6,80	2,72
71	Mindelo	0,90	0,36	0,54	3,60	1,44
72	Mindelo	2,10	0,84	1,26	8,40	3,36
73	Mindelo	1,20	0,48	0,72	4,80	1,92
74	Mindelo	1,40	0,56	0,84	5,60	2,24
75	Mindelo	2,30	0,92	1,38	9,20	3,68
76	Mindelo	1,40	0,56	0,84	4,20	1,68
77	Mindelo	0,70	0,28	0,42	2,80	1,12
78	Mindelo	0,60	0,24	0,36	0,66	0,26
79	Mindelo	1,20	0,48	0,72	1,66	0,66
80	Mindelo	2,00	0,80	1,20	2,92	1,17
81	Mindelo	1,50	0,60	0,90	2,24	0,89
82	Mindelo	1,10	0,44	0,66	1,76	0,70
83	Mindelo	1,80	0,72	1,08	3,10	1,24
84	Mindelo	0,70	0,28	0,42	1,24	0,50
85	Mindelo	1,40	0,56	0,84	2,56	1,02
86	Mindelo	1,50	0,60	0,90	2,76	1,10
87	Mindelo	0,60	0,24	0,36	1,12	0,45
88	Mindelo	0,90	0,36	0,54	1,69	0,68
89	Mindelo	0,70	0,28	0,42	1,39	0,55
90	Mindelo	0,80	0,32	0,48	1,60	0,64

91	Mindelo	0,80	0,32	0,48	1,68	0,67
92	Mindelo	1,90	0,76	1,14	4,20	1,68
93	Mindelo	2,00	0,80	1,20	4,54	1,82
94	Mindelo	0,80	0,32	0,48	1,91	0,76
95	Mindelo	1,70	0,68	1,02	4,17	1,67
96	Mindelo	2,30	0,92	1,38	5,68	2,27
97	Mindelo	1,30	0,52	0,78	3,38	1,35
98	Mindelo	1,20	0,48	0,72	3,24	1,30
99	Mindelo	1,70	0,68	1,02	4,81	1,92
100	Mindelo	2,20	0,88	1,32	6,38	2,55
101	Mindelo	2,00	0,80	1,20	6,00	2,40
102	Mindelo	2,10	0,84	1,26	6,45	2,58
103	Mindelo	0,60	0,24	0,36	1,86	0,74
104	Mindelo	2,00	0,80	1,20	6,38	2,55
105	Mindelo	1,30	0,52	0,78	4,55	1,82
106	Mindelo	2,30	0,92	1,38	8,14	3,26
107	Mindelo	1,30	0,52	0,78	4,65	1,86
108	Mindelo	1,60	0,64	0,96	5,76	2,30
109	Mindelo	1,70	0,68	1,02	6,24	2,50
110	Mindelo	1,70	0,68	1,02	6,29	2,52
111	Mindelo	2,40	0,96	1,44	9,60	3,84
112	Mindelo	1,80	0,72	1,08	7,25	2,90
113	Mindelo	0,60	0,24	0,36	2,46	0,98
114	Mindelo	1,70	0,68	1,02	7,31	2,92
115	Mindelo	1,60	0,64	0,96	7,20	2,88
116	Mindelo	0,90	0,36	0,54	4,14	1,66
117	Mindelo	0,90	0,36	0,54	4,31	1,72
118	Mindelo	1,50	0,60	0,90	7,79	3,11
119	Mindelo	0,90	0,36	0,54	4,73	1,89
120	Mindelo	1,00	0,40	0,60	5,70	2,28
121	Mindelo	1,40	0,56	0,84	8,15	3,26
122	Mindelo	1,50	0,60	0,90	9,33	3,73
123	Mindelo	1,90	0,76	1,14	12,96	5,18
124	Mindelo	1,50	0,60	0,90	10,41	4,16
125	Mindelo	1,30	0,52	0,78	11,01	4,40
Mean		1,41515	0,56606	0,84909	5,00394	2,00076
SD		0,49404	0,19762	0,29642	2,66494	1,06596