

CrossRef DOI of original article:

# A Retrospective Study on the Complications of Ilizarov Technique in the Treatment of Neurogenic Talipes Equinovarus

Xinjiang Lin,<sup>1</sup> Shouping Chen<sup>2</sup> and Shengke Xie<sup>3</sup>

<sup>1</sup> Annual Ring Orthopaedic Hospital Group, Changsha, Hunan 410000, China

*Received: 1 January 1970 Accepted: 1 January 1970 Published: 1 January 1970*

---

## Abstract

In order to discuss the postoperative complications of Ilizarov technique in the treatment of neurogenic talipes equinovarus, we conducted a retrospective study on the postoperative complications of neurogenic talipes equinovarus patients treated with Ilizarov technique from January 2013 to December 2020. 182 patients (228 feet), 134 males (171 feet) and 48 females (57 feet). The age ranged from 4 to 70 ( $\bar{x}23.6 \pm 13.9$ ) years old. There were 44 cases of sequelae of cerebral palsy (24.2

---

**Index terms**— Ilizarov technology; Horseshoe varus foot; complications; tissue displacement syndrome; Retrospective study.

## 1 Introduction

neurogenic talipes equinovarus is a disease with the highest prevalence and the greatest harm among the modern disabled people. It is one of the common orthopedic malformations, and also a disease with the longest treatment cycle, greater difficulty and poor patient satisfaction. The main pathological change is due to the dislocation of calcaneus, talus and scaphoid, which leads to imbalance of muscle force and local tendon contracture. It is a three-dimensional deformity caused by complex pathological changes of soft tissue and bone and joint, mainly manifested as foot drop, high arch, varus, adduction and other foot and ankle deformities. Due to the relative lack of subcutaneous soft tissue and poor blood supply in the ankle, traditional surgery is easy to cause skin ischemic necrosis, postoperative secondary infection, soft tissue scar contracture and other complications, and the recurrence rate is as high as about 20% [1]. In order to reduce the complications of correction of neurogenic talipes equinovarus and reduce the recurrence rate after operation, we applied Ilizarov technology for treatment. Ilizarov technology is to install a special external fixator in the ankle, and gradually correct the deformity of horseshoe foot through slow tissue drafting. The clinical application of many scholars has proved that [2-3], not only the deformity correction is satisfactory, but also the shape and function of the foot can be preserved to the maximum extent, and at the same time, serious complications can be avoided and reduced. It is a safe and reliable method for the correction of neurogenic clubfoot. Due to the different treatment methods and various methods, it is necessary to carry out personalized force line analysis, osteotomy plane and angle design, neuromyoelectric test and analysis, determine tendon transposition, repair of nerve, blood vessel and skin tissues, and long-term, systematic and comprehensive rehabilitation training after surgery according to the patient's different age, sex, occupation, deformity, etc. Therefore, orthopedic physicians need to have comprehensive knowledge of basic medicine and clinical medicine. However, at present, there are not many orthopedic doctors who have received standardized training in China, and the amount of orthopedic operations is large, with many therapeutic effects and postoperative complications. Therefore, this paper conducts a retrospective clinical study on the treatment effect and postoperative complications of 182 patients with complete data from 262 patients with neurogenic talipes equinovarus treated with Ilizarov technology in our hospital from January 2013 to December 2020. 2) Plantar flexion deformity is less than 400 and there is no complaint of discomfort;

## 2 II.

### 3 Proposed Method

3) Patients with incomplete clinical data and follow-up less than 3 times. c) General data 182 patients (228 feet) in this group. Among them, 136 were unilateral and 46 bilateral. 134 males (171 feet) and 48 females (57 feet). The age ranged from 4 to 70 ( $23.64 \pm 13.96$ ) years, 12 cases were  $< 7$  years old (6.6%), 52 cases were 8-17 years old (28.6%), 72 cases were 18-30 years old (39.6%), 44 cases were 31-59 years old (24.2%), and 2 cases were  $> 60$  years old (1.1%). Figure 1. Classification of diseases: 44 cases (24.2%) of sequelae of cerebral palsy, 112 cases (61.5%) of sequelae of poliomyelitis, 18 cases (9.9%) of congenital clubfoot, and 8 cases (4.4%) of traumatic clubfoot.

### 4 d) Treatment method

According to the degree of bone deformity of the patient's foot, muscle strength, age, the degree of cooperation of the patient and his family, the ability of the doctor to master this technology and other factors, a personalized surgical plan is formulated. First, according to the degree of deformity of the patient's foot, the posterior medial soft tissue release, Achilles tendon lengthening, external transfer of tibial anterior muscle, osteotomy of the three joints of the foot, internal rotation osteotomy under the tibial tubercle and other soft tissue release, muscle force balance, osteotomy correction and joint fusion were selected, and then the Ilizarov external fixator was used for correction.

### 5 e) Data collection and sorting

During the hospitalization, the responsible physician (with 5-10 years of clinical experience) is responsible for tabulating, statistical analysis and sorting out the clinical symptoms, signs and relevant examinations of patients after using the Ilizarov external fixator. After the patient leaves the hospital, the responsible physician and the customer service department personnel will conduct telephone or on-site follow-up to the patient and his family members 1 month, 3 months, 6 months, 1 year or 2 years after the patient leaves the hospital. The responsible physician focuses on understanding the patient's disease, guiding the later rehabilitation, prevention and treatment of complications, frame adjustment, frame removal, reexamination, etc. The customer service department staff mainly understand the patient's recent situation, physical recovery, and the evaluation and suggestions on the hospital's work.

### 6 f) Statistical methods

All data were analyzed by SPSS 20.0 software. The measurement data are expressed as mean  $\pm$  standard deviation ( $\pm S$ ). Take the percentage of 5 groups of data of the same variable, calculate the 99% confidence interval and the correlation between the variables. The P value of the detection level is less than 0.05 on both sides, which is considered to be statistically significant.

## 7 III.

### 8 Results and Analysis a) Evaluation method and efficacy

According to the ICFSG scoring standard, the patients were scored according to the 2-year follow-up after surgery. 228 feet, ICFSG score: excellent 136 feet, good 67 feet, fair 11 feet, poor 8 feet, the excellent and good rate is 89.04%.

### 9 b) Postoperative complications and analysis

The time of using Ilizarov external frame for this group of cases was 36-381 days ( $\bar{x}86.3 \pm 56.5$ ). The patients were followed up for 1 month, 3 months, 6 months, 1 year or 2 years after discharge. The follow-up time ranged from 1 to 24 months, with an average of 16.2 months. Among them, 65 people (77 feet) had 16 kinds of complications, the incidence was 33.77%. It is significantly higher than 22.5% reported in the data [4]. We think it is related to different statistical caliber. English literature records the frequency of various complications during limb lengthening of Ilizarov, which may reach 100% [5,6].

### 10 i. Early complications

In this group, 147 patients (178 feet) had pain, swelling, numbness and other symptoms after operation, accounting for 78.07%. It lasted for 2-12 ( $\bar{x}5.85 \pm 2.41$ ) weeks, 13 feet had needle infection, 17 feet had loose connecting rods, 5 feet had broken needles, 3 feet had nerve injury and 2 feet had skin necrosis. Needle infection is a common complication in the process of Ilizarov external fixation device orthopedics, with an incidence of 21%-42% [7,8,9,10]. The incidence of cases in this group is low (5.7%). The main causes are thermal burns to tissues during operation, skin and muscle injuries caused by long-term traction, exposed

---

## 94 11 ??

95 needle mouth pollution, and skin diseases of a few patients themselves. The causes of this group of cases were  
96 prevented in advance, such as using a protective sleeve when threading the needle during the operation, paying  
97 attention to the direction and strength of the steel needle and the condition of skin displacement, ensuring the  
98 skin is clean, and timely dealing with early infection. Therefore, the needle infection rate is far lower than that  
99 reported in the literature. In this group, one patient suffered from allergic dermatitis with infection and finally  
100 osteomyelitis due to untimely treatment of early needle infection. The loose connecting rod is mainly due to the  
101 loose screw fixation or more patient activities, especially in the rehabilitation training of patients with spastic  
102 cerebral palsy. order to relieve local tension pain, individual patients adjust the screw by themselves. Two cases  
103 of common peroneal nerve injury and one case of posterior tibial nerve injury were caused by intraoperative  
104 traction. After the application of neurotrophic drugs, rehabilitation physiotherapy and other treatments, they  
105 all recovered completely. Two patients suffered from skin necrosis within 1 cm around the anterior medial tibial  
106 needle path due to skin heat injury, and recovered after dressing change. 5. The needle breaks at the edge of  
107 the fixed screw. Pull out the broken end and fix the broken pin again with the connecting piece, which does not  
108 affect the fixing effect.

109 ii.

## 110 12 Late complications

111 The application of Ilizarov technology has created unique technical advantages for limb orthopedics, but there may  
112 be a variety of complications in the application of Ilizarov technology, including needle infection, osteomyelitis,  
113 footswelling, toe flexion deformity, metatarsophalangeal joint subluxation, foot stiffness and even recurrence  
114 [11,12,13]. It is difficult to treat adult neurogenic talipesquinovarus, especially in patients with long course  
115 and severe deformity. Although there are many methods of surgical treatment, it is difficult to correct all  
116 malformations in one operation [14]. Repeated soft tissue release and osteotomy orthopaedic surgery are more  
117 likely to cause stiffness, small and pain of the foot and ankle [15,16]. Beaty JH. Freedman JA et al. [12,13,]  
118 believed that ankle and subtalar joint stiffness, arthritis, pain and residual deformity existed for a long time. In  
119 this group, 1.8% of the patients had limited knee movement, 2.6% had ankle arthritis and 1.3% had subtalar  
120 joint stiffness, which was lower than that reported in the literature. The deformity of toe flexion contracture was  
121 5.7%. After orthopedic surgery, a kind of instinctive anti fall reflex causes the toe to flex and contract for a long  
122 time when the foot touches the ground, resulting in toe flexion contracture deformity. Parmanand Gupta et al.  
123 [17] believed that toe flexion contracture deformity is a complication that is difficult to treat, and once it occurs,  
124 it will not be able to participate in sports competitions as a professional athlete. In this group, 43 patients (49  
125 feet) had limb pain, swelling, numbness and other symptoms more than 3 months after surgery, which we call  
126 "tissue displacement syndrome", accounting for 21.49%. The incidence of complications in different age groups is  
127 shown in Table 1. iii. Tissue displacement syndrome The clinical characteristics of equinovarus foot are mainly  
128 ankle plantar flexion, heel varus and forefoot adduction [18]. During the surgical correction, tendon transposition,  
129 such as Achilles tendon extension and tibial anterior tendon insertion, should be done. A few patients need to  
130 do rectangular shortening osteotomy of the lateral column of the calcaneus, then use a steel needle to cross the  
131 calcaneus and metatarsal, connect the half ring and fix it on the calcaneus and foot back, and then slowly pull it  
132 for 2 to 3 months, so that the forefoot and midfoot gradually rotate outward and turn outward, so as to restore  
133 the normal appearance of the foot. During the whole operation and slow tissue traction process, the tissue has  
134 been displaced, local microvessels have been damaged, and circulation obstacles have occurred, leading to local  
135 swelling; Pain caused by tissue injury, hemorrhage, edema, and inflammatory stimulation; Tissue compression,  
136 nerve damage, numbness. This series of pathophysiological reactions is called "tissue displacement syndrome".  
137 The similar reaction in the earlier stage is "tissue displacement reaction". If the angle of the external frame  
138 is adjusted properly or the speed of frame adjustment is slowed down, drug treatment, physical therapy and  
139 other comprehensive treatments are carried out, and the symptoms still have no significant change and affect  
140 the normal walking function after more than 3 months, it is called "tissue displacement syndrome". The severity  
141 of the syndrome is related to surgical trauma, displacement angle, traction time, speed, patient age, and body  
142 regeneration and repair ability. In this group, most of the patients with "tissue displacement syndrome" occurred  
143 in adults over 18 years old, and there was a significant positive correlation with age ( $P < 0.05-0.01$ ). Among  
144 the 43 patients with "tissue displacement syndrome" in this group, 26 (60.47%) were followed up 1 year after  
145 discharge, and 12 (27.91%) were followed up 2 years after discharge. The symptoms such as swelling and pain  
146 of the patients' limbs basically disappeared, and some patients felt a little numbness locally, but the walking  
147 function of the limbs was not affected. Five patients were not followed up. The symptoms of patients with  
148 "tissue displacement syndrome" persist, but the prognosis is good.

## 149 13 c) Correlation between complications and age

150 The correlation test was conducted between the first 6 complications with high incidence rate and different age  
151 variables. Among them, pain, swelling and numbness (tissue displacement syndrome) were positively correlated  
152 with age ( $P < 0.05-0.01$ ). There was no correlation between needle infection, loose connecting rod and toe flexion  
153 deformity and age ( $P > 0.05$ ). Table 2.

## 14 d) Prevention measures

The distraction osteogenesis theory of Ilizarov technology has proved that the external fixator is beneficial to the shape recovery of various bone tissues, the adjustment and maintenance of limb length during the slow traction process, so that the correction of talipes equinovarus deformity can obtain satisfactory results for clinicians and patients [18,19]. However, due to the wide variety of configurations of external fixation devices, wide surgical indications, and long learning curve of postoperative management process and doctors, errors are inevitable in the treatment process, and problems in any link, such as needle threading and installation of external fixators, needle bag, postoperative management and guidance of patients' functional training, and the time to remove external fixators, may occur large and small complications [4]. However, through our efforts, most of these complications can be avoided. Paley [20] divided the problems arising from the application of Ilizarov technology into three categories: one is called problems, which can be solved without surgery; The second type is called obstacle, which needs to be solved by reoperation, but will not leave sequela; The three types are called complications, which will still leave morphological abnormalities or dysfunction after treatment. According to the Paley classification, there are 6 kinds of "problems" in this group, of which 3 are "tissue displacement syndrome"; 6 "obstacles"; There were 4 kinds of "complications", 25 feet, accounting for 10.96%.

Orthopedic surgery (including peripheral nerve surgery) is recommended by the bone and joint professional committee of the Chinese Rehabilitation Medical Association, the China Brain Palsy Multidisciplinary Cooperation Alliance, and the surgical treatment experts of spastic cerebral palsy by consensus as the second stage surgery of spastic cerebral palsy, an important supplement to SPR surgery, and is not recommended to take corrective surgery first. It is suggested that rehabilitation training is an important guarantee for postoperative functional improvement. Advocate the concept of "three points operation, seven points training" [21]. Therefore, the key to the successful treatment of neurogenic clubfoot is to objectively predict the surgical effect, fully communicate with patients, reduce patients' expectations, improve patients' compliance, strengthen the sense of responsibility of medical personnel, scientifically, rigorously and strictly control the surgical indications, reduce complications, strengthen long-term and standardized rehabilitation training for patients after surgery, and cooperate with doctors and patients.

IV.

## 15 Conclusion

In this group, 182 patients (228 feet) with neurogenic clubfoot were treated with Ilizarov external fixator, and the excellent and good rate was 89.04%. There were 16 kinds of complications, accounting for 33.77%. In the early stage, pain, swelling, numbness and other "tissue displacement reactions" were the main symptoms (78.07%). In the later stage, "tissue displacement syndrome" occurred (21.49%), but the prognosis was good. Among them, 88.37% of the patients had basically recovered from follow-up data within two years. The loose connecting rod accounted for 7.5%, toe flexion deformity and needle infection accounted for 5.7% respectively. The main complications increased with age, and there was a significant positive correlation between complications and age ( $P < 0.05$ ). However, through our efforts, most of these complications can be avoided. Therefore, in the process of applying Ilizarov technology to correct neurogenic talipes equinovarus, we should strengthen the sense of responsibility of medical personnel and improve their professional skills. Scientific, rigorous and strict control of surgical indications. Do a good job of communication between doctors and patients before surgery to improve patients' compliance with treatment. Personalized installation and adjustment of the external frame, strengthening long-term postoperative rehabilitation training, and other factors are the key to reduce complications and successfully treat neurogenic clubfoot.




Figure 1: a) Inclusion criteria 1 )

1

complication	77 year	8717 year	18730 year	31759 year	760 year	Total(%)
Pain (>3 months)	1	2	12	9	1	25(10.96)
Swelling (? 3 months)	1	2	10	9	1	23(10.09%)
Numbness (? March)	1	5	16	12	1	35(15.35%)
Needle infection	1	1	5	6	0	13(5.7%)
Broken needle	0	0	2	3	0	5(2.2%)
Loose connecting rod	1	2	6	8	0	17(7.5%)
osteomyelitis	0	0	1	0	0	1(0.4%)
Restricted knee movement	0	0	2	2	0	4(1.8%)
Nonunion of bone	0	0	1	1	0	2(0.9%)
Skin necrosis	0	1	0	1	0	2(0.9%)
Nerve injury	0	0	1	1	0	3(1.3%)
Ankle dislocation	0	0	2	1	0	3(1.3%)
Toe flexion deformity	1	3	5	4	0	13(5.7%)
recrudescence	0	1	1	1	0	3(1.3%)
Ankle arthritis	0	0	3	3	0	6(2.6%)
Subtalar joint stiffness	0	0	2	1	0	3(1.3%)

Figure 2: Table 1 :

**2**

age (year)	pain	swelling	numbness	needle infection	Loose connecting rod	toe flexion deformity
?7	0.83	0.83	0.83	8.3	8.3	8.3
8?17	3.85	3.85	9.62	1.9	3.8	5.8
18?30	16.67	13.89	22.22	6.9	8.7	6.9
31?59	20.45	20.45	27.27	13.6	18.2	9.1
?60?	50.00	50.00	50.00	0	0	0
r	0.93	0.926	0.976	-0.144	0.51	-0.585
p	?0.05	?0.05	?0.01	?0.05	?0.05	?0.05

Figure 3: Table 2 :

## 1 Conflicts of Interest

The authors declare that there are no conflicts of interest.

## 2 Data Availability

The data used to support the study are included in the paper.

[ Orthopedics () ] , *Orthopedics* 1991. 14 (3) p. .

[ Actaorthopscand () ] , *Actaorthopscand* . 1999. 70 p. .

[ J Pediatr Or-thop () ] , *J Pediatr Or-thop* 2007. 2 p. .

[ Chinese Journal of Orthopedics () ] , *Chinese Journal of Orthopedics* DOI: 10.3760 ?cma. j.. 0253-2352.2012.03.010. 2012. 32 (3) p. .

[ Chinese Journal of Orthopedics () ] , 10.3977/j.issn.1005-8478.2020.01.15. *Chinese Journal of Orthopedics* 2020. 28 (1) p. .

[Al-?aggadm ()] Al-?aggadm . *Talectomy in the treatment of resistant talipesquinovarus deformity: the indications and results?J*, 2013. 4 p. .

[Liu et al. (2020)] *Complications of bone transport technique using the Ilizarov method in the lower extremity: a retrospective analysis of 282 consecutive cases over 10 years. BMC MusculoskeletDisord*, Y Liu , M Yushan , Z Liu , J Liu , C Ma , A Yusufu . 10.1186/s12891-020-03335-w. 32505174. PMC7276072. 2020 Jun 6. 21 p. 354.

[Beaty (ed.) ()] *Congenital clubfoot (talipesquinovarus)*, J H Beaty . Canale ST (ed.) 2003. Philadelphia, PA: Mosby. p. . (Campbell’s operative orthopaedics. 10th ed.)

[Ferreira et al. ()] ‘Correction of neglected clubfoot using the Ilizarov external fixator’. R C Ferreira , M T Costo , G G Frizzo . *Foot AnkleInt* 2006. 4 p. .

[Ippolito et al.] ‘et al??ong-term comparative results in patients with congenital clubfoot treated with two different protocols’. E Ippolito , P Farsetti , R Caterini . *J Bone Joint Surg* 2003 (85) p. .

[Shaofeng and Sihe] *etc Analysis of complications of Ilizarov technique in the treatment of limb deformity*, Jiao Shaofeng , Qin Sihe .

[Mahan et al.] *Factors in pin tract infections*, J Mahan , D Seligson , S L ?enry .

[El-Sayed ()] ‘Ilizarov external fixation for management of severe relapsed clubfeet in older children’. M El-Sayed . *J Foot Ankle Surg?* 2013. 19 (3) p. .

[Gupta and Nitinbithar ()] ‘Ilizarov in relapsed clubfoot: a necessary evil’. Parmanand Gupta , Nitinbithar . 10.1097/BPB.0b013e32836486de. *Journal of Pediatric Orthopaedics B* 2013. 22 p. .

[Hosny (2020)] ‘Limb lengthening history, evolution, complications and current concepts’. G A Hosny . 32140790. PMC7058770. *J OrthopTraumatol* 2020 Mar 5. 21 (1) p. 3.

[Camathias and Valderrabano ()] ‘Oberli??Routine pin tract care in external fixation is unnecessary: A randomized, prospective, blinded controlled study’. C Camathias , V Valderrabano . DOI: 10.1016 / j.injury.2012.08.010?. *Injury* 2012. 43 (11) p. .

[Dobbs and R?schoenecker] ‘P??ong-term follow-up of patients with clubfeet treated with extended soft tissue release’. M B Dobbs , Nunley R?schoenecker . DOI: 10.2106 / JBJS.E.00114. *Bone Joint Surg Am?2006?88* (5) p. .

[Ahlborg and Josefsson] *Pin-tract complications in external fixation of fractures of the distal radius, ?* G Ahlborg , P O Josefsson .

[Paley (1990 r2501)] ‘Problems, obstacles, and complications of limb lengthening by the Ilizarovtechnique’. D Paley . *ClinOrthopRelat Res* 1990 r2501. p. .

[Professional Committee of Bone and Joint of China Rehabilitation Medical Association, China Association for Multidisciplinary *Professional Committee of Bone and Joint of China Rehabilitation Medical Association, China Association for Multidisciplinary Collaboration of Cerebral Palsy, and consensus of surgical treatment experts for spastic cerebral palsy*,

[Prem et al.] *Soft tissue Ilizarov correction of congenital talipes equinovarus 5to 10 years post surgery*, H Prem , M Zenios , Farrelr .

[Grill and Franke ()] ‘The Ilizarov distractor for the correction of relapsed or neglected clubfoot’. F Grill , J Franke . *J Bone Joint Surg Br* 1987. 69 p. .

[Freedman et al. ()] ‘The Ilizarov method for the treatment of resistant clubfoot: is it an effective solution?’. J A Freedman , H Watts , N Y Otsuka . *J PediatrOrthop* 2006. 26 p. . (11)

[?assan and ?etts] ‘The management of the neglected congenital foot deformity in the older child with the Taylor spatial frame’. A ?assan , M? ?etts . *J PediatrOrthop?2012?32* (1) p. .

## 15 CONCLUSION

---

- 250 [Bradish and Noor ()] 'TheIlizarov method in the management of relapsed club feet?J'. C F Bradish , S Noor .  
251 *J Bone Joint Surg( Br)* 2000. 82 (3) p. .
- 252 [Wallander HM? Congenital clubfoot? Aspects on epidemiology, residual deformity and patient reported outcome J]  
253 ctaOrthopSuppl?2010?81]b21 'Wallander HM? Congenital clubfoot? Aspects on epidemiology, residual  
254 deformity and patient reported outcome'. *J] ActaOrthopSuppl?2010?81* (339) p. .