



GLOBAL JOURNAL OF MEDICAL RESEARCH: J
DENTISTRY & OTOLARYNGOLOGY
Volume 23 Issue 3 Version 1.0 Year 2023
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Craniofacial Features of Cleidocranial Dysplasia (CCD) - A Case Report”

By Dr. B. V. Thimma Reddy, Dr. B. Udaykumar Chowdary,
Dr. Hemant Kumar. R, Dr. Raghavendra Kumar. J, Dr. N. Spandanareddy,
Dr. B. Simren & K. Sudheerkumar

Abstract- Cleidocranial dysplasia (CCD) is an autosomal-dominant malformation syndrome affecting bones and teeth. The most common skeletal and dental abnormalities in affected individuals are hypoplastic/aplastic clavicles, open fontanelles, short stature, retention of primary teeth, delayed eruption of permanent teeth, supernumerary teeth, and multiple impacted teeth. Treatment of CCD requires a multidisciplinary approach that may include dental corrections, orthognathic surgery and cranioplasty along with management of any complications of CCD. Early diagnosis of this condition enables application of the treatment strategy that provides the best quality of life to such patients. Notably, Runx2 gene mutations have been identified in CCD patients. Therefore, further elucidation of the molecular mechanism of supernumerary teeth formation related to Runx2 mutations may improve understanding of dental development in CCD. The insights into CCD pathogenesis may assist in the development of new treatments for CCD.

Keywords: *cleido cranial dysplasia, autosomal, runx 2 gene, mutations.*

GJMR-J Classification: *NLM Code: WN 430*



Strictly as per the compliance and regulations of:



© 2023. Dr. B. V. Thimma Reddy, Dr. B. Udaykumar Chowdary, Dr. Hemant Kumar. R, Dr. Raghavendra Kumar. J, Dr. N. Spandanareddy, Dr. B. Simren & K. Sudheerkumar. This research/review article is distributed under the terms of the Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0). You must give appropriate credit to authors and reference this article if parts of the article are reproduced in any manner. Applicable licensing terms are at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Craniofacial Features of Cleidocranial Dysplasia (CCD) – A Case Report

Dr. B. V. Thimma Reddy ^α, Dr. B. Udaykumar Chowdary ^σ, Dr. Hemant Kumar R. ^ρ,
Dr. Raghavendra Kumar. J ^ω, Dr. N. Spandanareddy [¥], Dr. B. Simren [§] & K. Sudheerkumar ^x

Abstract- Cleidocranial dysplasia (CCD) is an autosomal-dominant malformation syndrome affecting bones and teeth. The most common skeletal and dental abnormalities in affected individuals are hypoplastic/aplastic clavicles, open fontanelles, short stature, retention of primary teeth, delayed eruption of permanent teeth, supernumerary teeth, and multiple impacted teeth. Treatment of CCD requires a multidisciplinary approach that may include dental corrections, orthognathic surgery and cranioplasty along with management of any complications of CCD. Early diagnosis of this condition enables application of the treatment strategy that provides the best quality of life to such patients. Notably, Runx2 gene mutations have been identified in CCD patients. Therefore, further elucidation of the molecular mechanism of supernumerary teeth formation related to Runx2 mutations may improve understanding of dental development in CCD. The insights into CCD pathogenesis may assist in the development of new treatments for CCD.

Keywords: cleido cranial dysplasia, autosomal, runx 2 gene, mutations.

I. INTRODUCTION

The term cleidocranial dysplasia (CCD; OMIM 119600) is derived from the ancient Greek words cleido (collar bone), kranion (head), and dysplasia (abnormal formation). This rare hereditary skeletal disorder, which is also known as Scheuthauer-Marie-Sainton syndrome or cleidocranial dysostosis, is characterized by abnormal skeletal and dental development. The prevalence of CCD is an estimated one per million and does not differ by race or by gender.¹ In most cases, the disorder is an inherited autosomal dominant trait. In 20-40% of reported cases, however, the disorder occurs sporadically.¹ This syndrome is characterized by hypoplastic and/or aplastic clavicles, patent sutures and fontanelles, wormian bones, wide pubic symphysis, supernumerary teeth, short stature, and various other skeletal changes. Although clavicular defects have been reported in the literature as early as 1765,² Scheuthauer³ in 1871 was apparently the first to describe the syndrome accurately. Marie and Sainton⁴ in 1898 coined the term “dysostose cleidocranienne hereditaire” for this condition. The term

“cleidocranial dysostosis” was originally used because CCD was thought to involve only bones of intramembranous origin, i.e., bones of the skull, clavicles and flat bones. Subsequent studies showed that bones of endochondral ossification are also affected and that CCD is a generalized disorder of many skeletal structures. Therefore the term “cleidocranial dysostosis” was changed to “cleidocranial dysplasia” to reflect the more generalized nature of the condition.^{5,6}

II. CASE HISTORY

a) Clinical features

The clinical appearance of CCD is so distinct that it is pathognomonic. The main clinical features of CCD are recognized during early childhood and include a short stature, delayed closure of fontanelles, prominent forehead, and abnormal dental development. The head of a CCD patient usually shows frontal and parietal bossing and a groove along the metopic suture. The neck appears to be abnormally long, and the shoulders are narrow with marked drooping. (figure-1) Clavicular abnormalities with associated muscle defects allow excessive mobility of the shoulder girdle. For example, many CCD patients can approximate their shoulders in front of the chest for variable levels. The clinical spectrum is extremely variable even within families and ranges from mild cases with only dental abnormalities to severe cases with pronounced skeletal deformities.^{7,8}

b) Radiographic features

The distinctive radiological features of CCD are shortened or absent clavicles, delayed ossification of the skull bones and delayed ossification of pelvic bones.¹ The chest radiographs for CCD patients in show that the clavicles may be completely absent (aplasia) or smaller than normal (hypoplasia). The clavicles are typically hypoplastic or discontinuous, either unilaterally or bilaterally; the clavicles are completely absent in 10% of cases. Hypoplastic clavicles include hypoplasia of the acromial end or absence of the sternal end with the acromial end present. The missing segment may cause fibrous pseudoarthrosis or may be replaced by a fibrous tether or cord.⁹

c) Craniofacial Morphological Features

(Figure -1) shows that the skull in CCD is characterized by brachycephaly, delayed or failed

Author ^α ^σ ^ρ ^ω [¥] [§]: Department(s) and institution(s) Department of Paediatric and Preventive dentistry, SVS Institute of Dental Sciences, Appannapally, Mahbubnagar.

Corresponding Author^x: SVS institute of dental sciences.
e-mail: kothasudheerkumar266@gmail.com

closure of the fontanels, open skull sutures, and multiple wormian bones in the coronal and lambdoid suture regions. Defective fusion of frontal and parietal bones leading to open coronal and sagittal sutures are also visible. The nasal bones are missing or hypoplastic. Mandibular prognathism may be secondary to nasomaxillary deficiency. Dense alveolar crestal bone can be seen in the anterior mandible. Other craniofacial morphological features of CCD include abnormally small or non-existent maxillary sinuses, hypoplastic zygomatic bones, and patency of the mandibular symphysis.^{1,10} The zygomatic arch may be thin or even discontinuous at the zygomaticotemporal suture. The zygomatic arch has a characteristic downward bend.¹ The mandible is characterized by a narrow ascending ramus with nearly parallel anterior and posterior borders and by an abnormally slender and pointed coronoid process with an abnormally distal curvature.¹⁰ The trabecular pattern of the mandible is very coarse. (Figure-2)

d) Radiographic Features Associated with the Teeth

(Figure-2,3) shows that CCD is characterized by prolonged retention and delayed shedding of the primary teeth and multiple unerupted permanent and supernumerary teeth.¹⁰ Dentigerous cysts occasionally arise in association with these unerupted teeth. Although development of the primary teeth is rarely affected, root resorption and exfoliation of the primary teeth may be delayed. (figure-4)

e) Cone-Beam Computed Tomography (Cbct) Imaging

Other pertinent information provided by CBCT include the precise location of a supernumerary tooth in relation to important structures such as the cortex of the nasal floor, labial cortex of the nasal ridge, nasopalatine duct, and the mandibular canal and adjacent root apices.¹¹ Because CBCT clearly depicts the position and anatomy of impacted teeth, CBCT is useful for both diagnosis and treatment planning in CCD.

f) Histopathological Features

Tooth formation and eruption occur in a series of complex and highly regulated process. The reasons for failure of permanent tooth eruption and retention of the primary teeth in CCD patients are poorly understood. Absence of cellular cementum at the root apex is presumably one factor in failed or delayed eruption of permanent teeth and retention of the primary teeth in CCD.^{12,13} The lack of cellular cementum is presumed to increase the number of unerupted teeth in patients with CCD. However, recent reports of a lack of cellular cementum in normal teeth do not support this presumption.¹⁴⁻¹⁶ Studies of bone from the alveolus overlying unerupted teeth in CCD patients have a higher than normal density as well as reversal lines, which suggest an abnormal resorption pattern.^{17,18} Possible explanations for delayed eruption of teeth included increased density and coarse trabecular pattern of the

jaw bone, decreased resorption, and multiple reversal lines. A delayed eruption may also be attributable to various other factors such as mechanical obstruction of multiple supernumerary teeth. Therefore, the most likely causes of extreme delay or arrested eruption of permanent teeth in CCD are diminished bone resorption, delayed resorption of the roots of primary teeth, and less commonly multiple supernumerary teeth.¹⁹ One proposed hypothesis is that supernumerary tooth formation results from hyperactive dental lamina, i.e. over-proliferation or prolonged survival of dental lamina epithelial cells.²⁰ Another hypothesis is that formation of supernumerary permanent teeth in CCD patients results from markedly delayed resorption or from dental lamina of permanent dentition that is normal but does not resolve completely at the expected time.¹⁹

g) Molecular Genetics

The Runx2 gene is a master transcription factor of bone and plays a role in all stages of bone formation. Core binding factor (Cbf) plays crucial roles during skeletal development. Cbf consists of two subunits: Cbf alpha (Cbfa) and Cbf beta (Cbfb). Runt-related transcription factor 2 (Runx2) has been shown to be critical for the differentiation of osteoblasts and skeletal development.^{21,22} CCD results from a Runx2 gene mutation in the small arm of chromosome 6 at 6p21.1.^{23,24} A heterozygous mutation in the Runx2 gene encodes runt-related transcription factor 2, also termed core-binding factor alpha1 (CBFA1). Researchers generally agree that the underlying mechanism of CCD pathogenesis is haploin sufficiency or loss of Runx2 function.^{8,25} The Runx2 contains a DNA-binding domain (runt domain) which is necessary for transcriptional activation of target genes, a region of glutamine and alanine repeats in the N-terminal region (Q/A domain), and a region rich in proline-serine threonine (PST). The Runx2 is a key transcription factor involved in osteoblastic differentiation and skeletal morphogenesis.²⁶ Studies also suggest that Runx2 plays an important role in odontogenesis via participation in odontoblast differentiation, enamel organ formation, and dental lamina proliferation.²⁷ Disruption of these functions might explain the distinct dental anomalies associated with this disorder. To date, over 90 Runx2 gene mutations in 500 independent cases of CCD have been reported in the literature, including deletions, insertions, translocations, missense, frameshift, and splice mutations.²⁸ In most cases mutations occur in the runt domain.^{23,29} Mutations in Runx2 have a high penetrance and extreme variability. The Runx2 mutation is currently the only known molecular etiology of CCD. Notably, individuals who have CCD and identical Runx2 gene mutations show a wide variation in the number of asymmetrical supernumerary teeth in the maxilla and the mandible, which implies that the number and position of supernumerary teeth are not governed solely by Runx2

mutations. Runx2 mutations, which functions as a heterodimer with core binding factor b (Cbfb), are found in most individuals with CCD.^{21,22} Cbfb forms a heterodimer with Runx family proteins and enhances their DNA-binding capacity. Multiple functions of Cbfb are required for skeletal development and homeostasis in postnatal skeletogenesis. Cbfb deficiency reduced the expression of several key factors that mediate osteoblast formation and/or function. Cbfb is crucial for the later stages of chondrocyte differentiation as its deletion affects chondrocyte maturation and the formation of the growth plate. Although no Cbfb mutation has yet been identified in classical CCD patients, genetic alterations in the Cbfb gene may be responsible for CCD in those patients with no Runx2 mutation. Because Runx2 functions as a heterodimer with Cbfb, it has been suspected that Cbfb may be responsible for some cases of CCD. In terms of the pathogenesis of CCD, Cbfb deficiency may be equivalent to Runx2 haploinsufficiency as it relates to the function of the Runx2/ Cbfb complex in skeletogenesis.²¹ Fibroblast growth factor (FGF) signaling is one molecular mechanism of supernumerary teeth formation in CCD patients.³⁰ Runx2 might indirectly inhibit FGF signaling by antagonizing Twist1 function. Twist1 is a basic helix-loop-helix-containing transcription factor that is expressed in the dental mesenchyme in early stages of tooth development. A relative abundance of unbound Twist1 caused by Runx2 haploinsufficiency may elevate FGF signalling, which then causes formation of supernumerary teeth in human CCD.³⁰

h) Treatment

Managing the dental and orofacial manifestations of CCD is a challenging long-term process that requires careful planning and execution by an interdisciplinary team. The treatment strategies may differ according to the age of the patient. Surgical exposure of unerupted permanent teeth with orthodontic guided eruption is the preferred treatment for adolescent CCD patients. Generally, deciduous and supernumerary teeth should be removed to improve the possibility of orthodontic guided eruption.^{31,32} Bone overlying permanent teeth should also be removed since histology studies show that alveolar bone in CCD has abnormal dense trabeculation with multiple reversal lines.¹⁷ Orthodontic treatment with mini-implant screws for traction of impacted teeth can reduce the treatment time for CCD patients.³³ Leaving numerous deeply unerupted teeth in place is not an acceptable practice. The dentition associated with CCD is usually responsive to skillful orthodontic therapy and obviates the need for partial dentures. In adults with fully developed jaws, dental implants and fixed prostheses are the preferred therapeutic measures in adult CCD cases requiring multiple extractions of teeth. Calvarial defects in the

open anterior fontanelle, sagittal suture, and metopic suture have been successfully corrected by cranioplasty using bone cement.³⁴ Midface deficiency can be corrected by orthognathic surgery after growth is complete.^{35,36} In patients who meet the defined criteria, the above treatments can obtain substantial esthetic and functional benefits.

III. DISCUSSION

CCD is a generalized skeletal dysplasia affecting bones of intramembranous and endochondral ossification. The condition varies from mild cases presenting with only supernumerary teeth to cases with the phenotypic features that characterize CCD. Timely recognition of CCD and counseling for patients with hereditary risk factors are mandatory. Although CCD is associated with various skeletal abnormalities, CCD patients typically visit dental clinics only when they require treatment for dental and orofacial problems. Therefore, dentists have essential roles in identifying CCD and then planning and implementing a multidisciplinary therapeutic treatment aimed at improving quality of life in patients with this condition. Different approaches to the treatment of the dentition in CCD have been proposed in the past. The method suggested by Becker et al.^{31,32} may be viewed as the most promising.

The proposed method is founded on several premises:

- 1) Need for early removal of all obstacles to the eruption of the unerupted permanent teeth and application of traction forces at the biologically appropriate time,
- 2) Extraneous force needs to be provided to bring about an eruption of the teeth, along with an accompanying vertical alveolar development, and
- 3) Concentrating initial efforts towards bringing anterior teeth into mouth early, for the patient's psychological wellbeing.

Extract the anterior deciduous teeth and all supernumerary teeth, and expose unerupted permanent incisor teeth. The timing of surgical exposure of unerupted teeth is governed by appropriate root development. Root development should be two-thirds their expected length and is suitable for its active eruption. The approximately 3-year discrepancy in development of the dentition in these cases of dental age 7-8 years generally dictates that the chronological age of the patient is usually around 10-12 years.³¹ Further development of the roots of the posterior successional teeth will have increased their length to around two-thirds of the expected final length and are suitable for their active eruption at dental age of 10-11 years and chronological age 13 years. Surgical and orthodontic difficulties and complications abound during the treatment of CCD and there is a risk for the failure of one or other of the many aspects of the treatment or the

prognosis of the result. An inordinately long period involved in the completion of last orthodontic treatment stage.³² The displacement of the roots of several of the teeth is often extreme and many months of root torquing and uprighting are needed to bring them into their proper positions. Long-term retention of the treatment result is advised. FGF signalling is reportedly a molecular mechanism of supernumerary teeth formation in CCD patients.³⁰ However, wide variation in the dental phenotype of CCD patients suggests that genetic modifiers and interacting partners await discovery.³⁷ Twist1 is the functional antagonist of the Runx2. Excess of unbound Twist1 caused by Runx2 haploinsufficiency enhances FGF signaling, which then promotes formation of supernumerary teeth.³⁰ Runx2 haploinsufficiency in humans affects permanent dentition but not primary dentition.¹⁹ It is difficult to establish direct genotype-phenotype correlation for Runx2 because of very variable phenotypic penetrance of the mutations.³⁸ There is also a weak genotype-phenotype correlation in case of dental aspect of CCD phenotypes, especially with respect to teeth development.³⁹ Further elucidation of molecular mechanisms of supernumerary teeth formation related to Runx2 mutations will improve insight into dental development. The insights into CCD pathogenesis may assist in development of novel therapies for CCD.

REFERENCES RÉFÉRENCES REFERENCIAS

- McNamara CM, O'Riordan BC, Blake M, Sandy JR. Cleidocranial dysplasia: radiological appearances on dental panoramic radiography. *Dentomaxillofac Radiol* 1999; 28: 89-97.
- Martin S. Sur un déplacement naturel de la clavicule. *J Med Chir Pharmacol* 1765; 23: 456-60.
- Scheuthauer G. Kombination rudimentärer Schüsselbeine mit Anomalien des Schädels beim erwachsenen Menschen. *Allg Wien Med Ztg* 1871; 16: 293-5.
- Marie P, Sainton P. Sur la dysostose cleidocranienne héréditaire. *Rev Neurol* 1898; 6: 835-8.
- McKusick VA, Scott CI. A nomenclature for constitutional disorders of bone. *J Bone Jt Surg Am* 1971; 53A: 978-86.
- Rimoin DL. International nomenclature of constitutional diseases of bone. *J Pediatr* 1978; 93: 614-6.
- Chitayat D, Hodgkinson KA, Azouz EM. Intrafamilial variability in cleidocranial dysplasia: a three generation family. *Am J Med Genet* 1992; 42: 298-303.
- Quack I, Vonderstrass B, Stock M, et al. Mutation analysis of core binding factor A1 in patients with cleidocranial dysplasia. *Am J Hum Genet* 1999; 65: 1268-78.
- Mundlos S. Cleidocranial dysplasia: clinical and molecular genetics. *J Med Genet* 1999; 36: 177-82.
- Jensen BL, Kreiborg S. Craniofacial abnormalities in 52 schoolage and adult patients with cleidocranial dysplasia. *J Craniofac Genet Dev Biol* 1993; 13: 98-108.
- Dalessandri D, Laffranchi L, Tonni I, et al. Advantages of cone beam computed tomography (CBCT) in the orthodontic treatment planning of cleidocranial dysplasia patients: a case report. *Head Face Med* 2011; 7: 6.
- Rushton MA. An anomaly of cementum in cleidocranial dysostosis. *Br Dent J* 1956; 100: 81-3.
- Smith NH, Sydney NS. A histologic study of cementum in a case of cleidocranial dysostosis. *Oral Surg* 1968; 25: 470-8.
- Yamamoto H, Sakae T, Davies JE. Cleidocranial dysplasia: a light microscope, electron microscope, and crystallographic study. *Oral Surg Oral Med Oral Pathol* 1989; 68: 195-200.
- Counts AL, Rohrer MD, Prasad H, Bolen P. An assessment of root cementum in cleidocranial dysplasia. *Angle Orthod* 2001; 71: 293-8.
- Yang X, Zhang CY, Zheng SG. Analysis of root resorption and dental structure in patients with cleidocranial dysplasia. *J Peking Univ Heal Sci* 2011; 43: 98-101.
- Hitchin AD, Faily JM. Dental management in cleidocranial dysostosis. *Brit J Oral Surg* 1974; 12: 46-55.
- Kreiborg S, Jensen BL, Bjoerk A, Skieller V. Abnormalities of the cranial base in cleidocranial dysostosis. *Am J Orthod* 1981; 79: 549-57.
- Jensen BL, Kreiborg S. Development of the dentition in cleidocranial dysplasia. *J Oral Pathol Med* 1990; 19: 89-93.
- Jarvinen E, Tummers M, Thesleff I. The role of the dental lamina in mammalian tooth replacement. *J Exp Zool B Mol Dev Evol* 2009; 312B: 281-91.
- Chen W, Ma J, Zhu G, et al. Cbfb deletion in mice recapitulates cleidocranial dysplasia and reveals multiple functions of Cbfb required for skeletal development. *Proc Natl Acad Sci U. S. A* 2014; 111: 8482-7.
- Jaruga A, Hordyjewska E, Kandzierski G, Tylzanowski P. Cleidocranial dysplasia and Runx2-clinical phenotype-genotype correlation. *Clin Genet* 2016; 90: 393-402.
- Mundlos S, Mulliken JB, Abramson DL, Warman ML, Knoll JH, Olsen BR. Genetic mapping of cleidocranial dysplasia and evidence of a microdeletion in one family. *Hum Mol Genet* 1995; 4: 71-5.
- Lee B, Thirunavukkarasu K, Zhou L, et al. Missense mutations abolishing DNA binding of the osteoblast-specific transcription factor OSF2/CBFA1 in cleidocranial dysplasia. *Nat Genet* 1997; 16: 307-10.

25. Kim HJ, Nam SH, Kim HJ, et al. Four novel RUNX2 mutations including a splice donor site result in the cleidocranial dysplasia phenotype. *J Cell Physiol*2006; 207: 114-22.
26. Ducy P, Zhang R, Geoffroy V, Ridall AL, Karsenty G. *Osf2/Cbfa1* a transcriptional activator of osteoblast differentiation. *Cell* 1997; 89: 747-54.
27. Camilleri S, McDonald F. *Runx2* and dental development. *Eur J Oral Sci* 2006; 114: 361-73.
28. Shen Z, Zou CC, Yang RW, Zhao ZY. Cleidocranial dysplasia: report of 3 cases and literature review. *Clin Pediatr*2009; 48: 194-8.
29. Ducy P. *Cbfa1*: a molecular switch in osteoblast biology. *Dev Dyn* 2000; 219: 461-71.
30. Lu Y, Li Y, Cavender AC, Wang S, Mansukhani A, D'Souza RN. Molecular studies on the roles of *Runx2* and *Twist1* in regulating FGF signaling. *Dev Dyn* 2012; 241: 1708-15.
31. Becker A, Lustmann J, Shteyer A. Cleidocranial dysplasia: Part 1eGeneral principles of the orthodontic and surgical treatment modality. *Am J OrthodDentofacOrthop*1997; 111: 28-33.
32. Becker A, Shteyer A, Bimstein E, Lustmann J. Cleidocranial dysplasia: Part 2 General principles of the orthodontic and surgical treatment modality. *Am J OrthodDentofacOrthop* 1997; 111:173-83.
33. Kuroda S, Yanagita T, Kyung HM, Takano-Yamamoto T. Titanium screw anchorage for traction of many impacted teeth in a patient with cleidocranial dysplasia. *Am J OrthodDentofacOrthop*2007; 131: 666-9.
34. Kang N, Kim SZ, Jung SN. Correction of depressed forehead with bone source in cleidocranial dysplasia. *J CraniofacSurg*2009; 20: 564-6.
35. Hwang SM, Park B, Hwang MK, Kim MW, Lee JS. Aesthetic facial correction of cleidocranial dysplasia. *Arch CraniofacSurg*2016; 17: 82-5.
36. Madeira MF, Caetano IM, Dias-Ribeiro E, et al. Orthognathic surgery in patients with cleidocranial dysplasia. *J CraniofacSurg*2015; 26: 792-5.
37. Lee KE, Seymen F, Ko J, et al. RUNX2 mutations in cleidocranial dysplasia. *Genet Mol Res* 2013; 12: 4567-74.
38. Yoshida T, Kanegane H, Osato M, et al. Functional analysis of RUNX2 mutations in Japanese patients with cleidocranial dysplasia demonstrates novel genotype-phenotype correlations. *Am J Hum Genet* 2002; 71: 24-38.
39. Ryoo HM, Kang HY, Lee SK, et al. RUNX2 mutations in cleidocranial dysplasia patients. *Oral Dis* 2010; 16: 55-60.