



GLOBAL JOURNAL OF MEDICAL RESEARCH: C
MICROBIOLOGY AND PATHOLOGY
Volume 19 Issue 1 Version 1.0 Year 2019
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Distribution of Genital Mycoplasmas in pregnant and non-pregnant Women in Iraq of Basrah City

By Reham M. Al-Mosawi, Amin A. Al- Sulami & Ghaeda. J. Al-Ghizawi

Basrah University

Abstract- A total of 120 women attending the outpatient clinic of the obstetric and gynecology department of Basrah General Hospital, were evaluated for the role of genital mycoplasmas in certain infectious female genital conditions during the period from December, 1st 2003 to October, 2nd 2004. High vagina and endocervix swabs were cultured. Five species of genital mycoplasmas were isolated and identified on modified PPLO medium, the characteristics and biochemical properties of them were fit with those of *Mycoplasma hominis*, *Ureaplasma urealyticum*, *M. fermentans*, *M. genitalium* and *M. penetrans* in which the last three species were identified for the first time in Iraq. *M. hominis*, *U. urealyticum*, *M. genitalium* and *M. penetrans* were more frequently distributed in age group 20-39 years, and in women with lower socioeconomic status. A significant difference was found in the isolation of both *M. hominis* and *M. genitalium* from non-pregnant. Also, a significant difference was noted in isolation rates of *M. hominis*, *M. fermentans* and *M. penetrans* from pregnant women at different gestational periods..

GJMR-C Classification: LCC: QH301



Strictly as per the compliance and regulations of:



© 2019. Reham M. Al-Mosawi, Amin A. Al- Sulami & Ghaeda. J. Al-Ghizawi. This research/review article is distributed under the terms of the Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0). You must give appropriate credit to authors and reference this article if parts of the article are reproduced in any manner. Applicable licensing terms are at <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

Distribution of Genital Mycoplasmas in pregnant and non-pregnant Women in Iraq of Basrah City

Reham M. Al-Mosawi ^α, Amin A. Al- Sulami ^σ & Ghaeda. J. Al-Ghizawi ^ρ

Abstract- A total of 120 women attending the outpatient clinic of the obstetric and gynecology department of Basrah General Hospital, were evaluated for the role of genital mycoplasmas in certain infectious female genital conditions during the period from December, 1st 2003 to October, 2nd 2004. High vagina and endocervix swabs were cultured. Five species of genital mycoplasmas were isolated and identified on modified PPLO medium, the characteristics and biochemical properties of them were fit with those of *Mycoplasma hominis*, *Ureaplasma urealyticum*, *M. fermentans*, *M.genitalium* and *M.penetans* in which the last three species were identified for the first time in Iraq. *M.hominis*, *U.urealyticum*, *M. genitalium* and *M.penetans* were more frequently distributed in age group 20-39 years, and in women with lower socioeconomic status. A significant difference was found in the isolation of both *M.hominis* and *M. genitalium* from non-pregnant. Also, a significant difference was noted in isolation rates of *M.hominis*, *M.fermentans* and *M.penetans* from pregnant women at different gestational periods. A statistically significant difference at the level of ($P<0.01$) was found in isolation of both *U.urealyticum* and *M.genitalium* from endocervix region. This study revealed a significant association between the five species of genital mycoplasmas and some cases or symptoms in the female genital tract. Furthermore, different species of bacterial isolates other than mycoplasmas were identified in 12 infected women using contraception. Genital mycoplasmas were found as a single infection in frequent (16.9%) and mixed infection with other causative agents in frequent (17.5%). Also, this study shows the conjunction of *M.hominis* with both *U.urealyticum* and *M.fermentans* in (5.0%), (1.6%) respectively.

I. INTRODUCTION

The mycoplasmas are the smallest cell free-life microorganisms, characterized by their small cell size (0.3-0.8 μ m) and thus can pass through some filters used to remove bacteria. They have the smallest genome size and, as a result, lack many metabolic pathways and lack of a rigid bacterial cell wall (Mayer and Murray *etal.*, 2002). Mycoplasmas are fastidious organisms that require specialized media for their isolation and identification (Razin and Freundt, 1984; Razin *etal.*, 1998). Ultrastructural studies showed that they are constructed of only three organelles: plasma membrane, ribosomes and prokaryotic chromosomes. There is no evidence of any intracellular membrane structure, and because of the lack of rigid cell wall,

mycoplasmas are pleomorphic (Lin, 1985). The family Mycoplasmataceae contains two genera that infect humans: Mycoplasma and Ureaplasma, which are usually referred to collectively as mycoplasmas. Seven species of mycoplasmas can be isolated from genitourinary tract, but in female genitourinary tract the most common mycoplasmas isolated are *U. urealyticum* and *M.hominis* have been implicated in human diseases (Cassell and Cole, 1981; Krause and Taylor-Robinson, 1992). *M.hominis* and *U.urealyticum* normally inhabit the urogenital tract in sexually mature men and women. Higher isolation rates are obtained from women than from men, possibly reflecting more favorable growth conditions in the vaginal tract. The degree of colonization with these species is related to sexual activity and to the number of sexual partners (Taylor-Robinson and McCormack, 1979). In 1980 new mycoplasma species were isolated from urethral specimens from 2 of 13 men with nongonococcal urethritis. It was later named *Mycoplasma genitalium* because of the host tissue location, however *M.genitalium* strains were originally isolated from the urogenital tract. This mycoplasma shared several properties with *M.pneumoniae* (Tully *et al.*, 1981,1983; Lind *et al.*, 1984; Taylor-Robinson, 1995).

M.genitalium and *M.penetans* are recent additions to the pathogenic human mycoplasmas, reviewed by Marmion and Harris, (1996). *M.penetans*. This newly-identified mycoplasma was isolated by (Lo *etal.*, 1991) from the urogenital tract of Human Immuno Virus of positive homosexual men. In a study in west Africa, *M.genitalium* was associated with nongonococcal urethritis (NGU), particularly in Trichomonas vaginalis negative patients (Pepin *et al.*, 2001).

Preliminary evidence for the involvement of the mycoplasma in the cervicitis and pelvic inflammatory disease was presented by Moller *etal.* (1984) and Uno *etal.* (1997). The etiological roles of genital mycoplasmas are: Acute pyelonephritis, bacterial vaginitis, pelvic inflammatory disease, chorioamnionitis, post- abortion and postpartum fever, pneumonia in newborns, non gonococcal urethritis, prostatitis and epididymitis (Mardh and Westrom, 1970; Shepard, 1970, 1980; Taylor-Robinson and McCormack, 1980; Krause and Taylor-Robinson, 1992; Taylor-Robinson, 1996). Furthermore, that mycoplasmas play a role in human infertility (Gnarpe and Fribreg, 1972; Stray-

Author α σ ρ : Lecturer, MSC Microbiology/Medical bacteriology, Dentistry College, Basic Science, Basrah University, Basrah, Iraq.
e-mail: rehamalmosawiaz13@gmail.com

Pedersen et al., 1978; Cassel et al., 1983; Toth et al., 1983), abortion and stillbirth (Embree et al., 1980; Kunds in et al., 1981), premature birth and low birth weight of infant (Braun et al., 1971; Embree et al., 1980; Kunds in et al., 1981).

Despite numerous studies of these organisms, their role in most genital infection remains controversial and ill defined. The ubiquity and low virulence of genital mycoplasmas make it difficult to evaluate their role in producing genital infections, a situation that is further complicated by their frequent isolation together with other sexually transmitted diseases (Watts and Eschenbach, 1988). Furthermore the presence of both genital mycoplasmas (*U.urealyticum* and *M.hominis*) in a large proportion of healthy women complicates the assessment of the pathogenic role of these organisms (Taylor-Robinson and Furr, 1998). However, circumstantial evidence including high frequency of isolation of mycoplasmas from females with urogenital conditions or infertility warns against minimizing the importance of these microorganisms in human disease (Kunds in and Driscoll, 1970).

Mycoplasmas were the causative agents of disease in human (Krause and Taylor-Robinson, 1992) over many years passed. Despite, numerous studies conducted in Iraq on different genera of bacteria, there have been few studies on the role of genital mycoplasmas in the infectious disease (Simhairi, 1990; Al-Bahli, 1993).

II. MATERIALS AND METHODS

a) Study population

The population under study was women attending the outpatient clinic of obstetric and gynecology department of Basrah General Hospital. The samples obtained consisted of pregnant women at different gestational periods and non pregnant women including women using contraceptive devices besides women not using contraceptive devices and those with infertility and those with various complaints. Also women presented with different types of abortions are included. The samples were selected randomly from the women above and collected from different parts of Basrah. A total number of two hundred and twenty two women (222) were investigated. Their age ranged from 17 to 50 years. The investigation period extended from 1st of December, 2003 to 2nd of October, 2004. A control group consisted of one hundred (100) healthy women.

b) Collection and Inoculation of samples

Two swabs one from endocervix and the other from high vagina were obtained from each woman and each was inoculated. Onto a suitable medium. A sterile speculum was used. All swabs were transported to the laboratory within (one) hour for culture. For the isolation of mycoplasmas, each specimen was directly inoculated into the liquid phase, mixed up well and tilted

for a while, once or twice, to cover the upper slanted portion in a simple monophasic - diphasic culture setup (MDCS) prior to incubation (Al-Sulami et al., 2002). For the isolation of bacteria other than mycoplasmas another two swabs from endocervix and high vagina were obtained from the same women included in the study. Then, each specimen after being transported to the laboratory was directly cultured onto MacConkey & Blood agar by the streaking method then incubation follows.

c) Culture Media

Media for isolation and identification species of genital mycoplasmas were applied depending on Marmion and Harris [5].

d) Cultivation and isolation of mycoplasmas (genital mycoplasmas)

Endocervix and high vaginal swabs were taken from the women under study, then, each specimen was directly inoculated into the liquid phase of the MDCS, mixed up well and then tilted once or twice to cover the upper portion of the slant for a while prior to incubation. All inoculated media were incubated aerobically at 37°C and observed daily for colour change from red to yellow in the liquid phase after 24 hrs. Then isolated colonies appeared after that on the slanted solid phase. We added sheep erythrocytes (7%) to the solid phase of the MDCS in several trials to detect the blood haemolysis by *M.fermentans* and *U.urealyticum*. as well, the egg yolk suspension (15 ml) was added to the standard PPLO agar for detect the lipolytic ability of *M.fermentans*.

III. RESULTS

In our knowledge this study performs the first isolation of *M.fermentans*, *M.genitalium* and *M.penetrans* in Iraq with MDCS system and modified PPLO media according to the previous studies. Besides, it is the first isolation to both species: *M.hominis* and *U.urealyticum* by this method. The colonial growth was observed on the upper portion of the slant together with changing in colour of the liquid phase from red to yellow after 24 hrs. However, approximately 96 hrs. Period was necessary for full development of colonies on modified media of PPLO with MDCS for show the fried-egg appearance Figure 1, 2. Distribution of genital mycoplasmas and occupation is shown in Table (1), there was no statistically significant difference for *M.hominis*, *M.fermentans*, *U.urealyticum*, *M.penetrans* but a difference was found to be just significant for *M.genitalium* ($X^2 = 11.333$; $P < 0.01$) when comparison between the two groups (housewife and employee). Concerning the usage of contraceptives (Table 2), a difference was found to be just significant for *M.fermentans* $P < 0.05$ and *M.genitalium* $P < 0.05$, in detection of genital mycoplasmas colonization in tested women under study that used and not used of

contraceptive devices. A difference was found to be significant in the isolation of both *M.hominis* and *M.genitalium* from non-pregnant when compared with pregnant women. A significant difference was also noted in the isolation rates of: *M.hominis*, *M.fermentans* and *M.penetrans* from pregnant women at different gestational periods (trimesters) as shown in Table 3, 4.

A statistically significant difference at the level of (P<0.01) was noted in the isolation of both *U.urealyticum* and *M.genitalium* from endocervix region

in comparison with high vaginal region as shown in Figure 3. Also, the bacteria (other than mycoplasmas) which isolated from women using contraception devices, they consisted of *S.aureus* as most frequently bacterial isolated in 20.0% followed by *K.pneumoniae*, *E.coli*, *P.aeruginosa* and *Proteus spp.* in (15.0%, 10.0%, 10.0%, 5.0%) respectively as shown in Table 5. Further more, the genital mycoplasmas were found as a single infection in 20 cases (16.9%) and mixed infection with other causative agents in 21 cases (17.5%) Table 6.




Figure 1: The fried egg colonies of *M.hominison* modified PPLO medium (X 160)




Figure 2: The fried egg colonies of *M.genitalium* on modified PPLO medium (X 160)

Table 1: Distribution of genital mycoplasmas in relation to occupation

Occupation	No. of tested women	No. and (%) of women + ve in				
		<i>M. hominis</i>	<i>M.fermentans</i>	<i>U.urealyticum</i>	<i>M.genitalium</i>	<i>M.penetrans</i>
House wife	90	12 (13.3)	4 (4.4)	9 (10.0)	5 (5.5)	3 (3.3)
Employee	30	4 (13.3)	1 (3.3)	3 (10.0)	0 (0)	0 (0)
Total	120	16	5	12	5	3
X ²		0	0.286	0	11.333	3.000
P		NS	NS	NS	0.01	NS

P> 0.05 (NS)

Table 2: Genital mycoplasmas colonization correlated with usage of contraception

Women subjects	No. of tested women	No. and (%) of women + ve in				
		<i>M. hominis</i>	<i>M. fermentans</i>	<i>U. urealyticum</i>	<i>M. genitalium</i>	<i>M. penetrans</i>
Used	20	3 (15.5)	0 (0)	2 (10.0)	0 (0)	1 (5.0)
Not used	100	13 (13.0)	5 (5.0)	10 (10.0)	5 (5.0)	2 (2.0)
Total	120	16	5	12	5	3
X ²		0.143	6.250	0	6.250	1.286
P		NS	0.05	NS	0.05	NS

P > 0.05 (NS)

Table 3: Prevalence of Genital Mycoplasmas in Pregnants Compared to Non-pregnant Women

Women subjects	No. tested	No. and (%) of women + ve in				
		<i>M. hominis</i>	<i>M. fermentans</i>	<i>U. urealyticum</i>	<i>M. genitalium</i>	<i>M. penetrans</i>
Pregnant	40	3 (7.5)	1 (2.5)	4 (10.0)	0 (0)	2 (5.0)
Non-pregnant	50	8 (16.0)	2 (4.0)	5 (10.0)	2 (4.0)	1 (2.0)
X ²		12.040	0.346	0	4.000	2.25
P		0.01	NS	NS	0.05	NS

Table 4: Prevalence of Genital Mycoplasmas in Pregnant Women at Different Gestational Periods

Pregnant women	No. tested	No. and (%) of women + ve in					Total of G.M (%)
		<i>M. hominis</i>	<i>M. fermentans</i>	<i>U. urealyticum</i>	<i>M. genitalium</i>	<i>M. penetrans</i>	
First trimester	10	1 (10.0)	1 (10.0)	1 (10.0)	0 (0)	1 (10.0)	4 (40.0)
Second trimester	10	0 (0)	0 (0)	1 (10.0)	0 (0)	0 (0)	1 (10.0)
Third trimester	20	2 (10.0)	0 (0)	2 (10.0)	0 (0)	1 (5.0)	5 (25.0)
X ²		10.110	23.333	0	0	10.000	34.000
P		0.01	0.01	NS	NS	0.01	0.01




Figure 3: Correlation between genital mycoplasmas and sources of isolation

Table 5: Bacteria which isolated from women using contraception (20 women).

Bacteria	No. and (%) of infected women + ve in
<i>Staphylococcus aureus</i>	4 (20.0)
<i>Klebsiella pneumoniae</i>	3 (15.0)
<i>Escherichia coli</i>	2 (10.0)
<i>Pseudomonas aeruginosa</i>	2 (10.0)
<i>Proteus spp.</i>	1 (5.0)
Total	12 (60.0)
X ²	6.84
P	0.01

Table 6: Presence of genital mycoplasmas alone or in conjunction with other bacteria

Total of genital mycoplasmas	Alone	In conjunction with				
		<i>S.aureus</i>	<i>K.pneumoniae</i>	<i>P.aeruginosa</i>	<i>E.coli</i>	<i>Proteus. spp</i>
41	20	7	5	4	3	2

IV. DISCUSSION

Prevalence of mycoplasmas in the female genital tract depends on numerous factors such as age, level of socioeconomic status, sexually active, disorders in the menstruation, pregnancy, infertility, urogenital complaints and use of contraceptive device [19-21, 38-39]. According to our knowledge, this study was the third in Iraq to determine the prevalence of genital mycoplasmas in Iraqi women and the second in Basrah. Further, the current study shows for the first time the isolation and identification of: *M.fermentans*, *M.genitalium* and *M.pneumoniae* in Iraq besides the first isolation of both *M.hominis* and *U.urealyticum* by using modified PPLO medium and MDCS system for the primary isolation of genital mycoplasmas from clinical samples then, detection with biochemical test [18, Holt et al, 1994]. The modified PPLO medium consist of antimicrobial agents: such as thallium acetate and penicillin, the other supplements consist of glucose solution; sodium deoxyribonucleate (DNA calf thymus) solution; hydrogen phosphate solution and cresol red solution as an indicator of the mycoplasmal growth in addition to PPLO broth/agar media; yeast extract; sodium chloride and horse serum, this method offered multi merit symbolized by the less contamination as a

result of elimination of transport media, fast of results appearance, supply a pliability in the kinds of media represented by liquid and solid phases, it is inexpensive because its consumed a small quantities of liquid and solid media furthermore, it has a short incubation time period within one test tube. It is the first technique used to determine the prevalence of genital mycoplasmas in women in Basrah City in comparison with studies of Simhairi (1990 and Al-Bahlis' (1993).

Table 1 shown, there was no statistically significant difference for *M.hominis*, *M.fermentans*, *U.urealyticum*, *M.pneumoniae* but a difference was found to be just significant for *M.genitalium*($P < 0.05$), when comparison between the two groups (housewife and employee) in related of occupation, this could be association with educational levels of women, awareness, culture and interest in personal hygiene [21, 26]. Concerning the usage of contraceptives (Table 2) which shows, a significant difference for *M.fermentans*($P < 0.05$) and *M.genitalium* ($P < 0.05$) in detection of genital mycoplasmas colonization when, tested women under study that used and not used of contraceptive devices, that is due to, the colonization of genital mycoplasmas in women is linked to younger age, contraceptive devices use, lower socioeconomic

status, and sexual activity with multiple partners and other factors [19, 21]. A difference was found to be significant in the isolation of both *M.hominis* and *M.genitalium* from non-pregnant when compared with pregnant women. A significant difference was also noted in the isolation rates of: *M.hominis*, *M.fermentans* and *M.penetrans* from pregnant women at different gestational periods (trimesters) as shown in Table 3, 4. Simhairi (1990), Al-Bahli (1993) found no significant difference in the prevalence of *M.hominis* and *U.urealyticum* among pregnant women as compared to non pregnant, as well as, to gestational stage of pregnancy. Their results were somewhat consistent with the results of the study. Similar as well as contrary observations regarding the prevalence of *M.hominis* of other strains in pregnant women have been reported by many investigators (Csonka et al., 1966; Jones, 1967; Harwick et al., 1970; Delouvois et al., 1975; Taylor-Robinson and McCormack, 1979 and Iwaska et al., 1986a). This controversy may be attributed to several factors like strain variations of the microorganisms, difficulty in determining a precise matching control with respect to sexual experience, especially in relation to the number of partners and population peculiarities, like differences in the socioeconomic status and hygienic standard (McCormack et al., 1973b). In present study we found, a statistically significant difference at the level of ($P < 0.01$) in the isolation of both *U.urealyticum* and *M.genitalium* from endocervix region in comparison with high vaginal region (Figure 3), the results of present study in agreement with R.M. Al-Mosawi, 2009. And in study of Upadhyaya et al. (1983) the frequency of the isolation of *U.urealyticum* was significantly higher in the infertile group than in a group of pregnant women. Thereupon, the results of the current study which show that both *U.urealyticum* and *M.genitalium* were recovered in a higher isolation rates from endocervix than high vagina of infected women may be associated with the inflammatory disease leading to infertility. Several investigators have demonstrated some kind of association between female infertility and *U.urealyticum* infection (Gnarpe and Friberg, 1973a; Fowlkerson et al., 1975; Upadhyaya et al., 1983; Taylor-Robinson, 1986).

In present study, the bacteria (other than mycoplasmas) were isolated from women using contraceptions, they consisted of *S.aureus* as most frequently bacterial isolated in 20.0% followed by *K.pneumoniae*, *E.coli*, *P.aeruginosa* and *Proteus spp.* in (15.0%, 10.0%, 10.0%, 5.0%) respectively as shown in Table 5, those findings were in accordance with Al-Bahli, 1993 study.

Moreover, in the current study, the genital mycoplasmas were found as a single infection in 20 cases (16.9%) and mixed infection with other causative agents in 21 cases (17.5%), as showed in Table 6, these results were somewhat in agreement with Simhari, 1990; Al-Bahli, 1993; Al-Mosawi, 2009.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Mayer, G. and Murray, G. M., et al. (2002). Mycoplasma and Ureaplasma. Medical Microbiology, 3rd ed., Chapter 42., MBIM 650/720, South Carolina University.
2. Razin, S. and Freundt, E. A. (1984). The mycoplasmas. In: Krieg, R. N. & Holt, G. J. (ed.). Bergey's Manual of Systematic Bacteriology. Vol. 1: Williams & Wilkins, C. O. Baltimore London.
3. Razin, S., Yogeve, D., Naot, Y. (1998). Molecular biology and pathogenicity of mycoplasmas. Microbiol. Mol. Biol Rev. 62: 1094-1156.
4. Lin, J. L. (1985). Human macoplasma infection: Serologic observations. Rev. Infect. Dis. 2: 216-30.
5. Cassell, G. H. and Cole, B. C. (1981). Mycoplasmas as agents of human disease. N. Engl. J. Med. 2: 80-87.
6. Krause, D. and Taylor-Robinson, D., Elhane, R. Mc., Finch, L., Baseman, J., Maniloff, J., Washington, D. C. (1992). Mycoplasmas which infect humans "Molecular biology and pathogenesis" American Society for Microbiology., PP. 417-444.
7. Taylor-Robinson, D. and McCormack, W. M. (1979). Mycoplasmas in human genitourinary infections. In: The Mycoplasmas. Vol. II. Academic Press Inc. Orlando, Florida 32887 USA: Page 307.
8. Tully, J. G., Taylor-Robinson, D., Cole, R. M. and Rose, D. L. (1981). A newly discovered mycoplasma in the human urogenital tract. Lancet. i: 1288-1291.
9. Tully, J. G., Taylor-Robinson, D., Rose, D. L., Cole, R. M. and Bove, J. M. (1983). Mycoplasma genitalium, a new species from the human urogenital tract. Int. J. Syst. Bacteriol. 33: 387-396.
10. Lind, K., Lindhardt, B. Q., Schutten, H. J., Blom, J. and Christiansen, C. (1984). Serological cross-reactions between Mycoplasma genitalium and Mycoplasma pneumoniae. J. Clin. Microbiol. 20: 1036-1043.
11. Taylor-Robinson, D., (1995). The history and role of Mycoplasma genitalium in sexually transmitted diseases. Genito-urin. Med. 71: 1-8.
12. Marmion, B. and Harris, R. (1996). Mycoplasma pneumoniae and other medically important members of the family Mycoplasmataceae In: Mackie & Macroney Practical medical microbiology, Vol. 30, PP. 591-595; 600-601, Collee, J., Marmion, B., Fraser, A. and Simmonds, A., (Eds.), 14th ed. Churchill Livingstone, New York.
13. LO, S. C., Tsai, S., Benish, J. R., Shih, J. W., Wear, D. J. and Wong, D. M. (1991). Enhancement of HIV-1 cytotoxic effects in CD₄⁺ lymphocytes by the AIDS-associated mycoplasma. Science, 251: 1074-1076.
14. Pepin, J., Sobela, F., Deslandes, S., et al. (2001). Etiology of urethral discharge in West Africa: the role

- of Mycoplasma genitalium and Trichomonas vaginalis. Bull. World Health Organ. 79: 118-26.
15. Moller, B. R., Allen, J., Toft, B., Hansen, K. B. and Taylor-Robinson, D. (1984). Pelvic inflammatory disease after hysterosalpingography associated with Chlamydia trachomatis and Mycoplasma hominis. Br. J. Obstet. Gynecol. 91: 1181-7.
 16. Uno, M., Deguchi, T., Saito, A., Yasuda, M., Komeda, H., Kawada, Y. (1997). Prevalence of Mycoplasma genitalium in asymptomatic men in Japan. Int. J. STD-AIDS. University Gifu, Japan. 8 (4): 259-60.
 17. Mardh, P.A. and Westrom, L. T. (1970). T-mycoplasma in the genitourinary tract of the female. Microbiol. Scand. Sect. B. 78: 367-374.
 18. Shepard, M. C. (1970). Nongonococcal urethritis associated with human strains of T-mycoplasma. JAMA. 211: 1335-40.
 19. Shepard, M. C. (1980). Current status of Ureaplasma urealyticum in nongonococcal urethritis. Asean. J. Clin. Sci. 1: 198-209.
 20. Taylor-Robinson, D. and McCormack, W. M. (1980). The genital mycoplasma. Part II. N. Engl. J. Med. 320: 1003-1063.
 21. Krause, D. and Taylor-Robinson, D., Elhaney, R. Mc., Finch, L., Baseman, J., Maniloff, J., Washington, D. C. (1992). Mycoplasmas which infect humans "Molecular biology and pathogenesis" American Society for Microbiology., PP. 417-444.
 22. Taylor-Robinson, D. (1996). Infections due to species of Mycoplasma and Ureaplasma: an update. Clin. Infect. Dis. 23: 671-682.
 23. Gnärpe, H. and Friberg, J. (1972). Mycoplasmas and Human reproductive failure. Am. J. Obstet. Gynecol. 114: 727-731.
 24. Stray-Pedersen, B., Erg, J., and Reikram, T. M. (1978). Uterine T-mycoplasma colonization in reproductive failure. Am. J. Obstet. Gynecol. 3: 306-11.
 25. Cassell, G. M., Younger, J. B., Brown, M. B., Blackwell, R. E., Davis, J. K., Marriott, P. A. and Stagno, S. (1983). Microbiologic study of infertile women at the time of diagnostic laparoscopy: association of Ureaplasma urealyticum with defined sub population. N. Engl. J. Med. 308: 502-4.
 26. Toth, A., Lesser, M. L., Brooks, G. and Labriola, D. (1983). Subsequent pregnancies among 161 couples treated for T-mycoplasma genital tract infection. N. Engl. J. Med. 308: 505-7.
 27. Embree, J. E., Krause, V. W., Embil, J. A. and Macdonald, S. (1980). Placental infection with Mycoplasma hominis and Ureaplasma urealyticum: Clinical correlation. Obstet. Gynecol. 56: 475-81.
 28. Kundsinn, R. B., Driscoll, S. G. and Pelletier, P. A. (1981). Ureaplasma urealyticum incriminated in perinatal morbidity and mortality. Science, 213: 474-6.
 29. Braun, P., Lee, Y. H., Klein, J. O., Marcy, S. M., Klein, T. A., Charles, D., Levy, P. and Kass, E. H. (1971). Birth weight and genital mycoplasmas in pregnancy. N. Engl. J. Med. 284: 167-171.
 30. Watts, D.H. and Eschenbach, D. A. (1988). Treatment of chlamydia, mycoplasma and group B-Streptococcal infections. Clin. Obstet. Gynecol. 2: 435-52.
 31. Taylor-Robinson, D. and Furr, P. M. (1998). Update on sexually transmitted mycoplasmas. Lancet. 351 (Suppl 3): 12-15.
 32. Kundsinn, R. B. and Driscoll, S. G. (1970). Mycoplasmas and human reproductive failure. Surg. Gynecol. Obstet. 131: 89-92.
 33. Simhairi, R. Z. (1990). Microbiological study of some mycoplasma species of female genitalia and evaluation of their role in some disease conditions. M. Sc. thesis, Al-Mustansiriyah University. (Arabic).
 34. Al-Bahli, S. A. H. (1993). Prevalence of genital Mycoplasma in women with selected obstetric and gynecological conditions. M. Sc. thesis, College of Medicine, Basrah University. (Arabic).
 35. Al-Sulami, A. A., Hammadi, S. S. and Al-Gizawi, G. J. (2002). A single method for rapid isolation and identification of Mycoplasma pneumoniae from clinical specimens. Eastern Mediterranean Health Journal, 8 (1): 1-7. (Arabic).
 36. McCormack, W. M., Rosner, B. and Lee, Y. H. (1973b). Colonization with genital Mycoplasma in women. Am. J. Epidemiol. 97: 240-245.
 37. McCormack, W. M. (1983). Epidemiology of Mycoplasma hominis. Sexually Transmitted Diseases. J. Amer. Ven. Dis. Assoc. 10: 261-262.
 38. Al-Mosawi, R.M. (2009). Signs and symptoms of urethritis and cervicitis among women with or without genital mycoplasma infection in governorate of Basrah. Journal of Basrah Researches ((Sciences)) Vol. 35, No.3, p: 48-57.
 39. Al-Sulami, A., Al-Gizawi, G. J. and Al-Mosawi, R.M. (2007). Prevalence of genital mycoplasmas among women in the province of Basrah and an evaluation of their role in some cases. J. Basrah Researches (Sciences) Vol. 33. No.2, p: 129 -136.
 40. Holt, J. G., Krieg, N. R., Sneath, P. H., Staley, J. T. and Williams, S. T. (1994). The Mycoplasmas or (Mollicutes): cell wall-less bacteria In: Bergey's Manual of determinative bacteriology, Vol. 30, PP. 705-707, 9th ed. Williams & Wilkins, U.S.A.
 41. C. R. Lilia., G. Constantino., Z. Ivonne., Y. Antonio., and G. Silvia. Association of Mycoplasma hominis and Ureaplasma urealyticum with some indicators of

- nonspecific vaginitis. *Revista Latinoamericana de Microbiologia*. 42: 1-6. 2000.
42. Csonka, G. W. and Williams, R. E. O. (1966). T-strain mycoplasma in non-gonococcal urethritis. *Lancet*, 1: 1292-6.
 43. Jones, D. M. (1967a). Mycoplasma hominis in pregnancy. *J. Clin. Pathol.* 20: 633-635.
 44. Harwick, H. J., Purcell, R. H., Lippa, J. B. and Felkety, F. R. (1970). Mycoplasma hominis and abortion. *J. Infect. Dis.* 121: 260-268.
 45. De-Louvois, J., Hurley, R. and Stanley, V. C. (1975). Microbial flora of the lower genital tract during pregnancy: relationship to morbidity. *J. Clin. Pathol.* 28: 731-735.
 46. Iwasaka, T., Wada, T., Kidera, Y. and Sugimori, H. (1986a). Hormonal status and mycoplasma colonization in the female genital tract. *Obstet. Gynecol.* 86: 263-267.
 47. McCormack, W. M., Rosner, B., Lee, Y. H. (1973b). Colonization with genital Mycoplasma in women. *Am. J. Epidemiol.* 97: 240-245.
 48. Upadhyaya, M., Hibbard, B. M., Walker, S. M. (1983). The role of Mycoplasmas in reproduction. *Fertil. Steril.* 39: 814-817.
 49. Gnarpe, H. and Friberg, J. (1973a). Mycoplasma and human reproductive failure: The occurrence of different mycoplasma in couples with reproductive failure. *Am. J. Obstet. Gynecol.* 114: 727-31.
 50. Fowlkers, D. M., Macleod, J. and O'Leary, W. M. (1975). T-mycoplasma and human infertility: Correlation of infection with alteration in seminal parameters. *Fertil. Steril.* 26: 1212-8.
 51. Taylor-Robinson, D., (1986). Evaluation of the role of *Ureaplasma urealyticum* in infertility. *Pediatric infect. Dis.* 5: S262-S265.
 52. Bercovici, B., Persky, S., Rozansky, R., Razin, S. (1962). Mycoplasma (pleuro pneumonia-like organisms) invaginitis. *Am. J. Obstet. Gynecol.* 84: 687-691.
 53. Mendel, E. B., Rowan, D. F., Graham, J. H. M., Dellinger, D. (1970). Mycoplasma species in the vagina and their relation to vaginitis. *Obstet. Gynecol.* 35: 104-108.