

# Parathyroid Preservation during Total Thyroidectomy and its Outcome

Sharfuddin Mahmud<sup>1</sup>

<sup>1</sup> Sir Salimullah Medical College Mitford Hospital

*Received: 12 April 2015 Accepted: 30 April 2015 Published: 15 May 2015*

---

## Abstract

**Objectives:** To know the incidence of hypocalcaemic tetany at post surgical state of total thyroidectomy. **Methods:** It was a cross sectional observational study among 50 patients who underwent total thyroidectomy in the Dept. of Otolaryngology and Head Neck Surgery, Sir Salimullah Medical College and Mitford Hospital Dhaka Medical College Hospital during the period from January 2014 to December 2014. **Results:** In this study out of 50 patients 14 (28

---

*Index terms*— arathyroid glands, total thyroidectomy, hypocalcaemic tetany.

## 1 I. Introduction

In the past century, a lethal outcome occurred in more than 40% of patients undergoing thyroidectomy and morbidity levels were also consequently very high. Through the contribution of several pioneers in thyroid surgery such as Billroth, Halsted and Kocher, thyroid surgery has become a relatively safe operative procedure. Nonetheless, there are still many complications related to thyroid surgery. Among them hypoparathyroidism is perhaps the most distressing. Parathyroid preservation during thyroidectomy is not only desirable, but essential for the effective management of surgical diseases of the thyroid gland. Most individuals possess at least two pairs of parathyroid glands. To the unaided eye, the glands are a yellow brown orange colour. At operation brown fat, yellow fat, sequestered thyroid tissue, thymus, lymph node and autonomic ganglia may all mimic these appearances. Each weighs about 50 mg. The lowest post-operative serum-calcium level that can be regarded as normal has not been agreed. Wade (1960) thought that any patient with a serum-calcium level between 8 and 9 mg/100 ml was in need of treatment, and those with a level below 7.5 mg/100 ml were in need of urgent treatment. Biochemical monitoring of serum calcium and PTH levels can assist in initiating early management for postoperative hypocalcemia and reducing hypocalcemic symptoms during the early weeks after surgery. The reasons for postoperative hypoparathyroidism are devascularizations of parathyroid glands during surgery owing to the close proximity to the thyroid capsule, the accidental removal of one or more parathyroid gland (s), destruction of the parathyroid glands as a result of lymphadenectomy or hypoparathyroidism due to hematoma formation. Permanent and transient hypoparathyroidism can be reduced by identification of the parathyroid glands, dissection close to the thyroid gland, preservation of the blood supply to the parathyroid glands and avoiding manipulation of parathyroid glands. Ligation of the inferior thyroid artery preferably should be performed close to the thyroid capsule to minimize the risk of postoperative hypoparathyroidism. Vascular injury probably far more important than inadvertent removal. The incidence of permanent hypoparathyroidism should be less than 1% and most cases present dramatically 1 -5 days after operation; however very rarely onset is delayed 2 -3 weeks or a patient with marked hypocalcaemia remain asymptomatic. The incidence of temporary hypoparathyroidism is approximately 10 -20 percent and is often an inevitable consequence of total thyroidectomy and this may be reduced by sound anatomical knowledge, surgical technique and experience. Removal of a single parathyroid gland is not associated with postoperative hypocalcemia, resection of at least 2 parathyroid glands increases the risk of transient and permanent hypoparathyroidism. Careful examinations of the surgical specimen intraoperatively decreases the incidence of inadvertent parathyroidectomy during thyroidectomy, ligation of the superior thyroid artery after identifying and saving its anastomotic vessel to inferior thyroid artery and ligation of the inferior thyroid artery close to the gland if at all possible.

## 2 Results

### 3 Table -I :

Age and sex distribution of the sample (n=50)

## 4 III. Discussion

In this series 50 patients were included in the study those who had undergone total or completion thyroidectomy with or without neck dissection. Among them 14 (28%) were male and 36 (72%) were female. Male to female ratio is 1:2.57 (Table-I). Similar sex distribution has been found in another study where male 22% and female 78% and male to female ratio was 1:3.57. 13 In our study out of 50 patients 29 (58%) were multinodular goiter, 19 (38%) patients were papillary thyroid carcinoma, 1 (2%) patients were medullary thyroid carcinoma and 1 (2%) and were Graves disease (Table-II). In a study shows that indication of thyroidectomy in thyroid cancer 52% case and Benign multinodular goiter 48%. 7 Regarding clinical presentation of the study population thyroid swelling was present in all cases (100%). Out of them 12 (24%) patients were presented along with cervical lymphadenopathy (Table-III). Lymph nodes involvement was most common in level III (90.9%) and level IV (72.72%). Clinically no palpable lymph nodes were found in level I. Unilateral lymphadenopathy was 9(75%), whereas bilateral involvement was 3 (25%) (Table-IV). Result correspond with the other study which showed in level III 82% and in level IV 75%. 15 In our study most hypocalcaemia developed in patients underwent thyroidectomy with neck dissection (58.33%); it was followed by completion thyroidectomy 40% and total thyroidectomy 18.18% (Table-V).

The difference between thyroidectomy with neck dissection and thyroidectomy alone in the development of hypocalcaemia was statistically significant.  $\chi^2 = 6.03, P < 0.05$  i.e. in case of more extensive disease, the chance of developing hypocalcaemia is significantly more. In one study shows that hypoparathyroidism occurs more frequently in patients with large extension and invasive thyroid cancer and when total thyroidectomy is associated with central and lateral neck dissection. 14 In this study (Table-VI) it is found that occurrence of hypocalcemic tetany is more in the group where parathyroid was not identified preoperatively.

The difference in the development of hypocalcemic tetany between identification and not identification of parathyroid gland was statistically significant ( $\chi^2 = 4.79, p < 0.05$ ).

So it can be concluded that there is a significant association between development of hypocalcaemia and not identification of parathyroid preoperatively which coincides with other study. 16 In our study it is found that maximum patients 10 (66.73%) were developed hypocalcemic tetany in second postoperative day (n= 15), 3(20%) cases developed hypocalcaemia in 3 rd post operative day (Table-VII). One study shows that symptoms of hypocalcaemia usually appear 24-48 hours after surgery. 13 This finding is similar to that of the current study.

A study conducted at King Abdul Aziz University. Jeddah, Saudi Arabia in 2005 revealed transient hypocalcaemia occurred 6.9% to 42% and permanent hypocalcaemia 4% to 29% . 18 One study shows that post operative hypocalcaemia is the most common complication with incidence ranging from 1.6 to 50% . 13 The result in this series showed that temporary hypocalcaemia occurred in 24% cases and permanent hypocalcaemia occurred in 6% cases (Table-VIII), which is within the range of other study.

## 5 IV. Conclusion

In this study we suggest that inferior thyroid artery should be ligated close to the thyroid capsule and at least two parathyroid gland should be identified and preserved .In the context of our country, now a days most of our thyroid surgeons are expert enough to identify and preserve recurrent laryngeal nerve and external branch of superior laryngeal nerve but identification and preservation of parathyroid gland during thyroid surgery is not well practiced. Even if all the parathyroid glands cannot be completely accounted for, the surgeon should make an attempt at identifying and preserving parathyroid glands without damaging their blood supply. Proper knowledge and expertness regarding parathyroid preservation can alleviate sufferings of the patient due to iatrogenic morbidity.. Now a days it is more concern all over the world.

Table-VII : Time interval		No. of patients		Percentage (%)	
1 st POD		1		6.66%	
2 nd POD		10		66.73%	
3 rd POD		3		20%	
4 th POD		1		6.66%	
Age		Sex			
		male	female	Total	
10-20yrs		0	3	3	
21-30yrs		4	9	13	
31-40yrs		5	12	17	
41-50 yrss		2	5	7	
>50yrs		3	7	10	
Total		14	36	50	
Surgery		No. of patients		No. of Hypocalcemia	Percentage (%)
Total thyroidectomy		33	6	18.18	
Completion thyroidectomy		5	2	40	
Thyroidectomy and neck dissection		12	7	58.33	
Group		Hypocacaemia	No Hypocacaemia	Total	Percentage of Hypocalcaemia
Group A (Parathyroid Identified)	10	32		42	23.8%
Group B (Parathyroid not Identified)	5	3		8	62.5%
Total	15	35		50	

Figure 1: Table - V

Type	No. of patients	Percentage (%)
Temporary hypocalcaemia	12	24
Permanent hypocalcaemia	3	6
Total	15	30

Figure 2: Table -



---

91 Magnification during surgery is helpful in preservation of parathyroid gland.

92 [India] , India .

93 [Kuwait Medical Journal] , *Kuwait Medical Journal* 37 (3) p. . Aziz University Hospital (Jeddah

94 [Cjk and Connell ()] *Bailey and Love's short practice of surgery*, Williams Cjk , O Connell , PR . 2008. London:

95 Chapman and Hall. 2 p. 791. (5th ed)

96 [Glinoyer et al. ()] 'Clinical aspects of early and late hypocalcaemia after thyroid surgery'. D Glinoyer , G Andry

97 , G Chanrain . *European Journal of surgical oncology* 2000. 26 p. .

98 [Goepfert and Callender ()] 'Differentiated thyroid cancer-papillary and follicular carcinoma'. H Goepfert , D L

99 Callender . *American Journal of Otolaryngology* 1994. 15 p. .

100 [Qari ()] *Estimation of ionized calcium levels after thyroidectomy at King Abdul*, F A Qari . 2005.

101 [Asari et al. ()] 'Hypoparathyroidism after total thyroidectomy'. R Asari , C Passler , M K Kaazireh . *Arch Surg*

102 2008. 143 (2) p. .

103 [Kim ()] 'Impact of preserving the parathyroid gland on hypocalcaemia after total thyroidectomy with neck

104 dissection'. Y S Kim . *J Korean Surg Soc* 2012 august. 83 (2) p. .

105 [Aaron et al. ()] 'Incidental Parathyroidectomy During Thyroid Surgery Does Not Cause Transient Symptomatic

106 Hypocalcaemia'. R Aaron , M D Sasson , F James . *Arch Otolaryngol Head & Neck Surg* 2001. 127 p. .

107 [Sinnatamby ()] *Last's anatomy*, C S Sinnatamby . 2011. 12. (th ed)

108 [Lee et al. ()] B J Lee , S G Wang , J C Lee . *Archives of otolaryngology-Head and Neck Surgery*, 2007. 133 p. .

109 [Ashok et al. ()] 'Parathyroid autotransplantation during thyroid surgery'. R Ashok , C Vurnett , M Bernard .

110 *Journal of surgical oncology* 2006. 46 p. .

111 [Girling and Murley (1963)] 'Parathyroid insufficiency after thyroid surgery'. J A Girling , R S Murley . *Br Med*

112 *J* 1963 May. 1 (5341) p. .

113 [Shaha and Jaffe ()] 'Parathyroid preservation during thyroid surgery'. A R Shaha , B M Jaffe . *Am J Otolaryngol*

114 1998. 19 p. .

115 [Kapre ()] *Preservation of the Parathyroids in the Thyroid Surgery. Case study from*, Madan Kapre . 1994. 2002.

116 [Zarnegar et al. ()] 'Prevention, evaluation and management of complications of following thyroidectomy for

117 thyroid carcinoma'. R Zarnegar , L Brunaud , O H Clark . *Endocrinol metab Clin N Am* 2003. 32 p. .

118 [Ramsden and Watkinson ()] J Ramsden , J C Watkinson . *Scott-Browns Otolaryngology-Head and Neck Surgery*,

119 2008. 7 p. . (th edition)

120 [Tartaglia et al. ()] 'Randomized study on oral administration of calcitriol to prevent symptomatic hypocal-

121 caemia after total thyroidectomy'. F Tartaglia , A Giulion , M Sgueglia . *The American Journal of Surg* 2005.

122 190 p. .

123 [Marwah et al. ()] 'Routine parathyroid autotransplantation during subtotal thyroidectomy for benign thyroid

124 disease'. S Marwah , R Godara , A Kapoor . *The Internet Journal of Surgery* 2009. 7.

125 [Thomusch et al. ()] 'The impact of surgical technique on postoperative hypoparathyroidism in bilateral thyroid

126 surgery: a multivariate analysis of 5846 consecutive patients'. O Thomusch , A Machens , C Sekulla , J Ukkat

127 , M Brauckhoff , H Dralle . *Surgery* 2003. 133 p. .

128 [Toniato et al. ()] 'Thyroidectomy and parathyroid hormone; tracing hypocalcaemia prone patients. The Am'. A

129 Toniato , I M Boschini , A Pitto , M Pelizzo . *J. Surg. Pp* 2008. p. .

130 [Roh et al. (2007)] 'Total thyroidectomy plus neck dissection in differentiated papillary thyroid carcinoma

131 patients'. J L Roh , J Y Park , C Park . *Ann Surg* 2007 April. 245 (4) p. .

132 [Watkinson and Gilbert ()] *Tumours of the thyroid and parathyroid gland. Stell and Maran's Head and Neck*

133 *Surgery and oncology*, Jc & Watkinson , R W Gilbert . 2012. 5 p. . (th ed.)