

Bilateral Knee Dislocation with Tibial Shaft Fracture

Rafik Elafram

Received: 13 December 2015 Accepted: 4 January 2016 Published: 15 January 2016

Abstract

Acute dislocation of the knee is a limb-threatening injury that often results in extensive soft-tissue damage and disruption of the popliteal blood vessels. We report a case of traumatic bilateral open knee dislocation with a type 42 A2 closed right tibial shaft fracture and right common peroneal nerve palsy. We are not aware of any other reports of such a combination of injuries.

Index terms— bilateral, knee, dislocation, tibial shaft fracture.

1 I. Introduction

Acute dislocation of the knee is a limb-threatening injury that often results in extensive soft-tissue damage and disruption of the popliteal blood vessels [1,2].

We reported a case of traumatic bilateral open knee dislocation with a type 42 A2 closed right tibial shaft fracture and right common peroneal nerve palsy.

We are not aware of any other reports of such a combination of injuries.

2 II. Case Report

In April 2013, 25-year-old male was admitted to emergency medical service by an ambulance due to a traffic accident. He was hit by a car.

After initial examination, he had severe deformity of both knees with posterior skin injury in both popliteal fossa. Also, the patient had right limb deformity.

The pulses were palpable and symmetrical in the lower limb. The patient had right common peroneal nerve palsy.

Plain radiographies of the lower extremities were performed: they showed medial right knee dislocation with ipsilateral type 42 A2 tibial shaft fracture and fracture dislocation of the left knee.

A prompt reduction was performed. The arteriogram was made and had not shown any sign of occlusion.

In the operation room, exploration of the knees' injury revealed: in the right knee a section of the biceps femoris, bruises in the common peroneal nerve and the ACL and the PCL were intact, in the left knee, no noble element was affected. The fracture was fixed with screws.

Internal fixation of the tibial and fibular shaft were performed. The patient had bilateral knee immobilization for 45 days. Then, he begun rehabilitation. The consolidation fracture was obtained after 04 months.

In the final follow up, the range motion of the knee was 0 degree to 95 degree. He had a grade-1 anterior and posterior laxity and varus instability in the right knee. The lateral peroneal nerve palsy was recovered.

3 III. Discussion

Knee dislocations are uncommon, constituting less than 0.5% of joint dislocations [3]. The documented incidence of observed knee dislocations on admission per institution per year is even less and varies from 1/10,000 to 1/100,000 [4][5][6].

The exact mechanism responsible for the tibial shaft fracture and knee dislocation with disruption of all knee ligaments, popliteal vessels, and the common peroneal nerve was not clear [7].

In a large review by Green and Allen, 40% of 245 knee dislocations were anterior, while posterior dislocations are the second most common at 33% and are caused by direct application of a posterior force to the anterior

tibia. Lateral and medial dislocations are relatively uncommon, comprising 18% (lateral) and 4% (medial) of knee dislocations [8].

A thorough neurological examination is also essential, as peroneal nerve palsies have been noted in 14-35% of knee dislocations, most commonly in posterolateral dislocations [1,6,7].

Controversies over operative versus closed immobilization of traumatic complex, multiple ligamentous knee injury are still debated. In 2004, Chin-Ho Wong et al. investigated the results of surgical and conservative treatment of knee dislocation retrospectively. The international knee documentation committee (IKDC) scores of operatively treated patients and patient satisfactions were significantly better than conservatively treated group [10]. The most important fear in this kind of associations is to miss diagnose one dislocated joint because the other is more spectacular or threatening the limb's vitality, especially in polytrauma patients. Moreover, these associations present an evident problem of interference in management.

Two cases of combination of ipsilateral knee dislocation and tibial shaft fracture were reported in the litterature. But no bilateral open knee dislocation was noted. Therefore, we found that it would be interesting to report such an association [7,11].

4 IV. Conclusion

These injuries are orthopedic emergencies that have high complication rates. An awareness of the possibility of such association should lead to an appropriate treatment.



Figure 1: Figure 1 :



2

Figure 2: Figure 2 :



3

Figure 3: Figure 3 :

45



Figure 4: Figure 4 :Figure 5 :

-
- 61 [Ayd?n A 1 , Atmaca and Müezzino?lu (2013)] ‘Anterior knee dislocation with ipsilateral open tibial shaft
62 fracture: a 5-year clinical follow-up of a professional athlete’. H Ayd?n A 1 , Atmaca , Ü S Müezzino?lu
63 . *Musculoskelet Surg* 2013 Aug. 97 (2) p. .
- 64 [Kennedy ()] ‘Complete dislocation of the knee joint’. J C Kennedy . *J Bone Joint Surg Am* 1963. 45 p. .
- 65 [Shields et al. ()] ‘Complete dislocation of the knee: experience at the Massachusetts General Hospital’. L Shields
66 , M Mital , E F Cave . *J Trauma* 1969. 9 p. .
- 67 [Sisto and Warren ()] ‘Complete knee dislocation. A follow-up study of operative treatment’. D J Sisto , R F
68 Warren . *Clin Orthop Relat Res* 1985. 198 p. .
- 69 [Hegyes et al. ()] ‘Knee dislocation. Complications of nonoperative and operative management’. M S Hegyes , M
70 W Richardson , M D Miller . *Clin Sports Med* 2000. 19 p. .
- 71 [Richter et al. ()] ‘Knee dislocation. Long term results after operative treatment’. M Richter , P Lobenhoffer , H
72 Tscherne . *Chirurgia* 1999. 70 p. .
- 73 [Schenck et al. ()] ‘Knee dislocations’. R C SchenckJr , R E Hunter , R F Ostrum , C R Perry . *Instr Course*
74 *Lect* 1999. 48 p. .
- 75 [Wong et al. ()] ‘Knee dislocations-a retrospective study comparing operative versus closed immobilization
76 treatment outcomes’. C H Wong , J L Tan , H C Chang , L W Khin , C O Low . *Knee Surg Sports Traumatol*
77 *Arthros* 2004. 12 p. .
- 78 [Gupta et al. ()] ‘Traumatic anterior knee dislocation and tibial shaft fracture: a case report’. S Gupta , Ma
79 Fazal , Haddad . *Journal of Orthopaedic Surgery* 2007. 15 (1) p. .
- 80 [Meyers et al. ()] ‘Traumatic dislocation of the knee joint’. M H Meyers , T M Moore , J P HarveyJr . *J Bone*
81 *Joint Surg Am* 1975. 57 p. .
- 82 [Green and Allen ()] ‘Vascular injuries associated with dislocation of the knee’. N E Green , B L Allen . *J Bone*
83 *Joint Surg Am* 1977. 59 p. .