Artificial Intelligence formulated this projection for compatibility purposes from the original article published at Global Journals. However, this technology is currently in beta. *Therefore, kindly ignore odd layouts, missed formulae, text, tables, or figures.*

Assessment of Parent-Adolescent Communication about Sexual and Reproductive Health among High School Students in Mekelle Town, Northern Ethiopia

Desta Hailu¹, Zemenu Yowhanes² and Hailemariam Berhe³

¹ Arba Minch University

Received: 16 December 2015 Accepted: 1 January 2016 Published: 15 January 2016

7

Δ

5

6

8 Abstract

⁹ Parents have the potential to protect against adolescent sexual risk, including early sexual

¹⁰ behavior and its outcomes such as unwanted pregnancy and sexually transmitted infections.

¹¹ The objective of this study therefore was to assess the status of adolescent-parent

¹² communication about sexual and reproductive health and associated factors among school

¹³ students. Institution based cross sectional study was conducted. Data were entered, cleaned

¹⁴ and analyzed using SPSS.A total of 521 students were included in the study, making a

¹⁵ response rate of 97

16

17 Index terms—adolescent, parent, communication, sexual, reproductive, health, northern ethiopia.

¹⁸ 1 I. Introduction

19 ccording to world health organization (WHO), the term "adolescence is defined as period of life between 10-19 years [WHO, 2013; Society for public health education; ??HO, 2012]. It represents a unique period of life 11 characterized by significant physical, cognitive, emotional and social changes ??CDC, 2004]. It is a transitional 12 period from child hood to adult hood ??WHO, 2014] marked by increasing levels of individual autonomy, growing 13 sense of identity and selfesteem and progressive independence from adults [Save the ??hildren, 2009].

Traditionally, these group of people have commonly been regarded as healthy segment of the community and little attention have been given since the health care system was almost exclusively adult centered ??Georg, 2012]. However, it is a period when health problems that have serious immediate consequences and behaviors that could have long lasting adverse effects on their future are initiated ??WHO, 2010; ??DC, 2007]. Because of the rapid physical, cognitive, and emotional developments that takes place before adequate information, skills and experience of life is achieved, adolescence is a time when many health problems first emerge unless managed properly ??CDC, 2007; ??HO, 2006].

The health status of adolescents is strongly connected to a number of risk behaviors, which are often established during the adolescent years that end in chronic and non-chronic diseases. These behaviors include: substance abuse, unhealthy dietary behaviors, inadequate physical activity, violence, risky sexual behaviors that contribute to unintended pregnancy and sexually transmitted infections including HIV infection [Society for Public Health Education,2012; ??HO,2006].

One third of women worldwide give birth before the age of 20 each year and are at increased risk of morbidity and mortality due to obstetric complications. Annually, 5,000,000 and 70,000 adolescents between the ages of 15 and18 have unsafe abortions and abortion related deaths respectively ??EFDR, 2004]. Moreover, adolescents are more likely to engage in a wide range of high risk sexual behaviors that can result in sexually transmitted diseases, including HIV ??WHO, 2010].

Globally, One-third of all currently infected individuals are young people acquiring HIV infection every minute and more than 2.6 million each year [National center for Health Statistics, 2012].

In Ethiopia, like in other sub Saharan African countries, sexual and reproductive health problems of adolescents 43 are high. Twelve percent of adolescent women are already mothers or pregnant with their first child [Central 44 statistical agency, 2011] and 60% of adolescent pregnancies are unwanted or unintended ??WHO, 2010].According 45 to EDHS report, 1.9% and 3.8% of female and male adolescents had premarital sex respectively. Among those 46 who have premarital sex only 45.5% women and 60% men used condom. This unsafe sexual intercourse leads to 47 HIV prevalence of 0.2% among adolescent women [Central statistical agency, 2011]. 48

The health, safety and well-being of adolescents are affected by a complex interplay of factors between the individual and their surroundings, such as parents, peers, neighborhood, schools, community, health care systems, 50 media, social norms, policies and laws. These factors collectively impact young people's ability to appropriately make healthy decisions ?? Despite its importance, communication about sexual and reproductive health is greatly 52 influenced by the culture and social environment. Parent-adolescent communication remains challenging in many sub-Saharan African countries including Ethiopia as the social milieu in many traditional communities still limit. 54 Initiating conversations about the sexual issues may be difficult for parents as they may be unsure as to how to approach such issues, doubt their competence in handling sexuality topics and the questions that may be raised 56

by their adolescence or feel confused about the proper amount of information to offer ?? EFDR, 2004]. 57

58 While parent-adolescent communication has critical role in preventing adolescents from engaging in risky 59 sexual behaviors and promoting access to contraceptives and maternal health services, earlier studies in Ethiopia 60 have paid limited attention to adolescent health and the level of communication is not well documented. The 61 objective of this study therefore is to determine the level of parent-adolescent communication about reproductive

and sexual health and associated factors among secondary and preparatory school students in Mekelle University. 62

2 II. Methods and Materials a) Study setting and period 63

This study was conducted from Jan 2013-August 2013 in Mekelle Town. Mekelle is located in the northern part 64 of Tigray regional state, 783 Kilometers away from Addis Ababa, capital city of Ethiopia. It covers 28 square 65 kilometers and has an estimated population of 289,756 ??UNFPA, 2003]. The town has a total of 24 secondary 66

and preparatory schools (eight governmental and sixteen private schools). The total number of students for the 67 academic year 2012/2013 was 21,173. Of those, 45.1% and 54.89 were males and females respectively [Tigray 68

region education bureau, 2013]. 69

49

51

53

55

3 b) Study design and Population 70

A school based cross-sectional study involving qualitative method was undertaken among randomly selected 71 unmarried adolescent students aged 10-19 who was attending secondary and preparatory school during the time 72

of data collection. Mentally and/or physically incapable adolescents' students were excluded from the study. 73

c) Sample size determination and sampling procedure 4 74

Single population proportion formula was used to calculate the required sample size. Proportion of parent-75 adolescent communication about sexual and reproductive health [Tesso et al, 2012], margin of error, confidence 76 interval, design effect and non-response rate were assumed to be 69.5%, 5%, 95%, 1.5 and 10%, respectively.n i 77 = Z (?/2) 2 P (1-P)- d**2** 78

Where: n i = Sample size; Z (?/2) 2 = confidence interval; P = proportion of parent adolescent communication 79 (0.695.); D = marginal of errorThus, Ni = (1.96) 2 (0.695) (1-0.695) (0.05) 2 = 325 80

Finally, taking 10 % non-response rate and 1.5 design effect: $10 \% \ge 325 = 358 \times 1.5 = 537$ Multistage sampling 81 technique was used to select the study subjects. First, all the schools in the town were stratified in to private and 82 governmental. Then 3 out of 16 private and 3 out of 8 governmental schools were randomly selected. Then schools 83 were further stratified by grade and section. Calculated sample size was proportionally allocated to private and 84 governmental schools according to their number of students. Then, frames of students were developed from 85 student roster of each grade in collaboration with instructors of respective classes. Students whose age range 86 10-19 years were selected using simple random sampling from the existing sampling frame (students' roster). In 87 every step of selection simple random sampling technique was used. participants were selected. 88

5 d) Data collection tool and process 89

Structured self-administered questionnaire was prepared and utilized after reviewing relevant literatures. The 90 main contents of the questionnaire were sociodemographic characteristics, knowledge of major reproductive and 91 92 sexual health, sexual attitude and behavior, parental monitoring and communication with adolescents. The 93 questionnaire was prepared in English and then translated to Tigrigna (local language). To check its consistency, 94 it was back translated to English by an expert of both languages. After extensive Volume XVI Issue II Version

95 IYear 2016 (DDDD)

evaluation, final version of the questionnaire was developed. 96

Six 12th grade completed female interviewers who were fluent in the local language (Tigrigna) and are familiar 97 with the local customs collected the data. Two diploma holder health care workers with similar work experience 98 were assigned to supervise the data collection process. 99

Training was given for both data collectors and supervisors by the principal investigator for two days. 100

The training session includes the general objective of the study; content of the questionnaire, ways how to keep confidentiality and privacy and ways how to resolve when a problem arises.

¹⁰³ 6 e) Data Quality Control

An individual who have the ability to speak and write both English and Tigrigna languages translated the questionnaire. In order to identify the clarity and consistency of the questionnaire it was pretested on 5% of the sample in a similar population in Wukro high school and preparatory school other than the schools in the study area and necessary modifications such clarity and consistency of questions and evidence based time allocation for each respondent were made accordingly. All the data collected from each respondent were checked for completeness, clarity and consistency by the principal investigator and the supervisors immediately at the end of each data collection days.

¹¹¹ 7 f) Data processing and analysis

Data were entered, coded and cleaned using SPSS version 16 software. Univariate analysis was computed for each independent variable to assess their individual proportion. Then, bivariate analysis was done to examine crude association of predictors on parent adolescent communication. Finally, by selecting eligible variables using forward logistic regression, the independent effect of predictors on parent-adolescent communication about sexual and reproductive health were examined. Odds ratio and 95% CI were used to measure the statistical association. P value 0.05 was used to determine the statistical significance of the tests. Finally, the results were presented in

118 texts, tables and graphs.

¹¹⁹ 8 g) Ethical Consideration

After it was thoroughly reviewed, ethical approval was secured from Mekelle University college of Health Science

(MUCHS) and permission to conduct the study was obtained from Tigray regional education bureau and management body of the respective school facilities. Informed written consent was obtained from each respondent.

management body of the respective school facilities. Informed written consent was obtained from each respondent. Involuntary participants were free to withdraw from the study. Questionnaires were coded instead of using names

as identification and hence, confidentiality was assured throughout the study.

125 9 III. Results

¹²⁶ 10 a) Socio-demographic characteristics of adolescents

A total of 521 respondents included in the study making a response rate of 97%. Sixteen questionnaires which 127 were initially administered to respondents were excluded as they were incompletely &/or inconsistently filled. 128 Thus, analysis was made based on 521 respondents. Among the study subjects, 231 (44.3%) were males and 129 290(55.7%) were females making the male to female ratio of 1 to 1.3. Mean age of respondents' was $16.59\pm$ 130 1.33SD years ranging from 13-19 years. Tigray by ethnicity and Orthodox by religion were found to be 502 131 (96.4%) and 464 (89.1%) respectively. Majority 388 (74.5%) of the respondents were living with both of their 132 fathers and mothers. Regarding the educational status of their parents, Only 108(20.8%) and 37(7.1%) of their 133 mothers and fathers were illiterate respectively (table1). 134

¹³⁵ 11 c) Sexual Behavior of adolescent students

One hundred nineteen (22.8%) of the students believed that it is normal and acceptable to have sexual feeling during adolescent period. Majority of the respondents 341(65.5%) believed that sexual intercourse should not be made before marriage. Regarding their practice, Eighty three (15.9%) of the students had made sexual intercourse. The mean age when sexual practice started was 15.9+1.6. Among those who practiced sexual intercourse, 22(26.5%) reported, they made sexual intercourse with unknown person. Most 60 (72.28%) of the students made sex using condom.

¹⁴² 12 d) Attitude and practice of adolescent students towards ¹⁴³ parental monitoring

Most 344(66%) of the respondents agreed that parental monitoring to adolescents activities should be in place. Ninety (38.1%) of females reported that they were not allowed to have relationship with opposite sex, while the rest 146(61.86%) were allowed. Three hundred forty eight (66.8%) of the students reported that their parents knew with whom their son or daughter are when out of home, while 173(33.2%) of the respondents reported that parents did not know.

13 e) Parent-adolescent Communication & source of informa tion about sexual & reproductive health issues

Two hundred ninety six (56.8%) of respondents got information on sexual &RH issues from school followed by media 239 (45.9%) (fig2). Three hundred (57.6%) participants reported that they had discussed about at least one topic of sexual and /or reproductive health with either of their parents. Cultural unacceptability, being ashamed and lack of knowledge were major reasons cited by participants for not discussing about family planning (Table3) In multivariate logistic regression analysis, parents' educational status, living arrangement and level of education of respondents were found to be significantly associated with communication of adolescents with their parents about sexual and reproductive health matters (table 4). (AOR=1.80, 95% CL: [1.03-3.13]) more likely

to communicate about sexual and reproductive health matters with their parents as compared to those who had no formal education.

¹⁶⁰ 14 Volume XVI Issue II Version I

161 On the contrary, adolescents' level of education was found to be negatively related with the outcome variable.

Adolescents whose level of education was 12 were less likely to communicate about sexual and reproductive health with their parents as compared to those whose educational level was grade nine (AOR: 0.55, 95%CI: [0.31-0.97]).

¹⁶⁴ 15 IV. Discussion

204

Parents are important role models in adolescents' lives; they can directly or indirectly transmit values, traditions and life styles to their children. Positive family communication helps teens develop the values, security, and sense of worth that can lead to healthy decision making. This school based cross-sectional study has therefore examined adolescent communication with their parents concerning sexual and reproductive health matters among adolescent students in Mekelle secondary and preparatory school students, Tigray region, Ethiopia.

In line with the evidence from a study done in Nekmete town, and Myanmar [Seme & Wirtu, 2008; Nuoo et al, 170 171 2011], the finding of the present study indicated that the prevalence of adolescents' communication about at least 172 one sexual and reproductive health matters with their respective parents was 57.6%. This finding was higher when compared to the result of a cross sectional study done in Benishangul gumuz regional state 28.9% [Yesus & 173 Fantahun, 2010] and Debremarkos town 36.9% ??Shiferaw et al, 2014]. This might be due to the time gap that 174 there could be improvement in accessing and utilizing sexuality and reproductive health service. However, it is by 175 far lower than study findings from United States and Mexico [Jerman & Constantine, 2010; Erika, 2009] where 176 85% and 81.5% adolescents communicated with their parents about SRH matters respectively. This difference 177 could be attributed to the fact that proportion of parents in the United States was more educated, reproductive 178 health services were accessible in-terms of minimal distance and transport availability, and adolescents could have 179 better decision making autonomy. 180

In multivariate analysis, level of education showed strong statistical association with communication about 181 sexual and reproductive health. Adolescents whose mothers' educational level 12 and above were about 3 times 182 more likely to communicate about sexual and reproductive health matters with their parents as compared to 183 those whose parents had not attended formal education. Similarly, adolescents whose father's educational level 184 12 and above were about 2 times more likely to communicate about sexual and reproductive health matters with 185 their parents when compared to those whose parents had no formal education. This is congruent with previous 186 finding from study done in Rwanda ??Bushaija et al,2013].The possible explanations might be educated parents 187 have better access to health service information, improved perceptions of the causes and treatment of sexual 188 and reproductive system related disease and play critical role in helping their children use quality reproductive 189 healthcare services. 190

In contrary to evidence from a study done in Nekemte, Debremarkos and Lincoln [Seme & Wirtu, 2008; Shiferaw et al, 2014; Raffaelli & Green, 2003], this finding revealed that adolescents whose level of education was 12 were less likely to communicate about sexual and reproductive health with their parents as compared to those whose educational level was grade nine. This might be due to the reason that as educational level of adolescents increase they start underestimating the idea of their parents as they believe that they are more knowledgeable than their mothers or fathers and may not get inspired to communicate about sexual and reproductive health.

Among respondents who made communication regarding SRH matters, 12.9% and 31.9% adolescents discussed about premarital sex with their father and mother respectively. This is higher when compared with the findings from study in Deberemarkos ??Shiferaw et al, 2014], where 11.3and 15.3 of adolescents' father and mother discussed about premarital sex respectively. In line with a finding from study done in Deberemarkos ??Shiferaw et al, 2014], this study showed that the major reason for not discussing premarital sex was being ashamed. This can be attributed to the taboo nature of sexual and reproductive communication in traditional African settings including Ethiopia.

When interpreting the finding of this study, some limitations should be considered.

Communication on sexual and reproductive health as well as sexual behaviors was based on the self-report of respondents, and provided no validation of obtained information with any objective source, which was likely to be subjected to reporting biases. Moreover, this study was based on cross-sectional data, which implies that the direction of causal relationships cannot be determined.

²⁰⁹ 16 V. Conclusion and Recommendations

In this study almost half of school adolescent has never had communication with their parents regarding sexual and reproductive health matters. Educational status of their mother and father, living arrangement and level of education of respondents were found to be significantly associated with communication of adolescents with their parents about sexual and reproductive health matters. Among the respondents 12



Figure 1:

213

 $^{^{1}}$ © 2016 Global Journals Inc. (US)

 $^{^2 \}rm Assessment$ of Parent-Adolescent Communication about Sexual and Reproductive Health among High School Students in Mekelle Town, Northern Ethiopia



Figure 2: Figure 1 :

1

31 Volume XVI Issue II Version I Medical Research Global Journal of

Figure 3: Table 1 :

Year 2016		
Volume XVI Issue II Version I		
D D D D)		
Variable	Numb	
Sex b) Knowledge of adolescents about contraceptive Male Female meth-	231	44.3
ods & sexually transmitted infections Among the adolescent students,	290	55.7
409 (78.5%) Age were aware of at least one contraceptives methods.		
10-14 Three hundred four (64.1%) & 301 (57.8%) participants	22	4.2
15-19 reported that they had heard about condom and pill	499	95.8
Grade respectively (fig1).		
Grade 9 Grade 10 Grade 11 Grade 12 Adolescent spontaneously mention	144	27.6
STI. Accordingly, 415(79.7%) students were asked to respondents men-	157	30.1
tioned at least one type of STI. Among the respondents, 399(76.6%), 302	117	22.5
(58%), 293(56.2%),	103	19.8
Type of school $270(51.8\%)$ and $114(21.9\%)$ knew about HIV/AIDS,		
Private Public Gonorrhea, syphilis, Chancroid and lymphgranulo-	66	12.67
mavenerum respectively.	455	87.33

[Note: ? Aunt, grandparents, uncle, sister, brother; ? Protestant, catholic; ? Oromo, Guragie;?carpenter, tela (local bear) seller]

Figure 4: E

Religion		
Orthodox Muslim Others ?	$464 \ 30 \ 27$	$89.1 \ 5.8 \ 5.1$
Ethnicity		
Tigray	502	96.4
Amhara	16	3.1
Others ?	3	0.5
Living arrangement		
With both parents With mother only With father only	$388\ 88\ 13$	$74.5 \ 16.9 \ 2.5$
With other relatives With others?	$10 \ 21$	$1.9 \ 4$
Mother's educational status		
Illiterate	108	20.8
Read and write	122	23.4
Primary (1-8)	46	8.8
Secondary (9-12)	87	16.7
12+	158	30.3
Father's educational status		
Illiterate Read and write Primary (1-8) Secondary (9-	$37 \ 105 \ 42$	$7.1 \ 20.2 \ 8.1 \ 15$
12) 12+	$78 \ 206$	28.4
Mothers Occupation		
House wife Employed Merchant Farmer Others?	208 157	$39.9 \ 30.1 \ 22.6$
	$118 \ 7 \ 6$	$1.3 \ 1.2$
Family size		
<5 years	262	50.2
>=5years	259	48.8
Family income		
<500	4	0.8
500-1000	6	1.2
>1000	69	13.2
Don't know	442	84.8

Figure 5: E

Pills Condom Depo Implant IUCD Abistence Calender

19.60%

27.40%

45.90%

 $0.00\% \ 10.00\% \ 20.00\% \ 30.00\%$

40.00% 50.00%

56.80%60.00%

Figure 6: 64.10% 57.80% 55.90% 51.80% 50.90% 50.70% 44.10%

1

Ethiopia, 2013

Figure 7: Table 1 :

4	
Variable	Discussed on SRH
C	Yes No
Sex Male Female	$129(55.8\%) \ 102(44.2\%) \ 171(59\%) \ 11$
Age category 10-14	$13(59.1\%) \ 9(40.9\%)$
15-19	$287(57.5\%) \ 212(42.5)$
Living arrangement With both parents With mother only With father only With others ?	227(58.5%) 161(41.5%) 43(48.9%) 4
Grade 9 th 10 th 11 th 12 th	70(48.6%) $74(51.4%)$ $85(54.1)$ $72(45)$
Mothers' education No formal education Elementary 12+	$120(52.2\%) \ 110(47.8\%) \ 27(58.7\%) \ 1900$
Fathers' education No formal education Elementary 12+ Mothers' occupation	$112(57.4\%) \ 83(42.6\%) \ 24(57.1\%) \ 18$
Employed.	$93(59.2\%) \ 64(40.8\%)$
House wife	$113(54.3\%) \ 95(45.7\%)$
Merchant	68(57.6%) $50(42.4%)$
Others	26(68.4%) $12(31.6%)$
Fathers' occupation Employed Merchant Oth- ers ?	$147(58.1\%) \ 106(41.9\%) \ 105(56.1\%)$
Family size <5	$84(58.3\%) \ 60(41.7\%)$
>=5	$216(57.3\%) \ 161(42.7\%)$
Know at least one contraceptive method Yes No	59(52.7%) $53(47.3%)$ $241(58.9%)$ 16

who made communication regarding SRH matters, only very small number of them discussed about early sexual practice and sexual transmitted infections with their parents. The major reason respondents expressed for not discussing early sexual practice, contraception, sexual transmitted infections and unwanted pregnancy was

217 being ashamed.

Based on the findings of the study, the following recommendations were made.

This study showed that a significant proportion of adolescents did not discuss with their parents about at least one sexual and reproductive health matters. Therefore, the MOH, and other concerned stalk holders that are working in areas of reproductive health should strengthen existing strategies including provision of information, education and communication targeting adolescents and their parents to increase their awareness mainly about the impact of early sexual practice and its outcomes such as unwanted pregnancy and sexually transmitted infections (STIs).

As educational level of parents is a critical factor in advancing parent-adolescent communication about 225 SRH matters, policy makers and program managers should focus on encouraging parents to pursue formal 226 education. Moreover, health care providers should be effective in equipping adolescents with adequate sexual 227 and reproductive health information, being respectful and keeping their privacy during their visit to health 228 facility. Finally, further research with robust analytical studies on timing of parents-adolescents communication 229 230 regarding sexuality and reproductive health related issues and its effect on safe sexual behaviors. Abbreviations 231 EDHS: Ethiopian Demographic and Health Survey; SRH: Adolescent Sexual and Reproductive Health; MDG: Millennium Development Goals; AIDS: Acquired immune deficiency syndrome; RH: Reproductive Health; STI: 232 Sexual transmitted infection; WHO: World Health Organization. 233

²³⁴ .1 Competing interests

235 The authors declare that they have no competing interests.

²³⁶.2 VI. Acknowledgements

237 We would like to thank Mekelle University for financial assistance. We are also grateful to Tigray regional

- education bureau, supervisors, data collectors and study participants for their assistance and cooperation during the study.
- [A companion to the interagency field manual on reproductive health in humanitarian settings adolescent sexual and reproductive
 'A companion to the interagency field manual on reproductive health in humanitarian settings adolescent
- sexual and reproductive health toolkit for humanitarian settings'. *Save the children* 2009.
- [Ethiopia Demographic and Health Survey: Central statistical agency report ()] Ethiopia Demographic and
 Health Survey: Central statistical agency report, 2011. Ethiopia. Central Statistical Agency
- ²⁴⁵ [Federal democratic republic of Ethiopia (2004) National youth policy, Ministry of youth, Sport culture]
- Federal democratic republic of Ethiopia (2004) National youth policy, Ministry of youth, Sport & culture,
 Addis Ababa; Ethiopia.
- [Health in the united states, with special feature on en emergency care ()] Health in the united states, with special feature on en emergency care, 2012. Washington: Library of Congress. National center for health statistics
- [Helping parents improve adolescent health. Family health international. USA. Available at www.youthwg.org 15. World Health C
 Helping parents improve adolescent health. Family health international. USA. Available at www.youthwg.org
- 254 15. World Health Organization Regional Office for Africa Child and Adolescent Health (2012) Healthy and
- Happy Children, 2007. Interagency youth Working Group (Investing in our Common future. swizerland.
 WHO)
- [Improving the Health of Adolescents Young Adults: A Guide for States and Communities Atlanta. Available at ()]
 'Improving the Health of Adolescents & Young Adults: A Guide for States and Communities'.

http://nahic.ucsf.edu/wpcontent/uploads/2011/11/Complete2010Guide.pdf Atlanta.

- Available at 2004. Centers for Disease Control and Prevention, National Adolescent Health Information
 Center, University of California
- 262 [Progress in reproductive health research World health organization ()] 'Progress in reproductive health re-263 search'. http://www.WHO.int/reproductivehealth/hrp/progress/58/news World health organi-264 zation 2013.
- 265 [Strengthening the health sector response to adolescent health and development World health organization ()]
- 'Strengthening the health sector response to adolescent health and development'. http://www.who. int/maternal_child_adolescent/documents/cah_adh_flyer_2010_12_en.pdf World health organization 2010.
- [World health organization (2006) Helping parents in developing countries improve adolescents' health ()]
- 270 World health organization (2006) Helping parents in developing countries improve adolescents' health,

16 V. CONCLUSION AND RECOMMENDATIONS

- http://www.cdc.gov/nchs/data/misc/adolescent2007.pdf10 2007. (Adolescent health in the United States. Switzerland. WHO)
- [World health organization (2014) Maternal, newborn, child and adolescent health] World health organization (2014) Maternal, newborn, child and adolescent health,
- [World health organization. Introduction to adolescence to adolescent health (2012) Training Course in Sexual and Reproductive
 World health organization. Introduction to adolescence & to adolescent health (2012) Training Course in Sexual
- and Reproductive Health Research, http://www.gfmer.ch/SRH-Course-2012/adolescent-health/
- pdf/Introduction-adolescence-adolescent-health-WHO-2012.pdf 2012. Society for Public
- Health Education. (Adolescent Health: Planting Seeds for a Healthier Generation)