

Fungal and Yeast Involvement in Skin Diseases

Nwachukwu O.N¹, Onyeagba R.A² and Nwaugo V.O³

¹ Abia State University

Received: 8 December 2015 Accepted: 3 January 2016 Published: 15 January 2016

Abstract

A mycological investigation of skin scrappings, blood and aspirates from apparent clinical cases of cutaneous infection involving 100 patients was made. All samples were cultured on Sabouraud dextrose agar (SDA) for three weeks. A total of 27(43.5

Index terms— fungi, yeast, skin disease.

1 Fungal and Yeast Involvement in Skin Diseases

Nwachukwu O.N ?, Onyeagba R.A ?, Nwaugo V.O ?, Ugbogo O.C ? & Ulasi, A.E ¥ Abstract-A mycological investigation of skin scrappings, blood and aspirates from apparent clinical cases of cutaneous infection involving 100 patients was made. All samples were cultured on Sabouraud dextrose agar (SDA) for three weeks. A total of 27(43.5%) and 18(29.0%) fungi and yeasts respectively were isolated from the skin. Blood cultures yielded 8(9.9%) yeast species only from the patients while culture of aspirates showed more growths of yeasts (2 isolates) than filamentous fungi (1 isolate). The following dermatophytes and filamentous fungi were isolated from the skin: *Microsporum ferrugineum*, *Trichophyton mentagrophytes*, *T. rubrum*, *T. verrucosum* and *T. schoenleinii*; *A. fumigatus*, *A. niger*, *A. flavus* and *Penicillium* species. *Trichophyton mentagrophytes* had the highest occurrence (50%) while *Microsporum ferrugineum* and *Trichophyton verrucosum* were least in occurrence(9.1%) each. Five yeast species, *Candida albicans*, *C. tropicalis*, *C. krusei*, *Rhodorulla* species and *Torulopsis* species were also isolated from skin lesions. *Torulopsis* species was the most occurring yeast whereas *Candida krusei* and *Rhodotorulla* species showed the lowest occurrence (5.6%) each. *Torulopsis* species and *Candida tropicalis* were recovered from blood and aspirate of patients. Prevalence of fungi and yeast was similar in male and female patients. Fungi and yeast are involved in skin diseases.

Keywords: fungi, yeast, skin disease.

2 I. Introduction

kin is the most accessible organ of the body, the one most easily traumatized and therefore frequently subjected to infection. Normal human skin is colonized by large numbers of microorganisms that live harmlessly as commensals on its surface 1 . Skin diseases therefore is a complex subject involving diverse microorganisms that exhibit varying aetiological and pathogenic mechanisms.

Fungal diseases of the skin are a common public health problem worldwide. The prevalence of skin fungal diseases is expected to reach 20-25% of the world's population and its incidence continues to rise 2 . Fungal skin diseases constitute an important clinical and public health problem in tropical areas of the world where they are rarely managed 3 .

There are many species of fungi that cause skin diseases in man. These are mainly Dermatophytes (*Trichophyton* species, *Epidemophyton* sp and *Microsporon* so), *Malassezia furfur* and *Candida* species and less commonly *Aspergillus* species, *Trichothecium roseum*, *Cladosporium* sp and *Fusarium* sp 4 .

The diseases caused by fungi (mycoses) can be clinically classified as superficial, deep or systematic mycoses 5 . Dermatophytes are the most important microorganism which cause superficial mycosis and the lesions are characterized by circular disposition, desquamation alopecia and erythema of the edges 6 . They invade and destroy the skin, hair and nails. These diseases have been reported in various studies from developing countries as the most common dermatosed 7,8 . They are also responsible for most of the skin infections among school children 9 .

Fungal skin diseases constitute the majority of skin conditions seen by physicians in primary, secondary and tertiary health care centres in Nigeria [11][12] .

They make the individual uncomfortable, unsightly and present a cosmetically poor appearance [13] . The situation is enhanced in a tropical country like Nigeria by warm humid weather, crowded living and poor sanitary conditions which are prevalent and support infection on human skin [14] .

This study therefore seeks to investigate the rate of involvement of fungi and yeast in diverse kinds of skin diseases that present.

3 II. Materials and Methods

4 a) Study Area

This prospective study was carried out at Nnamdi Azikiwe University Teaching Hospital (NAUTH), Nnewi. NAUTH is the largest tertiary hospital and referral centre in Anambra State Nigeria.

5 b) Subjects

Individuals aged 1 year and older with varied skin disease referred to the mycology section of microbiology laboratory for further assessment were invited to participate and enrolled at the time of presentation to the laboratory.

6 c) Sample Collection

Skin scrapings, venous blood and aspirates (where applicable) from 100 patients with clinical cases of cutaneous infection were collected after thorough physical examination. The clinically apparent lesions described as dry, scaly and matted or seranguinous were cleaned with 70% alcohol. Epidermal scales at the active edges of the lesions were scrapped using sterile surgical blades. The scrapings were collected in a piece of paper, carefully folded and then placed in an envelope for storage in air-tight containers.

Aspirates were taken from pustular lesions or nodules where present. After cleaning the arm, a tourniquet was applied to dilate the upper arm and 1ml of venous blood was withdrawn from the cubital region of the forearm. These samples were properly labeled and a brief history of the disease taken.

7 d) Processing of Samples

Each skin scraping was inoculated onto Sabouraud Dextrose Agar (SDA) incorporated with 0.05mg/ml Chloramphenicol and Streptomycin 40mg/ml. Duplicate inoculations were made. They were incubated at room temperature (25-28 °C) and at 37 °C respectively for 21 days, examining daily for fungal and / or yeast growths.

Aspirates were also inoculated on SDA and incubated at 37 °C for 48 hours.

One millilitre (ml) amount of blood specimens were introduced aseptically into 5ml of Sabouraud Dextrose Broth contained in MacCathney bottles. Duplicate inoculations were made. They were incubated at 37 °C for 48 hours before being subcultured onto solid medium, SDA. Incubation of blood cultures and subsequent subculture onto SDA continued for 21 days after which negative cultures were discarded.

8 e) Characterization and Identification of Isolates

The mycological identification was based on macroscopic and microscopic examination of culture isolates. Macroscopic examination of dermatophytes was characterized by duration of growth, surface morphology and pigment production on the reverse [15] . Microscopic examination of fungal growth was observed with Lactophenol cotton blue stain. Nature of mycelium and conidia formation help to differentiate various genera and species [16] .

All cream to white-tan pasty colonies with characteristic yeast smell were stained by Gram's method and examined microscopically. Budding yeast cells of *Candida* species were identified by germ tube formation, sugar fermentation and sugar assimilation [17] .

Ethical approval for the study was obtained from the ethical committee of the hospital. All patients consented to participate in the study.

9 III. Results

A total of one hundred and fifty-seven samples consisting of 62 skin scrapings, 81 blood and 14 aspirates, collected from 100 patients were cultured. Culture of the skin scrapings revealed that 27(43.5%) isolates yielded fungi whereas 18(29.0%) of the isolates were yeast-like organisms. Eight (9.9%) of the eight-one blood samples yielded yeast-like organisms. Similarly, 1 (7.1%) of 14 aspirate samples cultured yielded a fungus and 2 (14.3%) yeast-like organisms (Table 1). Five species of dermatophytes were isolated from 100 patients who had superficial infections: *Microsporum ferrugineum*, *Trichophyton mentagrophytes*, *T. rubrum*, *T. verrucosum* and *T. schoenleinii*. *T. mentagrophytes* had the highest occurrence (50%) followed by *T. rubrum* (18.2%). Five yeast species including *Candida albicans*, *C. tropicalis*, *C. krusei*, *Rhodotorula* sp and *Torulopsis* sp were also recovered from skin lesions. *Torulopsis* sp was the most occurring yeast (44.4%) while *C. krusei* was the least occurring (5.6%) *Aspergillus*

niger was the most common filamentous fungi (44.4%) isolated from the skin whereas *A. flavus* was the least recovered (11.1%). *Torulopsis* sp (50%) and *Candida* tropicals (50%) were isolated from the blood as presented in Table 2 Fungi were isolated more from males (57.6%) than from females (42.4%). Similarly, yeasts were more frequently isolated from males (60.7%) than females (39.3%) Table 3. Different fungi and yeast species were isolated from various body sites (Table 4).

10 IV. Discussion

This study has revealed the skin as most susceptible to infections by fungi (43.5%) and yeast (29.0%) as compared to blood (0%) and (9.9%) respectively, lending credence to the work of Yahya et al. 18 that skin is the most accessible organ to infection. The result of the species of dermatophytes isolated from skin diseases which include *Microsporum ferrugineum*, *Trichophyton mentagrophytes*, *T. rubrum*, *T. verrucosum* and *T. schoenleinii* is in agreement with previous studies in Korea, Iran, India and Jos, Plateau State, Nigeria. [19][20][21][22][23] An earlier study carried out about 30 years ago in Eastern Nigeria 24 also isolated these dermatophytes. This implies that fungal infections are still highly prevalent in these areas. *Trichophyton mentagrophytes*, one of the several dermatophytes that cause cutaneous mycoses and *T. rubrum* were the most common dermatophytes isolated from skin lesions. This finding is in consonance with the work of Ta'ama et al 23 who recovered *T. mentagrophytes* constantly and that of Kannan et al 22 where *T. rubrum* was the most prevalent causative agent implicated in skin fungal infections.

Majority of dermatophytes were isolated from scalp (*Tinea capitis*) of patients. Among these were two members of a family. It had been reported. 2, 24,25 that frequent interchange, poor sanitary conditions, sharing of hair brushes, combs and hats have played some role in the spread of the disease. These conditions may be responsible for the observation made in the case of the two family members.

Non-isolation of *Trichophyton tonsurans* from any of the clinically observed lesions in this study is a deviation from earlier reports. 24,26 in Eastern and Northern parts of Nigeria where this fungus was frequently encountered. The relative small sample size of this study may have accounted for it.

In the screening for systemic involvements in patients with long standing skin diseases, *Candida tropicalis* (50% of the total yeast, isolates made), *Rhodotorula* species (37.5%) and *Torulopsis* species (12.5%) were isolated. Previous studies 5,27 had shown that *Torulopsis* species and *Candida* tropicals, both opportunistic pathogens, were capable of establishing themselves in the blood. The patient's defenses may have been weakened by some other processes for these pathogens to be recorded from blood. Petmy et al 28 reported that it may have been as a result of frequent usage of antibiotics, immunosuppressive drugs and various conditions like organ transplantations, lymphoma, leukemia and human immunodeficiency virus (HIV) infection. This study did not however, attempt to establish if there had been any previous but continuing disease, clinically or through laboratory diagnosis. However, some of the patients had been on antibiotics for long periods, a situation that may promote human infection by the yeasts. 17 In one of the cases studied, *Rhodotorula* species, was isolated from the skin of an infant (1 year old). This condition may be the result of certain complications such as napkin dermatitis or due to no clearly defined pre-disposing factor.

The pathogenic status of *Candida* tropicals and *Torulopsis* species was further highlighted in this study as these were isolated from aspirates.

Prevalence of fungal skin infections in males and females were similar (33.0% vs 28.0%). Similar studies 19,23 supports our finding showing that large numbers of people are often affected by fungal skin diseases irrespective of their gender.

11 V. Conclusion

Fungi and yeasts are involved in skin diseases. *Trichophyton mentagrophytes* was the most common dermatophyte while *Torulopsis* species was the highest occurring yeast involved in skin diseases. Prevalence in males and females were similar.

1

Nature of sample	No ex- amined	No positive for fungi (%)	No positive for yeasts (%)
Skin scrapings	62	27(43.5)	18(29.0)
Blood	81	0(0)	8(9.9)
Aspirate	14	1(7.1)	2(14.3)
Total	157	28(17.8)	28(17.8)

Figure 1: Table 1 :

2

Fungal species	Skin	Blood Aspirates	
Dermatophyte			
Microsporum ferrugineum	2(9.1)	-	-
Trichophyton mentagrophytes	11(50)	-	-
T. rubrum	4(18.2)	-	-
T. verrucosum	2(9.1)	-	-
T. schoenleinii	3(13.6)	-	-
Total	22	0	0
Filamentous fungi			
Aspergillus niger	4(44.4)	-	-
A. fumigates	2(22.2)	-	-
A. flavus	1(11.1)	-	-
Penicillium sp	2(22.2)	-	1(100)

Figure 2: Table 2 :

3

Gender	No of patients examined	No positive for yeast	No positive fungi (%)
Males	54	17(31.5)	19(35.2)
Females	46	11(23.9)	14(30.4)
Total	100	28(28.0)	33(33.0)

Figure 3: Table 3 :

4

Skin disease	Body site affected	Fungi isolated	Yeast isolated
Tinea capitis	Scalp	Microsporum ferrugineum Trichophyton mentaphytes T. rubrum Penicillium sp	Candida tropicalis Rhodotorulla sp
Tinea corporis	Hand, trunk face, groin leg, buttocks	T. mentagrophytes T. verrucosum Penicillium sp A. fumigatus ,A.niger	Candida krusei C. albicans Torulopsis sp
Tinea imbricate	Breast	T. rubrum	-
Paronychia (Tinea unguium)	Toe web Toe nail	T. mentagrophytes T. schoenleinii Penicillium sp T. mentagrophytes A. niger	- C. albicans C. tropicalis

Figure 4: Table 4 :

[Livingstone and London] , Churchill Livingstone , London . p. .

[Cheesbrough and Mycology In ()] , M Cheesbrough , Mycology In . *Medical Laboratory Manual for Tropical Countries* 1984. University Press. 2 p. .

[Souissi et al. ()] 'A study of skin diseases in Tunis. An analysis of 28,244 dermatological outpatient cases'. A Souissi , F Zeglaoui , B Zouari . *Acta Dermatoves APA* 2007. 16 p. .

[Yahya ()] 'Change in patterns of skin disease in Kaduna, north-central Nigeria'. M Yahya . *Int J Dermatol* 2007. 46 p. .

[Olasode et al. ()] 'Cutaneous infections in patients presenting in a skin clinic in the tropics'. O A Olasode , A A Otu , E B Henshaw . *Int J Trop Med* 2009. 4 p. .

[Ta'ama et al. ()] 'Dermatophytes and other agents of human skin infections in Jos, Plateau State'. L Ta'ama , A A Makinde , L H Lombin . *Nigeria J Med Lab Sci* 2001. 10 p. .

[Duguid et al. ()] J P Duguid , B P Marimion , Rha Swain . *Pathogenic fungi In: Medical Microbiology 13 th Ed*, 1978.

[Kim et al. ()] 'Epidemiological characterization of skin fungal infections between the years 2006 and 2010 in Korea'. S H Kim , H C Seung , K Y Seung . *Osong Public Health Res Perspect* 2015.

[Ameen ()] 'Epidemiology of superficial fungal infections'. M Ameen . *Clin Dermatol* 2010. p. .

[Egere and Gugnani ()] 'Etiology of dermatophytosis in Eastern Nigeria: a 7 year study'. J U Egere , H C Gugnani . *Mykosen* 1987. 30 p. .

[Gupta and Gupta ()] 'Evaluation of the incidences of dermatophilic infection in Rajastahan, India'. S Gupta , B L Gupta . *Int J Med Med Sci* 2013. 5 p. .

[Greenwood et al. ()] D Greenwood , Rcb Slack , J F Peutherer . *Fungi In: Medical Microbiology 16 th Ed*, (Churchill Livingstone, London) 2002. p. .

[Grover and Roy ()] 'Harlickova B, Czaika VA, Friedrich M. Epidemiological trends in skin mycoses worldwide'. Wcs Grover , Lcp Roy . *Med J Armed Forces India* 2003. 2008. 59 p. . (Mycoses)

[Finegold and Martin] 'Laboratory diagnosis of mycotic infections'. S M Finegold , W J Martin . *Diagnostic Microbiology. 6 th Ed. 1982. The CV Mosby Company USA*, p. .

[Pelczar et al. ()] M J Pelczar , Ecs Chan , N R Krieg . *Fungi, molds and yeasts In: Microbiology 5 th Ed*, 1986. Oxford Press London. p. .

[Koneman and Roberts] *Practical laboratory mycology. 3 rd Ed 1985. Baltimore Williams and Wilkins*, E W Koneman , G D Roberts . p. P188.

[Oyedeyi et al. ()] 'Prevalence and patterns of skin infections infestations among primary school pupils in Ijeshaland'. O A Oyedeyi , O Onayemi , G A Oyedeji . *Nig J Paed* 2006. 33 (1) p. .

[Kannan et al. ()] 'Prevalence of dermatophytes and other fungal agents isolated from clinical samples'. P Kannan , C Janaki , G S Selvi . *Indian J Med Microb* 2006. 24 (3) p. .

[Ahmadinejad et al. ()] 'Prevalence of fungal skin infections in Iranian wrestlers'. Z Ahmadinejad , A Razaghi , A Noori . *Asian J Sports Med* 2013. 4 (1) p. .

[Emodi et al. ()] 'Skin diseases among children attending the children outpatient clinic at the University of Nigeria teaching hospital Enugu'. I J Emodi , A N Ikefuna , U A Duru . *Afr Health Sci* 2010. 10 p. .

[Nnoruka ()] 'Skin diseases in South-east Nigeria: a current perspective'. E N Nnoruka . *Int J Dermatol* 2005. 44 p. .

[Oninia ()] 'Superficial fungi skin infections. The bane of dermatoses in Nigeria'. O A Oninia . *Med Microb Diagn* 2014. 3 p. 4.

[Petmy et al. ()] 'Superficial mycoses and HIV infection in Yadunde'. J L Petmy , A J Larido , V T Kaptue , M Folefack . *J Eur Acad Dermatol* 2004. 18 (3) p. .

[Monod et al. ()] 'Survey of Dermatophyte Infections in the Lauseanne Area (Switzerland)'. M Monod , S Jaccoud , C Zaugg . *Dermatology* 2002. 205 p. .

[Santos et al. ()] 'The prevalence of common skin infections in four districts in Tomor-Leste: a cross sectional survey'. Dos Santos , M M Amaral , S Harmen , SP . *BMC Infect Dis* 2010. 10 p. 61.

[Bissek et al. ()] 'The spectrum of skin diseases in a rural setting in Cameroon (sub-Saharan Africa)'. A Z Bissek , E N Tabah , E Kouotou . *BMC Dermatol* 2012. 12 p. 7.

[Jacyle and Lawande ()] 'Tinea capitis in Northern Nigeria'. W R Jacyle , Abram E Lawande , RV . *Mykosen* 1982. 25 p. .

[Gugnani and Njoku-Obi ()] 'Tinea capitis in school children in Eastern Nigeria'. H C Gugnani , Anu Njoku-Obi . *Mykosen* 1986. 29 p. .