

# Childhood Hypocalcemia: The Aetiological Pattern

Nasir A. M. Al Jurayyan<sup>1</sup>

<sup>1</sup> King Saud University

*Received: 13 December 2015 Accepted: 4 January 2016 Published: 15 January 2016*

---

## Abstract

Background: Hypocalcemia is not that rare condition and could be a potentially life threatening. Identifying the etiology is important for successful management. Results: A total of 60 patients were seen in the period under review, December 1989 and June 2016, with childhood hypocalcemia. Twenty-seven (45.0

---

*Index terms*— aetiology, childhood, hypocalcemia.

## 1 I. Introduction

ypocalcemia is a potentially life-threatening metabolic disturbance. It can result in severe symptoms that require rapid management. Hypocalcemia occurs most commonly as a result of deficiency of hormone (PTH). Though, there are many other potential etiologies of hypocalcemia, one usually does not consider them seriously unless the most common cause is ruled out or unless the initial evaluation suggest another cause is likely.

In primary hypoparathyroid, an assay that measures intact circulating PTH will be low, while in virtually all other causes associated with hypocalcemia. PTH levels are elevated. [1][2][3][4] This article focuses upon the etiology of hypocalcemia beyond the neonatal period, seen in a major teaching hospital, King Khalid University Hospital (KKUH), Riyadh, Saudi Arabia over three decades, December 1989 to July 2016.

Then KKUH is the main teaching hospital of the King Saud University (KSU) and considered as one of the major referral hospitals in the region, and provides primary, secondary, and tertiary health care services for the local population and also receives patients referred from all over the country.

## 2 II. Materials & Methods

During the period under review, December 1989 to June 2016, all patients who were diagnosed, beyond the neonatal period to have hypocalcemia were retrospectively reviewed.

Detailed history, clinical manifestations and results of all the laboratory, radiological and ancillary investigations were obtained. The aetiological diagnosis was based on specific investigations as recommended.

## 3 III. Results

During the period under review, December 1989 and June 2016, a total of 60 patients beyond neonatal period were seen by the author in the pediatric endocrine service, King Khalid University Hospital, Riyadh, Saudi Arabia. Table, showed the aetiological diagnosis of the group. In 27 (45.0%) patients, parathyroid hormone (PTH) deficiency was found while rickets was the diagnosis in 25 (41.7%) patients. Celiac disease was diagnosed in 6 (10.0%) patients.

## 4 IV. Discussion

Hypocalcemia beyond neonatal period is not that rare. It varies from an asymptomatic biochemical abnormality to a life threatening conditions, depending on the duration, severity and rapidity of development. Hypocalcemia is caused by loss of calcium into circulation. In a community with high prevalence of consanguinity mating and increased incidence of autosomal disorders, (5,6) various forms of hypoparathyroidism exist and constitute the major cause. Simple hypoparathyroidism usually occurs sporadically, though an autosomal dominant pattern of

42 inheritance has been reported. In most cases the pathogenesis is unknown, but agenesis, partial or complete  
43 atrophy, and inflammatory damage of the parathyroid glands are possible mechanisms. However the diagnosis  
44 of isolated hypoparathyroidism cannot be made with certainty in childhood, since children who first appears  
45 to have this disorder often develop additional endocrine or immunological abnormalities later on. (7)(8)(9)  
46 Damage to the parathyroid glands is a well-established risk of neck surgery, especially during total or subtotal  
47 thyroidectomy. Permanent parathyroid deficiency occurs in about five to ten percent of subtotal thyroidectomies  
48 and is significantly more common after total thyroidectomy for malignant thyroid disease. Hypoparathyroidism  
49 is usually caused by an interference with the blood supply of the glands and is rarely due to complete ablation of  
50 the parathyroid tissue. Non-surgical damage to the parathyroid glands can occur as a result of massive doses of  
51 external irradiation. However, the parathyroids are relatively radiation resistant so definite hypoparathyroidism  
52 following treatment for thyroid disease is exceptionally rare.

53 A well-established relationship exists between magnesium and calcium homeostasis. Magnesium deficiency  
54 may lead to hypocalcemia by either PTH synthesis or release and end-organ, bone, refractoriness to the effects of  
55 PTH. (11) Rickets was the second most common cause of hypocalcemia in our study. The various forms shown  
56 in table. The clinical and radiological is highly variable, depending on the age, the etiology and the duration  
57 and severity of the disorder. (3,4,(12)(13)(14)(15) Vitamin D deficiency was common, however, derangement in  
58 Vitamin D metabolites or action is not that rare.

59 Celiac disease should be considered in patients with hypocalcemia of unknown etiology, especially because  
60 gastrointestinal symptoms may be absent or mild. Six (10%) patients in our series were diagnosed to have celiac  
61 disease. (16) In conclusion, this study showed that parathyroid hormone (PTH) deficiency (45%) and rickets  
62 (41.7%) were the most common causes of childhood hypocalcemia

## 63 5 V. Acknowledgement

64 The author would like to thank Mrs. Cecile M. Sael for typing the manuscript, and extends her thanks and  
65 appreciations to Miss Hadeel N Aljurayyan for her help and support.

## 66 6 Conflict of Interest

67 The author have no conflict of interest to declare.

## 68 7 Volume XVI Issue V Version I

1

Diagnosis	Number	(%)
-----------	--------	-----

Figure 1: Table 1 :

- 
- 69 [Holick] , M F Holick .
- 70 [Wharton and Bishop ()] , B Wharton , N Bishop . *Rickets Lancet* 2003. 362 p. .
- 71 [Vitamin and Deficiency ()] , D Vitamin , Deficiency . *N Engl J Med* 2007. 357 p. .
- 72 [Shoback et al. ()] , D Shoback , Clinical Practice , Eng . *J Med* 2008. 359 p. .
- 73 [Sanjad et al. ()] ‘A new Syndrome of Congenital hypoparathyroidism, severe growth failure and dysmorphicfea-  
74 tures’. S A Sanjad , N A Sakati , Abu Osba , Y K Kaddoura , R Milner , RD . *Arch Dischild* 1991. 66 p.  
75 .
- 76 [Ali et al. ()] ‘Autoimmune polyendocrine Syndrome type I’. Al Ali , AO , Al Ayed , OA , Al Issas , A N  
77 Aljurayyan , N A Al Jurayyan . *Int J recent Sci Res* 2015. 6 (7) p. .
- 78 [Jurayyan et al. ()] ‘Celiac disease presenting as rickets in children’. Al Jurayyan , Azn , A L Otaibi , AM , Al  
79 Jurayyan , R N Al Assiri , A M Al Jurayyan , NA . *Pediatrics me* 2009. 14 (3) p. .
- 80 [Miller and Anthony ()] ‘Genetic disorders of Vitamin D biosynthesis’. W L Miller , A P Anthony . *Endocrinol-  
81 MetabClin N Am* 1999. 28 p. .
- 82 [Marx ()] ‘Hypoparathyroidism and hyperperithyroidism disorders’. S J Marx . *N Engl J Med* 2000. 343 p. .
- 83 [Anast et al. ()] ‘Impaired release of parathyroid hormone in magnesium deficiency’. C S Anast , J L Winnacker  
84 , L R Forte , T W Burns . *J clin Endocrine Metab* 1976. 42 p. .
- 85 [Linglart and Bastepe ()] ‘Lupponer HSimilar clinical and laboratory findings in patients with symptomatic  
86 autosomal dominant and sporadic pseudohypoparathyroidism type 1b despite different epigenetic changes  
87 at GNAS locus’. A Linglart , M Bastepe . *Clinic Endocrinol* 2007. 67 p. .
- 88 [Pettifor ()] ‘Nutritional and drug induced rickets and osteomalacia. In Favus MJ ed. Nutritional and induced  
89 rickets and osteomalacia: Primer on the metabolic and bone disorders and disorders of bone metabolism. 5  
90 th ed. Washington DC’. J M Pettifor . *American Society for Bone Mineral and research* 2006. p. .
- 91 [Thakker (ed.) ()] *Parathyroid disorders and diseases altering calcium metabolism*, R V Thakker . Worrall D,  
92 Cox T, Firan J, Benz E (ed.) 2003. Oxford University press.
- 93 [Jurayyan et al. ()] ‘Pseudohypoparathyroidism type 1b: A rare cause of femoral neck fracture’. Al Jurayyan ,  
94 N A Mohammed , S Saad-Uddin , MF , Al Jurayyan . *J Taiba university Med Sci* 2011. 6 (1) p. .
- 95 [Jurayyan et al. ()] ‘Rickets and osteomalacia in Saudi children and adolescents attending endocrine clinic,  
96 Riyadh Saudi Arabia’. Al Jurayyan , NA , MohamedS , Al Issa , SD , Al Jurayyan . *Sudan J Pediatric*  
97 2012. 12 (1) p. .
- 98 [Saedi-Wong et al. ()] ‘Socioeconomic epidemiology of consanguineous mating’s in Saudi Arabian population’. S  
99 Saedi-Wong , Al Frayh , A R Wong , Hym . *J Asian Afri* 1989. 24 p. .
- 100 [Jurayyan and Osman ()] ‘The increased prevalence of Congenital adrenal hyperplasia: The roles of consanguin-  
101 ity and multiple siblings involvement’. Al Jurayyan , Nam Osman , HA . *Eur J Res in Med Sci* 2015. 3 (1)  
102 p. .