

Major Causes of Liver Condemnation and Associated Direct Financial Losses in Bovine Saughtered at Assela Municipal Abattoir Arsi, South Eastern Ethiopia

Mukarim Abdurahaman¹, Hussein Aman² and Gobu Boru³

¹ Jimma University

Received: 6 December 2015 Accepted: 4 January 2016 Published: 15 January 2016

7

Abstract

A study was conducted from October, 2013 to March, 2014 on cattle slaughtered at Assela municipal abattoir with the aim of determining major causes of liver condemnation and to estimate the direct financial losses attributed to the condemned liver. Ante mortem and post mortem inspection procedures were followed throughout the study and abnormalities encountered were recorded. A total of 384 cattle were examined at ante mortem and 9 animals were found to have abnormalities/conditions like lameness, blindness, rough hair, amputated tail and branding. Out of 384 cattle slaughtered, 274 (71.4

16

Index terms— abattoir, assela, cattle, condemnation, financial loss, liver.

1 Introduction

One of the losses from endemic disease is expressed in terms of organ condemnation. The most commonly affected organs are liver and lung due to fasciolosis (for liver) and hydatidosis (for both) (Teka, 1997). Each year significant loss results from death of animals, inferior weight gain, condemnation of edible organs and carcasses at slaughtered. This production loss to the livestock industry is estimated at more than 900 million USD annually (Abebe, 1995).

A thorough meat inspection procedure requires two-steps namely antemortem and postmortem inspection. The importance of ante mortem inspection in the abattoir has long been recognized in an attempt to avoid the introduction of clinically diseased animals into the slaughterhouse. Ante mortem inspection should be done within 24 hours of slaughter and repeated when slaughter has been delayed over a day (Teka, 1997; Mezgebu, 2003).

Postmortem inspection is screening or sorting process to separate the normal from abnormal. It is the center around which meat hygiene revolves since it provides information indispensable from the scientific evolution of clinical signs and pathological process that affect wholesomeness of meat. Routine post mortem inspection of carcass and organs should be carried out as soon as possible after completion of dressing (Gracey, 1986).

The final judgment to be taken with an organ and carcass or parts of a carcass is based on the total evidence produced by observation, palpation, incision, smell, and any ante mortem signs (Teka, 1997). It is necessary to be aware the extent to which the public is exposed to certain zoonotic diseases detected in abattoirs and financial losses attributed to condemnation of affected organs and carcass (Nfi and Alonge, 1987). This is due to the fact that, meat is the main source of particular importance to the public such as hydatidosis, fasciolosis, tuberculosis and cysticercosis (Sirak, 1991).

Bovine liver is one of the largest visceral organs in the animal body which performs numerous functions and very rich sources of vitamins and minerals ??Radostitis et al., 2007). The tissue is much sought by consumers due to its palatability and easy to consumption. However, it is one of the most commonly condemned visceral organs during routine meat inspection (Phiri, 2006).

9 III. FINANCIAL LOSS ASSESSMENT

43 Parasites in the tropics are responsible for greater losses to the meat industry than any other diseases (Jobre
44 et al, 1996). Similarly like many other tropical countries of Africa, it is well known that parasitic diseases are
45 among the major factors responsible for the low productivity of livestock in Ethiopia (Abebe, 1995;Jobre et al.,
46 1996).

47 Cystic echinococcosis/Hydatidosis is a parasitic infection caused by larval stage of *Echinococcus granulosus*,
48 which is small tapeworm, for which dogs and other canids are typical definitive hosts. The adult parasite found
49 in small intestine of carnivores while the metacestode (hydatid cyst) is found in different organs of a wide varieties
50 of herbivores including (sheep, goats and cattle), pig, horse and man (Soulsby, 1982).

51 Fasciolosis in cattle is chronic wasting disease caused by the presence in the liver and bile ducts respectively
52 of immature and adult Trematode of the genus *fasciola*. The disease is found in vast areas of the world with
53 the smaller *fasciola hepatica* (3.5x1cm) in temperate countries and the large *fasciola gigantea* (7.5cm) in tropical
54 regions (Andrews et al., 1999).

55 Various researchers have undertaken studies at abattoir surveys to determine the prevalence and economic
56 importance and cause of meat condemnation (hydatidosis, Fasciolosis,Cystercosis) as these are mainly to be of
57 major economic and public health importance in meat inspection (Jobre et al., 1996). Therefore, the objective
58 of this study was to identify the major causes of liver condemnation in cattle slaughtered in Assela municipal
59 abattoir and to estimate the magnitude of direct economic losses attributed to this condemnation.

60 2 II.

61 3 Materials and Methods

62 4 a) Study area

63 The study was conducted at Assela municipal abattoir, from October 2013 to March 2014. Assela is located 175Km
64 South East of Addis Ababa at an altitude of 2350-2400 meters above sea level and has a climatic condition of
65 "Woynadega". The annual average rainfall is 1300-1350mm. A day and night temperature of the area ranges
66 from 10-25 0 c and 10-20 0 c respectively. The area has a bimodal rainfall occurring from March to April (short
67 rainy season) and from July to October (long rainy season) with mean annual rain fall of 1300-1350mm with the
68 relative humidity of 43-60 0 c (CSA, 2009).

69 5 b) Study animals

70 A total of 384 randomly selected cattle were inspected at Assela municipal abattoir. Out of which, 373 (97.14%)
71 were males and only 11 (2.86%) were females. From the total of 384 cattle slaughtered, 348 were old and 36 were
72 adults. Majorities of cattle came to Assela municipal abattoir for slaughters were originated from the market
73 places of Sagure, Kersa and Assela.

74 6 c) Study Design

75 A cross-sectional study was employed to identify the major cause of liver condemnation and to evaluate the direct
76 financial losses. The study animals were selected using simple random sampling method by taking the age, body
77 condition and origin of the animals into consideration. The desired sample size for this study was calculated
78 by using the formula given by Thrushfield (1995) with 95% confidence interval, 5% absolute precision and 50%
79 expected prevalence

80 7 d) Study Methodology i. Ante-Mortem Examination

81 Ante-mortem examination was conducted on individual animals, while the animals were entering in to the lairage
82 and in mass after they entered into the lairage. Both sides of the animals were inspected at rest and in motion.
83 Moreover, the general behavior of the animals, sign of diseases and abnormality of any type were recorded
84 according to the standard antemortem inspection procedures (Gracey, 1986). Following the judgment guideline
85 by ??FAO, 2003), animals fit for human consumption were allowed for slaughter.

86 8 ii. Post-mortem Examination

87 Post-mortem examination involved visual inspection, palpation and making systemic incision of liver to look
88 for the presences of cysts, adult parasites and other abnormalities. Pathological lesions were differentiated and
89 judged based on (FAO, 2003) guidelines on meat inspection for developing countries.

90 9 iii. Financial Loss Assessment

91 The direct financial losses due to liver condemnation from market were considered. The analysis was based on
92 the annual slaughter capacity of the abattoir considering market demand, the current average price of one liver in
93 Assela Butcherhouse and the rejection rate of liver. The direct financial loss incurred due to liver condemnation
94 was estimated by using the formula indicated below (Ogunrinade and Adesoke, 1982).

95 10 Results

96 11 a) Ante-Mortem Examination

97 Of the total 384 cattle examined at antemortem, 9 cattle were found to have the abnormalities listed below (Table
98 1). Some abnormalities encountered during antemortem inspection were rough hair coat, lameness, branding on
99 skin, blindness and amputated tail. Comparison of rejection rate of liver was carried out for different age, animal
100 origin and sex groups. From the total of 384 sampled animals, 340 were from sature while only 44 animals were
101 from (Assela and near surrounding). The proportion of liver condemnation was 71.8% and 68.2% from sature
102 and (Assela & near surrounding) respectively. However, no statistically significance difference was observed in
103 the liver condemnation rate between the two areas.

104 Regarding sex, the overall liver condemnation rate was 71.8% in male and 54.5% in female and no statically
105 significant difference was observed between the two sexes (Table 3). This study also showed that a highest liver
106 condemnation rate in older age groups (> 5 years) than adults (< or = 5 years). The rejection rate was 73.3%
107 in old and 52.8% in adult (Table 3). From the analysis, it was observed that as age increase, the rejection rate
108 was also found to increase. This difference in the rejection rate between the age groups was significant (Table
109 3). The annual slaughter rate of the abattoir for the last three years were 6800, 6900 and 6998 in 2010, 2011
110 and 2012, respectively. So the direct annual financial loss due to rejection of liver was calculated based on the
111 price of a liver at Assela (Table 4). By using necessary information and formula, the annual direct financial loss
112 incurred due to condemnation of liver was calculated to be 418,761 ETB or 22,040 US\$ per annum.

113 12 Discussion

114 In the present study, routine ante-mortem and post-mortem inspection was carried out to detect any abnormalities
115 encountered in Assela municipal abattoir. Branding, lameness, rough hair, blindness and amputated tail were
116 found in some animals during ante mortem examination and animals with such abnormalities were passed for
117 slaughter by considering that the problems were localized and simply related to (rough or low management
118 system) and not due to bad pathological condition.

119 From the total of 384 cattle slaughtered, 274 (71.4%) liver were condemned due to various causes. This finding
120 is in agreement with one of previous study, where 66.55% liver was condemned in Kombolcha ELFORA industrial
121 abattoir (Nurit et al., 2012). But slightly higher than 53.7% that have been done in Kombolcha (Jemal, 2009)
122 and 51.95% in Mekelle (Shegaw et. al., 2009). On the other hand, it is significantly higher than 31.1% and
123 17.61% reported by (Yifat et. al., 2011) in Gonder and (Mellau et. al., 2011) in Tanzania respectivitly.

124 In the present study, the major cause of liver condemnation were hydatidosis (64.6%) followed by fasciolosis
125 (20.8%). In the present study, the rejection rate of liver due to hydatidosis is 64.6%, which is highly greater than
126 the findings of (Nurit et. al., 2012) with 14.2% and (Jemal, 2009) with 9.2% in both are at Kombolcha ELFORA
127 industrial abattoir, 4.2% in Tanzania (Mellau et. al., 2011) and (Yifat et. al., 2011) with 3.7% in Gondar.

128 The rejection rate due to faciolosis is high when it is compared with the rejection rate of 12.7, 14.05 and 8.6%
129 by (Fufa et al., 2009)] at Welaita Sodo, (Swai and Ulicky, 2009)] at Hawi and (Mellau et al., 2011) at Tanzania,
130 respectively. On the other hand it was slightly similar with 24.32% by (Gebretsadik et al., 2009) at Mekelle, but
131 significantly lower than 68.7% in Kombolcha (Jemal, 2009) and 86.4 % in Gondar (Yifat et. al., 2011). This may
132 be due to climate and ecological condition of the study areas considered.

133 The finding of 1.5% of liver condemnation due to cirrhosis was significantly lower than 16.06% reported by
134 (Nurit et al., 2012) at ELFORA and (Raji, et al., 2010) with 10.4% at Zaria abattoir. However, it is almost
135 similar with 1.1% reported by (Yifat et al., 2011) at Gondar. A 0.7% liver condemnation due to abscess was
136 recorded in the present study. This is a little bit smaller than the report by (Cadmus and Adesokan, 2009) with
137 2.9% condemnation rate of liver due to hepatic abscess in western Nageria and (Ahmedullah et al., 2007), who
138 reported 3.8% liver condemnation rate in Bangladesh. On the other hand it was similar to the report of (Mellau
139 and Nongaond, 2010) with 1.1% at Arusha abattoir.

140 Calcifications were also among the lesions which significantly contributed to the liver condemnation in the
141 study abattoir. During this study, 7.7% of liver were condemned as a result of calcification, which is relatively
142 higher than the finding of (Mellau and Nongaond, 2010) which is 1.9% at Arusha, Tanzania. On the other hand,
143 it is similar with the finding of (Nurit et. al., 2012) with 8.18% done at ELFORA.

144 The analysis of the result on the bases of age indicated the total liver rejection rate was higher in older animals
145 and a significant difference was observed between the two age groups. This may be due to most of liver diseases
146 are chronic and the older animals are mostly affected by many diseases.

147 The direct financial loss incurred as result of condemnation of liver in the present study was 418,761 ETB or
148 22040 US\$ per annum. This is so much higher than 1800 US\$ per annum due to fasciolosis liver condemnation
149 reported by (Mwabonimana, 2008) at Arusha abattoir, Tanzania.

150 V.

151 13 Conclusion

152 In general, liver condemnations as have been reported in this study impact negatively on the economic status of
153 the traders and the livestock industry at large. This constituted a substantial loss to the economy of the slaughter

13 CONCLUSION

154 stock owners under study as such an amount of money would have been harnessed into livelihood improvements.
155 Though infected livers were condemned and rendered unfit for human consumption, there exist some public
156 health threats from animals slaughtered at the abattoir due to the possibility of some missed cases as a result
157 of poor cooperation between butchers and meat inspectors and other malpractices including hiding of infected
158 meat from meat inspectors to avoid economic losses on their side. Indeed, the condemnation of cattle livers
159 at slaughterhouse in Assela municipal abattoir represents a significant economic loss. Some of the conditions
160 described however can be prevented. Cases of hydatidosis could be reduced by better control of stray dogs. Since
161 most liver conditions were caused by parasites, deworming programmes coupled with good animal husbandry
162 would likely be effective in lowering their incidence. Some of the limitations, however; encountered in this study
163 included the use of only gross pathology in the diagnosis of the diseases, thus only those diseases with gross
164 pathological lesions that are pathognomonic were likely to be diagnosed. In spite of the limitation mentioned,
165 the public health implications of the quantity of infected livers condemned at Assela municipal abattoir on the
166 consumers and the role which postmortem inspection plays in safeguarding the health of the public cannot be
167 overemphasized. Therefore, there is a need for adequate meat inspection in Assela municipal abattoir in order
168 to reduce wastages, identify diseases and thereby minimize associated public health risks.

VI.

1

Abnormalities	No. of animals with disease condition	Judgment
Lameness	2	Passed for slaughter but with precaution
Blindness	1	Passed for slaughter but with precaution
Rough hair coat	4	Passed for slaughter but with precaution
Branding on skin	1	Passed for slaughter but with precaution
Amputated tail	1	Passed for slaughter but with precaution
Total	9	Passed for slaughter

b) Post-Mortem Examination

Out of 384 cattle slaughtered in Assela municipal abattoir, 274 livers were totally condemned due to various reasons (Table 2). The current study revealed the overall proportion of liver condemnation

rate due to various pathological findings was 71.4%. occurrence of hydatidosis was the highest (64.6%); followed by fasciolosis (20.8%), calcification (7.7%), coinfection of hydatid cyst with fasciolosis (4.7%), cirrhosis (1.5%) and liver abscess (0.7%).

Figure 1: Table 1 :

2

Major cause of liver condemnation	Frequency	proportion
Hydatid cyst	177	64.6
Fasciolosis	57	20.8
Calcification	21	7.7
Coinfection of hydatid cyst with fasciola	13	4.7
Cirrhosis	4	1.5
Abscess	2	0.7
Total	274	100

Figure 2: Table 2 :

3

Variables		No. exam	No. (+)	%	χ^2	p- value
Origin	Sagure	340	244	71.8	0. 245	0.621
	A&S	44	30	68.2		
Sex	Male	373	268	71.8	1.565	0.211
	Female	11	6	54.5		
Age	old	348	255	73.3	6.707	0.010
	Young	36	19	52.8		
BCS	Good	250	184	73.6	1.768	0.184
	Medium	134	90	67.2		

A&S = Assela and near surrounding

No. exam = number of animals examined

No. (+) = number of positive animals

BCS = body condition score

Figure 3: Table 3 :

4

Figure 4: Table 4 :

170 .1 Acknowledgments

171 The authors acknowledge Jimma University College of Agriculture and Veterinary Medicine for supporting the
172 study

173 [Gracey (ed.) ()] , J F Gracey . Meat hygiene.8 th ed. Baillere Tindal. London, Philadel Phia. Toronto (ed.)
174 1986. p. .

175 [Nfi and Alonge ()] *An economic survey of abattoir data in Fako division of south west, Cameron, Bulletin Animal*
176 *Health and Production in Africa*, A Nfi , D O Alonge . 1987. 35 p. .

177 [Swai and Ulicky ()] 'An evaluation of the economic losses resulting from condemnation of cattle livers and loss
178 of carcass weight due to fasciolosis: A case study from Hai town abattoir'. E S Swai , E Ulicky . *Livestock*
179 *Res. Rural Dev* 2009. 21 (11) .

180 [Ogunrinade and Adesoke ()] 'Bovine fasciolosis in Nigeria, intercurrent parasitic and bacterial infection'. A
181 Ogunrinade , G O Adesoke . *Trop. Alm. Health Prod* 1982. 14 p. .

182 [Fufa et al. ()] *Bovine fasciolosis: coprological, abattoir survey and its economic impact due to liver condemnation*
183 *at Soddo municipal abattoir, Southern Ethiopia. Tropical Animal Health and Production*, A Fufa , A Loma ,
184 M Bekele , R Alemayehu . 2009. 42 p. .

185 [Mwabonimana ()] *Cattle liver condemnation at Arusha meat company Ltd, Tanzania: causes and its financial*
186 *implication. A Research Paper Submitted in Partial Fulfilment of the Requirements for the Degree of*, M
187 F Mwabonimana . 2008. Morogoro, Tanzania. p. . Master of Preventive Veterinary Medicine of Sokoine
188 University of Agriculture

189 [Sirak ()] 'Cause of organ condemnation at Bahir Dar abattoir'. A Sirak . *Proceeding of the fourth national*
190 *livestock improvement Conference, Institute of Agricultural Research*, (eeding of the fourth national livestock
191 improvement Conference, Institute of Agricultural ResearchAddis Ababa, Ethiopia) 1991. p. .

192 [Central Statistical Authority livestock number of breed, age, sex and purpose report on livestock Characteristics (private peasant
193 'Central Statistical Authority livestock number of breed, age, sex and purpose'. *report on livestock*
194 *Characteristics (private peasant holdering) statistical bulletin*, (Addis Ababa) 2009. II.

195 [Phiri ()] 'Common conditions leading to cattle carcass and offal condemnation at south west province'. A M
196 Phiri . *Cameroon. Bull. Anium. Hlth. Prod. Afr* 2006. 35 (3) p. .

197 [Abebe ()] 'Current status of vet. Education and Animal Health research in Ethiopia, in vet'. G Abebe .
198 *Proceeding of an international IRLI, AA, and Pp*, (eeding of an international IRLI, AA, and Pp) 1995.
199 p. . (Impact of human health and nutrition in Africa)

200 [Jobre et al. ()] *Hydatidosis In three selected region in Ethiopia an assessment trial on its prevalence, Economic*
201 *and public health importance*, Y Jobre , F Labago , R Tiruneh , G Abebe , Phd Dorchies . 1996. 147 p. .
202 (Revne de medicine veterinary)

203 [Nurit et al. ()] 'Major Cause of Liver Condemnation and Associated Financial Loss at Kombolcha Elfora
204 Abattoir'. M Nurit , Zerihun , M Serkalem . *European Journal of Applied Sciences* 2012. 4 (4) p. .

205 [Yifat et al. ()] 'Major cause of organ condemnation and financial significance of cattle slaughtered at Gondar
206 ELLFORA abattoir'. D Yifat , D Gedefaw , S Desie . *Global Veterinarian* 2011. 7 (5) p. .

207 [Mezgebu ()] *Major cause of organ condemnation in ruminants slaughtered at Gondar Abattoir, North western*
208 *Ethiopia, DVM thesis*, Y Mezgebu . 2003. Ethiopia. p. 17. FVM, AAU, Debre-Zeit

209 [Jemal ()] *Major causes of organ condemnation in cattle slaughter at Kombolcha ELLFORA meat factory abattoir*
210 *south Wollo*, T Jemal . 2009. Ethiopia. Haramaya University (DVM thesis)

211 [Manual Diagnosis on Meat Inspection for Developing Country, FAO/UN. Viale deile termai caralla FAO ()]
212 'Manual Diagnosis on Meat Inspection for Developing Country, FAO/UN. Viale deile termai caralla'. *FAO*
213 2003. p. 49.

214 [Teka ()] *Meat Hygiene and Food Hygiene principles and methods of food borne disease control with special*
215 *reference to Ethiopia Pp*, G Teka . 1997. p. .

216 [Shegaw et al. ()] 'Organ condemnation and economic loss at Mekelle municipal abattoir'. S Shegaw , K Ashwani
217 , A Kassaw . *Ethiopia. Global veterinarian* 2009. 48 p. .

218 [Raji et al. ()] 'Pathological condition and lesions observed in slaughtered cattle at Zaria abattoir'. M A Raji , S
219 O Solam , J A Ameh . *J. clinical pathology and Forensic Medicine* 2010. 1 p. .

220 [Ahmedullah et al. ()] 'Pathological investigation of liver of the slaughtered buffaloes in Barisal district'. F M
221 Ahmedullah , M G Akbor , M M Kaider , K Hossain . *Bangladesh. J. Vet. Med* 2007. 5 p. .

222 [Gebretsadik et al. ()] *Prevalence and economic significance of fasciolosisincattlein Mekelle Area of Ethiopia.*
223 *Tropical Animal Health and Production*, B Gebretsadik , B Kassahun , T Gebrehiwot . 2009. 41 p. .

224 [Radostits et al. ()] O M Radostits , C C Gay , K W Hinchcliff , P D Constable . *Veterinary Medicine: A*
225 *Textbook of the Diseases of Cattle, Horses, Sheep, Pigs and Goats, 10 th ed*, (Philadelphia, PA, USA) 2007.
226 Elsevier Health Sciences. p. .

13 CONCLUSION

227 [Mellau and Nongaond ()] 'Slaughter house survey liver lesions in slaughtered cattle, sheep and goats at Arasha,
228 Tanzania'. L S Mellau , E D Nongaond . *Journal of Vet Sci* 2010. 3 p. .

229 [Mellau et al. ()] *Slaughter stock abattoir survey of carcasses and organ/offal condemnations in Arusha Region,*
230 *northern Tanzania. Tropical Animal Health and Production*, B L Mellau , H E Nonga , E D Karimuribo .
231 2011. 43 p. .

232 [Soulsby ()] E J L Soulsby . *Helminths, arthropods and Protozoa of domesticated animals*, (London, UK, Pp)
233 1982. 7 p. . (th ed. Birilliare Tindall)

234 [Andrews ()] 'The life cycle of hepatica in fasciolosis Dalton'. S J Andrews . *CABI publishing walling ford*, J P
235 Ed (ed.) (UK, Pp) 1999. p. .