

Phytochemical Screening and in Vitro Antioxidant Activity of Aqueous Extract of Blighia Sapida Stem Bark

Amira Philip Olaniyi¹

¹ Federal Polytechnic,

Received: 9 December 2016 Accepted: 2 January 2017 Published: 15 January 2017

Abstract

Blighia sapida is a plant belonging to the family of sapindaceae. In this study we aimed to carry the phytochemical screening of aqueous extract of Blighia sapida stem bark and evaluate its in vitro antioxidant activity. It was found that the aqueous extract of Blighia sapida stem bark showed the presence of tannin, saponin, flavonoid, alkaloid, phenols and ascorbic acid. It also contains trace elements zinc and selenium. Furthermore it showed some scavenging activity but not as such could be compared with the various standards used except for nitric oxide scavenging activity. These are indications that the aqueous extract of Blighia sapida contains some physiologically significant secondary metabolites and also possesses appreciable in vitro antioxidant activity.

Index terms— blighia sapida, phytochemicals, antioxidant, metabolites.

1 I. Introduction

Phytochemicals are naturally occurring in the medicinal plants, leaves, vegetables and roots that have defense mechanism and protect from various diseases. In fact, phytochemicals (or antioxidants) such as phenolic compounds (tocopherols, flavonoids and phenolic acids), nitrogen compounds (alkaloids, chlorophyll derivatives, amino acids and amines), carotenoids and ascorbic acid (Hall and Cuppett, 1997; Larson, 1988) compounds inhibit, or delay the oxidation of other molecules by inhibiting the initiation or propagation of oxidizing chain reaction.

Blighia sapida is a plant belonging to the family of Sapindaceae. It is commonly known as ackee. In Nigeria, it is called Gwanja Kusa (Hausa), Isin (Yoruba) and Okpu (Igbo). It is an evergreen tree of about 33 to 40ft (10-12m) with a dense crown of spreading branches. It is rather hand usually with a short trunk to 6ft (1.8m) in circumference. Its bark is grey and nearly smooth. The leaves are compound with three to five pairs of oblong, ovate-oblong, or elliptical leaflets 1.5-3.0cm long. The seed of the fruit is not edible, whereas the fleshy aril is edible. The fruit is known to contain Most of the earlier studies on Blighia sapida have been on the nutritional qualities of the root (Abolaji et al, 2007) and the leaves as a dry season feed resource for West African dwarf goats in the Northern savanna zone of Nigeria (Aderinola et al, 2007). The repellent potential of the fruit part components against stored-product insect pests (Khan and Gumbs, 2003) as well as neutropenia and thrombocytopenia effects of the aqueous and lipid extracts of the unripe fruit have been investigated in mice (Gardiner et al, 1996). More recently, the physicochemical properties of the oil from the fruit of the species and toxicological evaluation of the oil -based diet in Wistar rats have been investigated (Oladiji et al, 2009).

Tree bark is an important component of African traditional medicine as herbal medicine is still the main source of health care for the majority of Africans and in particular, Nigerians. There has been increasing demand for the use of plant products with therapeutic activity. The high cost, availability, uncertainty of use during pregnancy and undesirable side effects of synthetic drugs or drugs from other animal sources are some of the factors leading to a strong preference for drugs of plants origin.

This study is thus aimed at investigating the phytochemicals and in vitro antioxidant activity of the readily available Blighia sapida stem bark.

2 II. Materials and Methods

Chemicals: All chemicals used were of analytical grade and items are products of BDH and Sigma Chemical Ltd., UK and Roche Diagnostic, Germany.

Sourcing for the Tree Bark of *Blighia sapida*: A sizeable quantity of the tree bark of *Blighia sapida* was obtained from the compound of the Federal Polytechnic, Ado Ekiti, Nigeria.

Identification of Plant: The fruits and leaves of *Blighia sapida* plant were obtained from the compound of the The air-dried tree bark of *Blighia sapida* was pulverized.

100g of the pulverized sample was extracted with 800ml of distilled water for seventy-two (72) hours in an extractor. The aqueous extract was obtained by filtering with Whatman filter paper and subsequently freeze-dried in Armfield freeze-drier for ten (10) days.

Qualitative Phytochemical Analysis: Chemical tests were carried out on the aqueous extract using modified standard procedures to identify the constituents as described by Sofowora(1993), Trease and Evans(1989) and Harborne (1973).

Quantitative Determination of Phytochemicals Determination of Tannin: The tannin content of the extract was determined by using the modified procedure of Makkar (1994).

Determination of Saponin: The method used was that described by Obadoni and Ochuko (2001).

Determination of Flavonoid: The method of Boham and Kocipal-Abyazam (1994) was used.

Determination of Alkaloid: The total alkaloid content of the extract was determined using the method described by Harborne (1973).

Determination of Total Phenols: The total phenolic content was determined using the method described by Singleton and Rossi(1965) using Folin-Ciocalteu's phenol reagent.

Determination of Zinc and Selenium: The level of zinc and selenium were determined by the method described by AOAC (2006).

In vitro Antioxidant Assay DPPH free radical scavenging assay: The hydrogen or radical scavenging properties of the extract was determined using the stable radical DPPH (2, 2-Diphenyl-1-picrylhydrazyl hydrate) according to the method proposed and described by Blois (1958).

Hydroxyl radical scavenging assay: Deoxyribose assay was used to determine the hydroxyl radical scavenging activity in an aqueous medium .

Hydrogen peroxide decomposition assay: This activity was determined according to a method described by Long et al., (1999).

Nitric oxide (NO) scavenging assay: At physiological pH, nitric oxide generated from aqueous sodium nitroprusside (SNP) solution interacts with oxygen to produce nitrite ions, which may be quantified by the Griess Illosvoy reaction (Garrratt, 1964).

Ferric reducing antioxidant assay (FRAP): The FRAP assay uses antioxidants as reductants in a redox-linked colorimetric method with absorbance measured with a spectrophotometer (Benzie and Strain, 1999). The principle of this method was based on the reduction of a colourless ferric-tripyridyltriazine complex to its blue ferrous colour formed owing to the action of electron donation in the presence of antioxidants.

Determination of total antioxidant capacity: The total antioxidant capacity was determined in accordance with the method described by Prieto et al., (1999). Statistical Analysis: Data were expressed as mean + S.E.M. of five replicates.

3 III. Results

Phytochemical screening of extract: The result of phytochemical screening of aqueous extract of *Blighia sapida* stem bark shows the presence of tannin, saponin, flavonoid, alkaloid, phenols and ascorbic acid.

It also shows the presence of the trace elements zinc and selenium (Table 1 and 2).

In vitro antioxidant activity: In vitro antioxidant studies revealed that aqueous extract of *Blighia sapida* stem bark showed some scavenging activity by total antioxidant capacity (TAC) with the IC 50 value of $16.63 \pm 0.51 \mu\text{g/ml}$, DPPH radical scavenging activity with the IC 50 value of $23,938 \pm 169.28 \mu\text{g/ml}$, total phenol of $5.47 \pm 0.93 \mu\text{g/ml}$ GAE, hydrogen peroxide radical decomposition activity with the IC 50 value of $5,133.50 \pm 191.70 \mu\text{g/ml}$, hydroxyl radical scavenging activity with IC 50 value of $6904.00 \pm 751.90 \mu\text{g/ml}$ and nitric oxide scavenging activity with IC 50 value of $447.57 \pm 153.28 \mu\text{g/ml}$. However, apart from the nitric oxide scavenging activity value that compares with that of the standard (i.e. ascorbic acid), all other values are a lot higher than those of the standards used (Table 3).

4 IV. Discussion

Antioxidants such as phenolic compounds (tocopherols flavonoids, and phenolic acids), nitrogen compounds (alkaloids, chlorophyll derivatives, amino acids and amines), carotenoids and ascorbic acid (Hall and Cuppett, 1997;Larson, 1988) compounds inhibit or delay the oxidation of other molecules by inhibiting the initiation or propagation of oxidizing chain reactions.

In this study, the metabolites shown in Table ?? were known to show biological activity as well as exhibiting physiological activity (Sofowora, 1993). Flavonoids are potent water-soluble antioxidants and free radical scavengers which prevent oxidative cell damage and have strong anticancer activity (Okwuet al., 2006).

Flavonoids also lower the risk of heart disease. Saponins are capable of neutralizing some enzymes in the intestine that can become harmful, building the immune system and promoting wound healing. Alkaloids have been documented to possess analgesic, antispasmodic and bactericidal effects. Tannins hasten the healing of wounds and inflamed mucuous membrane (Okwu et al., 2006). The presence of these phytochemicals support the medicinal use of Blighia sapida (Saidu, 2012).

Zinc and selenium are the trace elements that have been found to be cofactors of antioxidant enzymes such as glutathione peroxidase and superoxide dismutase (Weydert and Cullen, 2009).

In vitro antioxidant studies revealed that aqueous extract of Blighia sapida stem bark only showed a comparative scavenging activity when compared to the standard (i.e. ascorbic acid), by nitric oxide scavenging activity. All other parameters considered in the in vitro antioxidant studies did not show promising results when compared with the standards.

5 V. Conclusion

A major finding of the study is that aqueous extract of Blighia sapida stem bark showed the presence of metabolites such as tannin, saponin, flavonoid, alkaloid, phenols and ascorbic acid. It was also found to contain trace elements zinc and selenium.

The results also revealed that Blighia sapida stem bark aqueous extract showed some in vitro antioxidant activity. Ofall the parameters studied, nitric oxide scavenging activity was the only one that showed a somewhat comparable IC 50 to that of the standard while all others did not compare well with the chosen standards. ¹

2

Species	Extract Type	Tannin	Saponin	Flavonoid	Alkaloid	Phenols	Ascorbic Add(mg/100g)	Zinc (ppm)	Selenium (ppm)
Blighia sapida	Aqueous	22.0±1.24	21.3±3.30	13.3±1.86	17.3±4.70	5.47±0.93	22.1±6.02	19.6±0.03	

[Note: © 2017 Global Journals Inc. (US)]

Figure 1: Table 2 :

3

Figure 2: Table 3 :

EXTRACT TYPE	Total Antioxi- dant Capacity (TAC)	Ferric reducing antioxidant (FRAP)	DPPH free radi- cal Scavenging IC 50 Values (µg/ml)	Total Phenol (GAE)	Hydrogen Peroxide	Hydroxyl radical	M o i
Aqueous	16.63 ± 0.81	9.13 ± 0.51	23938.00 ± 160.28	5.47 ± 0.93	5133.50 ±	6904.00 ±	4
Year 2017	Ascorbic acid Butylated	—	Standards ± 5.02	69.19 —	191.70	123.41 ± 4.84	-139.
	Hydroxytoluene (BHT)						
D							
D							
D							
D)							
B							
(							

[Note: signaling pathways mediators of insulin resistance and ?? ?cell dysfunction?, Diabetes, 52, 1-8. Volume XVII Issue 1 Version I]

Figure 3:

-
- [Am. J. Enol. Vitic] , *Am. J. Enol. Vitic* 16 p. .
- [Harbone ()] , J B Harbone . *Phytochemical Methods* 1973. Chapman and Hall Ltd. p. .
- [Halliwell et al. ()] , B Halliwell , M Grooveld , J M Gutteridge . *Methods of Biochemical Sciences* 1981. 33 p. .
- [Halliwell et al. ()] , B Halliwell , M Grooveld , J M Gutteridge . *Methods of Biochemical Sciences* 1981. 33 p. .
- [Holme and Peck ()] , D Holme , H Peck . *Analytical Biochemistry* 1998. Addison Wesley Longman Limited. (3rd Edition)
- [Sofowora] 1993): *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Book Ltd, E A Sofowora . Ibadan, Nigeria. p. 289.
- [Hall and Cuppet ()] 'Activities of Natural antioxidants'. C A Hall , S L Cuppet . *Antioxidant Methodology in vivo and in vitro Concepts*, O I Aruoma, S L Cuppet (ed.) (Champaign, II) 1997. AOCS Press. p. .
- [Blois ()] 'Antioxidant determination by the use of a stable free radical'. M S Blois . *Nature* 1958. 181 p. .
- [Boham and Kocipal-Abyazam ()] B A Boham , R Kocipal-Abyazam . *Flavonoids and Condensed Tannins from Leaves of*, 1994.
- [Teitze ()] 'Enzymatic method for the quantitative determination of nanogram amounts of total and oxidized glutathione: Applications to mammalian blood and other tissues'. F Teitze . *Anal. Biochem* 1969. 27 p. .
- [Gardiner et al. ()] 'Extracts from *Blighia sapida* (Koenig) produce neutropenia and thrombocytopenia in mice'. M T Gardiner , L A D William , T L The , C K Fletcher , P D A Singh , G Wharfe , E Choo-Kang , R N Sawh , E Rickards . *Phytother Res* 1996. 10 p. .
- [Benzie and Strain ()] 'Ferric Reducing/ Antioxidant Power Assay: Direct Measure of Total antioxidant Activity of Biological Fluids and Modified Version for Simultaneous Measurement of Total Antioxidant Power and Ascorbic Acid Concentration'. F F Benzie , J J Strain . *Methods in Enzymology* 1999. 299 p. .
- [Long et al. ()] 'Hydrogen peroxide in human urine: implications for antioxidant defense and redox regulation'. L H Long , P Evans , B Halliwell . *Biochem. Biophys. Res Commun* 1999. 262 p. .
- [Makkar ()] H P Makkar . *Quantification of tannins. A laboratory manual international for Agricultural Research in the Dry Area (ARDA)*, 1994.
- [Weydert and Cullen ()] 'Measurement of Superoxide dismutase, catalase, and Glutathione peroxidase in cultured cells and tissue'. C J Weydert , J Cullen . *Nat. Protoc* 2009. 5 (1) p. .
- [Aderinola et al. ()] 'Nutritional Potential of *Blighia sapida* K Konig (Ackee akkee) leaves as a dry season feed resources for West Africa dwarf goats in the derived savanna zone of Nigeria'. O A Aderinola , G O Farinu , J A Akinlade , T B Olayemi , O Ojebiyi , P Ogunniyi . *Livestock Res. Rural Dev* 2007. 19 (6) p. 78.
- [OA C ()] *Official methods of Association of Organic and Analytical Chemists*, OA C . 2006.
- [Oladiji et al. ()] 'Physicochemical properties of oil from the fruit of *Blighia sapida* and Toxicological Evaluation of the Oil-Based Diet in Wister Rats'. A T Oladiji , K L Shoremekun , M T Yakubu . *Journal of Medicinal Food* 2009. 12 (5) p. .
- [Saidu et al. ()] 'Phytochemical Screening and Hypoglycemic Effect of Aqueous *Blighia sapida* Root Bark Extract on Normoglycemic Albino Rats'. A N Saidu , A Mann , C D Onuegbu . *British Journal of Pharmaceutic Research* 2012. 156 p. .
- [Obadoni and Ochuko ()] 'Phytochemical studies and comparative efficacy of the crude extracts of some hemostatic plants in Edo and Delta States of Nigeria'. B Obadoni , P Ochuko . *Global J. Pure Appl. Sc* 2001. 8 p. .
- [Khan and Gumbs ()] 'Repellent effect of ackee (*Blighia sapida* Kaonig) component fruit parts against stored product insect pests'. A Khan , F A Gumbs . *Trop. Agric* 2003. 80 p. .
- [Singleton et al. ()] V L Singleton , J A Rossi , Jr . *Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents*, 1965.
- [Prieto et al. ()] 'Spectrophotometric quantitation of antioxidant capacity through the formation of a Phosphomolybdenum Complex: Specific application to the determination of vitamin E'. P Prieto , M Pineda , M Aguilar . *Analytical Biochemistry* 1999. 269 p. .
- [Larson ()] 'The antioxidants of higher plants'. R Larson . *Phytochemistry* 1988. 27 p. .
- [Garraat ()] *The quantitative analysis of drugs*, D C Garraat . 1964. Japan: Chapman and Hall Ltd. 3 p. .
- [Trease and Evans ()] G E Trease , W C Evans . *Pharmacognosy, 3rd Edition*, (London) 2006. p. .
- [Okwu et al. ()] 'Vitamin C improves basal metabolic rate and lipid profile in alloxan-induced diabetes mellitus in rats'. D U Okwu , A B Antai , K H Udofia , A O Obembe , K O Obasi , M U Eteng . *J. Biosci* 2006. 31 (5) p. .