

Physicochemical, Volatile Organic Composition, Phenolic, Flavonoid and Ascorbic Acid Contents, Antioxidant, Anti-Arthritic and Anti-Inflammatory Properties of Cocos nucifera Juice

Ololade Z.S

Received: 14 December 2016 Accepted: 5 January 2017 Published: 15 January 2017

Abstract

Various parts of Cocos nucifera are locally used for treatment of diseases and production of some foods and beverages for man and animals. This study examined the physicochemical properties, phytochemical and multi therapeutic potentials of juice of C. nucifera from Nigeria. These were measured using GC-MS, pH meter, specific gravity, UV-Vis spectrometry, Folin-Ciocalteu's, aluminium chloride, DPPH, PTAC and egg albumin methods respectively. GC-MS analysis revealed the presence of nitroisobutylglycerol as the most abundant volatile organic compounds in the juice. The pH, clarity, turbidity, TPC, TFC and TAA were 5.09, 1.34, 1.07, 2,261.5 $\pm$ 0.00 $\pm$ 0.0 μ gmg⁻¹ GAE, 20.00 $\pm$ 0.0 μ gmg⁻¹ QE and 66.75 $\pm$ 0.00 $\pm$ 0.0 μ gmg⁻¹ AAE, respectively. The antioxidant IC₅₀ and AAI values of the juice were 0.25mgml⁻¹ and 160 and it was capable of scavenging free radicals at a range between 33.09-76.26

Index terms—

1 I. Introduction

Natural products are good drug leads and phytochemical probes to explore mechanisms associated with infection of diseases (Kingston, 2011; Ibekwe and Ameh, 2014). Numerous plant polyphenols reportedly afford multiple health benefits. They are regarded as healthy food and nutrient sources because of their many beneficial components (Voravuthikunchai and Howe, 2014; Zhang et al., 2016). Secondary metabolites in the natural products are considered to be responsible for positive health outcomes (Gechev et al., 2014). Particularly, it is widely noted that plants produce a great deal of antioxidants to combat the oxidative stress induced by oxygen and light in the natural environment (Cartea et al., 2011; Qi et al., 2016). Natural products possess antimicrobial and antioxidant activities responsible for the prevalence of dermatophytosis (Thebo et al., 2016). Many approved therapeutics and drugs are derived from natural sources (Cragg and Newman, 2013; Lahlou, 2013).

Cocos nucifera Linn commonly known as coconut is an important fruit tree in the tropical regions and the fruit can be made into a variety of foods and beverages (Yong et al., 2009). C. nucifera is an important member of the family Arecaceae. The juice of C. nucifera has the ability to prevent diseases and sickness. This is due to the free radical scavenging abilities of the antioxidant phytochemicals in it. C. nucifera has long been used in traditional medicine for different kind of illness and almost all parts have their uses. The juice inside the fruit is sterile but when it is extracted and exposed to air, it becomes subjected to quick oxidation and microbial contamination leading to depletion of nutrients and spoilage (Matsui et al., 2008; ueiroz et al., 2008; ean et al., 2009; akono et al., 2012; dubofuor et al., 2016). Juice of C. nucifera is one of the natural food products to quench thirst and easily available in most of the countries. Both water and meat of coconut refresh the body by providing nutritious content. Traditionally, it has been used to protect the body against infection by dangerous diseases. It has been found to improve digestion and hasten the absorption of nutrients including vitamins, minerals, and amino acids. Recently, the health and medicinal uses of C. nucifera products get research interest because it contains several metabolites such as sugars, proteins, free amino acids, vitamins, minerals and

growth promoting factors (Reddy and Lakshmi, 2014). The juice of *C. nucifera* contains many enzymes including acid phosphatase, catalase, dehydrogenase, diastase, peroxidase and RNA polymerase. Juice of *C. nucifera* is locally consumed fresh, directly from the fruit. (Adubofuor et al., 2016). This study aimed at evaluations of physiochemical, phytochemical and therapeutic efficacies of *C. nucifera* juice.

II. Materials and Methods

The fruit of *C. nucifera* was gotten from Ota, Ogun State, Nigeria and the juice was collected and then stored in vial at 5 °C temperature to prevent contamination.

N

Abstract-Various parts of *Cocos nucifera* are locally used for treatment of diseases and production of some foods and beverages for man and animals. This study examined the physicochemical properties, phytochemical and multi therapeutic potentials of juice of *C. nucifera* from Nigeria. These were measured using GC-MS, pH meter, specific gravity, UV-Vis spectrometry, Folin-Ciocalteu's, aluminium chloride, DPPH, PTAC and egg albumin methods respectively. GC-MS analysis revealed the presence of nitroisobutylglycerol as the most abundant volatile organic compounds in the juice. The pH, clarity, turbidity, TPC, TFC and TAA were 5.09, 1.34, 1.07, 2,261.5±0.00 µgmg⁻¹ GAE, 20.00±0.0 µgmg⁻¹ QE and 66.75±0.00 µgmg⁻¹ AAE, respectively. The antioxidant IC₅₀ and AAI values of the juice were 0.25 mgml⁻¹ and 160 and it was capable of scavenging free radicals at a range between 33.09-76.26%. The TAC was 645.38±0.00 µgmg⁻¹ AAE. The protein denaturation inhibition capacity was at the range between 42.0-73.4%. Therefore, the juice of *C. nucifera* could be used as multi therapeutic agent.

Volume XVII Issue II Version I

b) Colour Determination

Colour of the juice was determined by physical observation in day light (Barkatullah et al., 2012).

c) Odour Determination

Odour of the juice was determined by organoleptic evaluation (Aloko et al., 2017).

d) Determination of pH

The pH of the juice of *C. nucifera* was determined immediately after extraction at room temperature using digital pH meter (Paz et al., 2016).

e) Determination of Specific Gravity (SG)

A clean specific gravity bottle was weighted (W_0). Then the bottle was filled to the brim with water and stopper was inserted. The water on the stopper and bottle were carefully wiped off and reweighed (W_1). Same process was repeated, but using juice samples instead of water and weighted again (W_2). The specific gravity of the juice was calculated using the formula below.

Specific gravity

$$= (W_2 - W_0) / (W_1 - W_0)$$

Where: W_0 = Weight of empty specific gravity bottle W_1 = Weight of water + specific gravity bottle W_2 = Weight of test sample + specific gravity bottle.

f) GC-MS Analysis

The juice of *C. nucifera* dissolved in methanol was analysed using Shimadzu GC-MS-QP2010 Plus (Japan). The separations were carried out using a Restek Rtx-5MS fused silica capillary column (5% diphenyl-95% dimethylpolysiloxane) of 30 m × 0.25 mm internal diameter (di) and 0.25 mm in film thickness. The conditions for analysis were set as follows; column oven temperature was programmed from 60-280 °C (temperature at 60 °C was held for 1.0 min, raised to 180 °C for 3 min and then finally to 280 °C held for 2 min); injection mode, Split ratio 41.6; injection temperature, 250 °C; flow control mode, linear velocity (36.2 cm/sec); purge flow 3.0 ml/min; pressure, 56.2 kPa; helium was the carrier gas with total flow rate 45.0 ml/min; column flow rate, 0.99 ml/min; ion source temperature, 200 °C; interface temperature, 250 °C; solvent cut time, 3.0 min; start time 3.5 min; end time, 24.0 min; start m/z, 50 and end m/z, 700. Detector was operated in EI ionization mode of 70 eV. Components were identified by matching their mass spectra with those of the spectrometer data base using the NIST computer data bank, as well as by comparison of the fragmentation

11 g) Determination of Total Phenolic Content (TPC)

The TPC of the juice of *C. nucifera* was determined using Folin-Ciocalteu method. 1 ml of juice was mixed with 1 ml of 10% Folin-Ciocalteu reagent in distilled water and then neutralized with 4 ml of 7.5% sodium carbonate solution. The sample was maintained at room temperature for 3 hrs with periodical mixing, the absorbance at 760 nm was measured using UV-vis spectrophotometer. Gallic acid was used as a reference and for the calibration curve; result was expressed in micrograms per gram of gallic acid equivalent (Vasudevarao and Sravanthi, 2017).

12 h) Total Flavonoid Concentration (TFC)

The total flavonoid content of the juice of *C. nucifera* was determined by spectrophotometry, using aluminium chloride method. Briefly, 1.0 ml of the juice, 0.10 ml of 10% aluminium chloride, 0.10 ml of sodium acetate (1 M) and 2.80 ml of distilled water. After incubation for 40 min, absorbance was measured at 415 nm using a UV-Vis-spectrophotometer. To calculate the concentration of flavonoids, we prepared a calibration curve using quercetin as standard. The index of TFC concentration is expressed as quercetin equivalents (QE) in µg per mg of juice. All assays were carried out in triplicate .

13 i) Determination of Total Ascorbic acid content (TAA)

1 ml of the juice was added to 1.0 ml 2,4-dinitrophenylhydrazine (2,4-DNPH). It was allowed to stand for 30 min. and the absorbance was read in triplicate at 515 nm, using distilled water as blank. Ascorbic acid was used as a reference and for the calibration curve; result was expressed in milligram per gram of ascorbic acid equivalent (Benites et al., 2015).

14 j) Determination of Free Radical Scavenging and Antioxidant Activities i. In vitro 2,2'-Diphenyl-1-picryl-hydrazyl Assay

The antioxidant and free radical scavenging of the juice of *C. nucifera* was measured by using DPPH. Briefly, the reaction mixture of 2.0 ml; consist of 1.0 ml of DPPH in methanol (0.004%) and 1.0 ml of various concentrations of juice. Then incubated for 30 min. in dark, and the absorbance was measured at 517 nm. The control was prepared by DPPH and methanol in place of sample. In this assay, the positive control is ascorbic acid. The percentage of inhibition can be calculated using the formula: $I\% = [(A \text{ blank} - A \text{ juc}) / A \text{ blank}] \times 100$ pattern with those reported in the literature (Ololade et al., 2017).

Where: A blank is the absorbance of blank solution and A juc is the absorbance of the juice. The dose response curve was plotted and IC 50 value for the juice and the standard were calculated (Ololade et al., 2016). AAI was classified as weak, when AAI < 0.5, moderate, when AAI ranged between 0.5-1.0, strong, when AAI ranged between 1.0-2.0, and very strong, when AAI > 2.0 (Arulpriya and Lalitha, 2014).

15 ii. Phosphomolybdate Total Antioxidant Capacity (PTAC) Assay

The PTAC of the juice of *C. nucifera* was determined with phosphomolybdenum using ascorbic acid as the standard. An aliquot of 1.0 ml of juice solution is combined with 1.0 ml of reagent (0.6 M sulphuric acid, 28 µM sodium phosphate and 4 µM ammonium molybdate). The tubes were capped and incubated in a boiling water bath at 95 °C for 90 min. after the samples had cooled to room temperature, the absorbance of the aqueous solution of each were measured at 695 nm in UV spectrophotometer. The blank solution contained 1.0 ml of reagent solution and the appropriate volume of the same solvent used for the sample and it was incubated under same conditions as the rest of the samples. The total antioxidant capacity was expressed as equivalents of ascorbic acid (Bulus et al., 2017).

16 In-vitro Anti-Arthritic and Anti-Inflammatory Activities of the Juice on Inhibition of Protein Denaturation (Egg Albumin

Assay): in vitro anti-arthritic/anti-inflammatory activity of the juice was evaluated against protein denaturation method using fresh hen's egg albumin. About 5 ml reaction mixtures (0.2 ml of egg albumin, 2.8 ml of phosphate buffered saline (PBS, pH 6.4) add 2 ml of test sample at 1000, 500, 250 and 125 µgml⁻¹). Distilled water with same volume (0.2 ml) was used as control. The mixtures were incubated at 37 °C in BOD incubator for about 15 min. followed by heating at 70 °C for 5 min. After cooling to the room temperature, absorbance was measured spectrophotometrically at 660 nm using vehicle as blank. Aspirin (3000 µgml⁻¹) was used as reference drug. The inhibition percentage of protein denaturation was calculated using the following formula: % inhibition = $100 \times (V_t/V_c) - 1$

Where:

V t = absorbance of test sample, V c = absorbance of control.

The drug concentration for 50% inhibition (IC 50) was determined by plotting percentage inhibition with respect to control against treatment concentration. (Smitha et al., 2017).

17 III. Results and Discussion

18 a) Physicochemical Properties of the Juice of *C. nucifera*

Determination of different physicochemical properties showed the practical importance and provides bases for suitability, consumption, utility, nutritional and physical qualities of the natural juice of *C. nucifera* in daily life (Angaye and Maduelosi, 2015). Physicochemical properties of the juice such as colour, odour, pH, clarity, turbidity, specific gravity (Table 1) showed the quality of the juice of *C. nucifera* from Nigeria.

19 b) Colour and Odour of the Juice of *C. nucifera*

The colour of the fresh juice of *C. nucifera* was milky in nature with sweet aromatic odour.

20 c) Clarity and Turbidity of the Juice of *C. nucifera*

The clarity and turbidity of the juice of *C. nucifera* were determined as 0.74 and 0.66 respectively using UV-Vis spectrophotometer.

21 d) pH of the Juice of *C. nucifera*

The pH value of the *C. nucifera* juice was 5.09; which was within the standard limit (pH 3.40-6.10) that insures freshness of the juice (El-Sohaimy et al., 2015). The acidity of juice might be due of present organic acid in the juice, which is responsible for important characteristics of juice: flavour and stability against microbial spoilage and this may confer longer keeping quality of the juice (Nadzirah et al., 2012; Offia-Olua and Ekwunife, 2015). Furthermore, it might also indicate that the juice of *C. nucifera* have high content of minerals. pH is a very important parameter in the conduct of fermentation. A pH of 4 is the optimum for the growth of fermentative yeast. This also inhibits the development of undesirable microbial flora (Ahoussi et al., 2015; Walker and Stewart, 2016).

22 e) Specific Gravity of the Juice of *C. nucifera*

The specific gravity of the juice of *C. nucifera* was 1.01 (Table 1). This was in line with the amount stated for beverages (including soft drinks and juices) and fruit drinks (low calories and undiluted) as 1.01-1.03. The more sugar present in a juice, the denser the juice becomes. Juice is low dense foods because of its high water content, which provides high volume and weight. To stay within low density guidelines, it is important to either consume natural fruit juice that has not been dehydrated than to eat processed fruit juice that contains added sugar ??Swinburn et al.,

23 g) Total Phenolic Content, Total Flavonoid Contents and Total Ascorbic Acid

The TPC, TFC and TAA analyses of the investigated juice of *C. nucifera* showed the presence of high amount phenolic, flavonoid compounds and ascorbic acid (Table 2). Natural phenolic compounds and ascorbic acid play many significant roles in human health as evident from their therapeutic properties ??Dimitros, 2006; ??nsari et al., 2013; ??zialo et al., 2016). Plants consumed by humans may contain thousands of different amounts of ascorbic acid, phenolic and flavonoid components (Saxena et al., 2013; Kasote et al., 2015; Zhang et al., 2015). The effect of dietary ascorbic acid and phenolics is currently of great interest due to their antioxidant and possible anticarcinogenic activities (Nahak et al., 2014; Pereira et al., 2009). Ascorbic acid, Phenolic and flavonoid compounds are chain breaking antioxidant, free radicals scavenger and quenchers of singlet oxygen formation in the process of formation of intracellular substances throughout the body (Mitra and Uddin, 2014; Ozcan et al., 2014). Moreover, ascorbic acid, phenolic and flavonoid components play important roles in the control of cancer and other human diseases (Ghasemzadeh and Ghasemzadeh, 2011). The percentage inhibitions of the juice at various concentrations (1000, 750, 500 and 250 µgml⁻¹) were 76.26, 71.22, 71.00 and 33.09% respectively; while the IC 50 value was found to be 0.25 mgml⁻¹ in comparison to ascorbic acid with IC 50 value of 9.0 ?gml⁻¹.

24 i) Phosphomolybdate Total Antioxidant Capacity (PTAC)

The PTAC of the juice of *C. nucifera* was 645.38±0.00 µgmg⁻¹ AAE (Table 3). The measure of the ability of natural products to delay oxidative stress in a controlled system is defined as total antioxidant capacity ??Apak et al., 2016; ??hang et al., 2016; ??ieme et al., 2017; Tyagi and Agarwal, 2017). The juice showed high antioxidant potential and this can be related to the high amounts of ascorbic acid, flavonoids and phenolic compounds in juice. Antioxidant play definite roles in many pathological conditions and they are known to fight against these free radicals and protects body from various diseases ??Aprioku et al., 2013; ??one et al., 2013). Their mechanism

of action is either by scavenging the reactive oxygen species or protecting the antioxidant defence mechanisms (Birben et al., 2012). The total antioxidant potential is a relevant tool for investigating the relationship between dietary antioxidants and pathologies induced by the oxidative stress (Pisoschi and Negulescu, 2011).

25 j) In-vitro Anti-Arthritic and Anti-Inflammatory Potential

The juice of *C. nucifera* possesses potentially useful anti-arthritic and anti-inflammatory activities at the doses tested. The juice exhibited significant inhibition of egg albumins denaturation of protein between 42.0-73.4% at concentrations between 125-1000 µgml⁻¹ and with the IC 50 value of 6.0 µgml⁻¹. The result was comparable to that of standard drug aspirin (89.4%) (Table 4). The juice showed inhibition of heat-induced protein (albumin) denaturation and prominent effects on protein denaturation was produced. Natural products that can prevent protein denaturation would be very useful for the development of anti-arthritic and antiinflammatory drugs (Sowjanya et al., 2013;Janakiraman and Parameswari, 2014;Obaseki et al., 2016). Therefore, the juice is a promising anti-arthritic agent of natural origin in the treatment of inflammatory disorders. It shows that the juice is capable of reducing the production of auto-antigen which indirectly reduces the protein denaturation and hence alleviate arthritis (Alamgeer et al., 2017;Boddupally et al., 2017). Protein denaturation is one of the leading causes of inflammatory as well as arthritic diseases, which led to production of auto antigens, progressing to certain rheumatic diseases (Jayaprakasam and Ravi, 2012 Kaliwal, 2017). The main mechanism involved in protein denaturation is characterized by changes or alterations in hydrophobic, electrostatic, hydrogen and disulphide bonding among the protein molecules (Zavodszky et al., 2001;Sangeetha and Vidhya, 2016).

26 IV. Conclusion

Conflict of Interest Statement: The authors declared that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The results of this study showed that the juice of *C. nucifera* can be an accessible source of promising therapeutic agents that can be used in Combating some infectious diseases caused. The study showed the presence of significant antioxidant, anti-arthritic antiinflammatory activities of the juice. The activities were due to the presence of pharmacologically active phytochemicals in the juice. Fresh juice of *C. nucifera* is a fluid that could be consumed for health, refreshment and nutritional purposes. Therefore the juice can include the production and commercialization of foods and drugs.

1

D D D D) B
(

Figure 1: Table 1 :

2

TPC	TFC	TAA
2,261.5±0.00	20.0±0.00	66.75±0.00
µgmg ⁻¹ GAE	µgmg ⁻¹ QE	µgmg ⁻¹ AAE

Data are presented as the mean value ± S.D. of triplicate

h) Free Radical Scavenging and Antioxidant Potentials

Figure 2: Table 2 :

3

Juice and Reference Drug	DPPH IC 50 mgml ⁻¹	AAI	PTAC µgmg ⁻¹ AAE
Juice	0.25	160	645.38±0.00

Figure 3: Table 3 :

4

Conc. μgml^{-1}	% Inhibition	IC 50 μgml^{-1}	% Inhibition of Aspirin 3000 μgml^{-1}
1000	73.4		
500	71.4	6.0	89.4
250	48.0		
125	42.0		

Figure 4: Table 4 :

- [Sulfone] , Dimethyl Sulfone . *IOSR Journal of Pharmacy and Biological Sciences* 12 (1) p. .
- [Pereira et al. ()] , D M Pereira , P Valentao , J A Pereira , P B Andrade . *Phenolics: From Chemistry to Biology* 2009. 14 p. . (Molecules)
- [Saxena et al. ()] , M Saxena , J Saxena , R Nema , D Singh , A Gupta . *Journal of Pharmacognosy and Phytochemistry* 2013. 1 (6) p. . (Phytochemistry of Medicinal Plants)
- [Sowjanya et al. ()] , K R R Sowjanya , R V Vishweswar , F Bushra , M Kapila , M Srilakshmi . *In-vitro Anti Inflammatory Activity of Ultra Sonic Bath Assisted, Methanol Extract of Lepidium Sativum Linn. Seeds, International Journal of Pharmaceutical Development and Technology* 2013. 3 (2) p. .
- [Pashikanti et al. ()] ‘Anti-arthritis activity of ethanolic extract from the leaves of Commiphora caudata (Linn.) in complete Freund’s adjuvant-induced arthritic rats’. G Pashikanti , U Kulundaivelu , V R Jupalli , S B Sheshagiri , V Eggadi . *Nigerian Journal of Experimental and Clinical Biosciences* 2014. 2 (1) p. .
- [Arulpriya and Lalitha ()] ‘Anti-inflammatory and antioxidant activity index of the aerial roots of Rhabdophora aurea (Linden Ex Andre) intertwined over different host trees’. P Arulpriya , P Lalitha . *Journal of Pharmacy Research* 2014. 8 (7) p. .
- [Zhang et al. ()] ‘Antioxidant Phytochemicals for the Prevention and Treatment of Chronic Diseases’. Y J Zhang , R Y Gan , S Li , Y Zhou , A N Li , D P Xu , H B Li . *Molecules* 2015. 20 p. .
- [Nahak et al. ()] ‘Antioxidant Potential and Nutritional Values of Vegetables: A Review’. G Nahak , M Suar , R K Sahu . *Research Journal of Medicinal Plants* 2014. 8 p. .
- [Tyagi and Agarwal ()] ‘Antioxidant Properties and Phenolic Compounds in Methanolic Extracts of Eichhornia crassipes’. T Tyagi , M Agarwal . *Research Journal of Phytochemistry* 2017. 11 p. .
- [Alamgeer et al. ()] ‘Appraisal of anti-arthritis and nephroprotective potential of Cuscuta reflexa’. S G N Alamgeer , M U Ambreen , N Q Muhammad , A Haseeb . *Pharmaceutical Biology* 2017. 55 (1) p. .
- [Ololade and Olawore ()] ‘Characterization of Essential Oil from the Seed of Eucalyptus cloeziana and Evaluation of its Modes of Medicinal Potentials’. Z S Ololade , N O Olawore . *Edorium Journal of Infectious Diseases* 2017. 3 p. .
- [Angaye and Maduelosi ()] ‘Comparative study of the physicochemical properties of some Refined vegetable oils sold in mile one market and some departmental stores in port Harcourt, Rivers State’. S Angaye , N Maduelosi . *Nigeria. Discourse Journal Agriculture and Food Sciences* 2015. 3 p. .
- [Benites et al. ()] ‘Contents of constituents and antioxidant activity of seed and pulp extracts of Annona coriacea and Annona sylvatica’. R S R Benites , A S N Formagio , E J S Argandoña , C R F Volobuff , L N F Trevizan , M C Vieira , M S Silva . *Braz. J. Biol* 2015. 75 (3) p. .
- [Rane and Anusha ()] ‘Determination of Bioactive components from the Ethanolic Peel extract of citrus reticulata by Gas chromatography-Mass Spectrometry’. Z A K P Rane , B Anusha . *International Journal of Drug Development and Research* 2012. 4 (4) p. .
- [Swinburn et al. ()] ‘Diet, nutrition and the prevention of excess weight gain and obesity’. B A Swinburn , I Caterson , J C Seidell , W P T James . *Public Health Nutrition* 2004. 7 (1A) p. .
- [Ledikwe et al. ()] ‘Dietary energy density is associated with energy intake and weight status in US adults’. J H Ledikwe , H M Blanck , L K Khan , M K Serdula , J D Seymour , B C Tohill , B J Rolls . *The American Journal of Clinical Nutrition* 2006. 83 (6) p. .
- [Zavodszky et al. ()] ‘Disulfide bond effects on protein stability: Designed variants of Cucurbita maxima trypsin inhibitor-V’. M Zavodszky , C W Chen , J K Huang , M Zolkiewski , L Wen , R Krishnamoorthi . *Protein Science* 2001. 10 p. .
- [Paz et al. ()] ‘Effect of pH at drainage on the physicochemical, textural and microstructural characteristics of mozzarella cheese from goat milk’. N F Paz , E G Oliveira , F J Villalva , M Armada , Ramon , AN . *Food Science and Technology* 2016. 37 (2) p. .
- [Jayaprakasam and Ravi ()] ‘Evaluation of Anti-Arthritis Activity of the Root Extract of Acalypha Indica Linn. Using In vitro Techniques’. Jayaprakasam , T K Ravi . *International Journal of Phytopharmacy* 2012. 2 (6) p. .
- [Janakiraman and Parameswari ()] ‘Evaluation of Anti-inflammatory effect of Plectranthus amboinicus leaf extract -An in vitro study’. D Janakiraman , C S Parameswari . *Journal of Advanced Pharmacy Education and Research* 2014. 4 (2) p. .
- [Obaseki et al. ()] ‘Evaluation of the antiinflammatory properties of the hexane extract of Hydrocotyle bonariensis Comm’. O E Obaseki , O I Adesegun , G N Anyasor , O O Abebawo . *Ex Lam. Leaves* 2016. 15 (49) p. .
- [Ololade et al. ()] ‘Free Radical Scavenging, Antioxidant and Antibacterial Activities of the Fruit-Pulp Essential Oil of Annona muricata and Its Phytochemical Composition’. Z S Ololade , O A Fakankun , F O Alao , O O Ajewole . *International Journal of Applied Research and Technology* 2016. 5 (2) p. .

- [Vasudevarao and Sravanthi ()] *GC/MS analysis and In-vitro Antioxidant activity of methanol extract of Ulothrix flacca and its main constituent*, B Vasudevarao , D J Sravanthi . 2017.
- [Slavin and Lloyd ()] 'Health Benefits of Fruits and Vegetables'. J L Slavin , B Lloyd . *Advances in Nutrition* 2012. 3 p. .
- [Mahabal and Kaliwal ()] 'In vitro Anti-Inflammatory Activity of L-Asparaginase from Soil Rhizosphere Fungus *Aspergillus tamarii*'. N S Mahabal , B B Kaliwal . *Current Trends in Biomedical Engineering and Biosciences* 2017. 4 (5) p. .
- [Smitha et al. ()] 'In vitro Anti-Inflammatory and Anti-Arthritic Activity in Methanolic Peel Extracts of *Persea americana*'. G S R Smitha , B C Jyoti , K K Hemanth . *World Journal of Pharmaceutical and Life Sciences* 2017. 3 (4) p. .
- [Sangeetha and Vidhya ()] 'In vitro antiinflammatory activity of different parts of *Pedaliium murex*'. G Sangeetha , R Vidhya . *International Journal of Herbal Medicine* 2016. 4 (3) p. .
- [Bulus et al. ()] 'In vitro Antioxidant Activity of N-Butanol Extract of *Curcuma longa* and Its Potential to Protect Erythrocytes Membrane against Osmotic-Induced Haemolysis'. T Bulus , S I David , L S Bilbis , A Babando . *Science World Journal* 2017. 12 (1) p. .
- [Boddupally et al. ()] 'In-vitro Anti-Arthritic Effect of the Leaves of *Cassia Fistula* Linn'. M Boddupally , S S Rani , A T Thirupathi , K N V Rao , R Vasanthi . *European Journal of Pharmaceutical and Medical Research* 2017. 4 (2) p. .
- [Birben et al. ()] 'Oxidative Stress and Antioxidant Defense'. E Birben , U M Sahiner , C Sackesen , S Erzurum , O Omer Kalayci . *World Allergy Organization Journal* 2012. p. .
- [Formagio et al. ()] 'Phenolic compounds of *Hibiscus sabdariffa* and influence of organic residues on its antioxidant and antitumoral properties'. A S N Formagio , D D Ramos , M C Vieira , S R Ramalho , M M Silva , N A H Zarate , M A Foglio , J E Carvalho . *Brazilian Journal of Biology* 2015. 75 (1) p. .
- [Ozcan et al. ()] 'Phenolics in Human Health'. T Ozcan , A Akpinar-Bayazit , L Yilmaz-Ersan , B Delikanli . *International Journal of Chemical Engineering and Applications* 2014. 5 (5) p. .
- [Nadzirah et al. ()] 'Physico-Chemical Properties of Pineapple Crown Extract Variety N36 and Bromelain Activity in Different Forms'. K Z Nadzirah , S Zainal , A Noriham , I Normah , A M Siti Roha . *APCBEE Procedia* 2012. 4 p. .
- [El-Sohaimy et al. ()] 'Physicochemical characteristics of honey from different origins'. S A El-Sohaimy , S H D Masry , M G Shehata . *Annal of Agricultural Sciences* 2015. 60 (2) p. .
- [Barkatullah et al. ()] 'Physicochemical Characterization of Essential and Fixed Oils of *Skimmia laureola* and *Zanthoxylum armatum*'. I M Barkatullah , A Rauf , Inyat-Ur-Rahman . *Middle-East Journal of Medicinal Plants Research* 2012. 1 (3) p. .
- [Aloko et al. ()] 'Physicochemical properties and lubricant potentials of *Blighia sapida* Sapindaceae seed oil in solid dosage formulations'. S Aloko , C P Azubuikwe , H A B Coker . *Tropical Journal of Pharmaceutical Research* 2017. 16 (2) p. .
- [Babajide et al. ()] 'Physicochemical Properties and Phytochemical Components of Spiced Cucumber-Pineapple Fruit Drink'. J M Babajide , A A Olaluwoye , T A Taofik Shittu , M A Adebisi . *Nigerian Food Journal* 2013. 31 (1) p. .
- [Offia-Oluan and Ekwunife ()] 'Production and evaluation of the physico-chemical and sensory qualities of mixed fruit leather and cakes produced from apple (*Musa pumila*), banana (*Musa sapientum*), pineapple (*Ananas comosus*)'. B I Offia-Oluan , O A Ekwunife . *Nigerian Food Journal* 2015. 33 p. .
- [Kasote et al. ()] 'Significance of Antioxidant Potential of Plants and its Relevance to Therapeutic Applications'. D M Kasote , S S Katyare , M V Hegde , H Bae . *International Journal of Biological Sciences* 2015. 11 (8) p. .
- [Elisha et al. ()] 'The anti-arthritis, antiinflammatory, antioxidant activity and relationships with total phenolics and total flavonoids of nine South African plants used traditionally to treat arthritis'. I L Elisha , J P Dzoyem , L J McGaw , F S Botha , J N Eloff . *BMC Complementary and Alternative Medicine* 2016. 16 p. .
- [Mitra and Uddin ()] 'Total Phenolics, Flavonoids, Proanthocyanidins , Ascorbic Acid Contents and In-vitro Antioxidant Activities of Newly Developed Isolated Soya Protein'. K Mitra , N Uddin . *Discourse Journal of Agriculture and Food* 2014. 2 (5) p. .
- [Ahoussi et al. ()] 'Valorization of pineapple produced in Benin: Production and evaluation of wine quality parameters'. L E Ahoussi , P E S Ahouandjinou , C T R Konfo , B B Yehouenou , M S Gnansounou , S N S Cakpo , Z Satchivi , E Ahoussi-Dahouenon . *International Journal of Advanced Research* 2015. 3 (11) p. .
- [Walker and Stewart ()] G M Walker , G G Stewart . *Saccharomyces cerevisiae in the Production of Fermented Beverages, Beverages*, 2016. 2 p. .