



Lung Cancer Screening: Beliefs and Recommendations of Primary Care Physicians at the National Guard Hospital (NGHA)

By Roaa R Amer, Adel F Yasky & Alia H Zawawi

King Saud Bin Abdulaziz University for Health Sciences

Abstract- *Background:* Early detection of lung cancers via screening may aid in decreasing the associated mortality; however, optimal screening methods have not yet been established.

Objectives: We aimed to explore the beliefs and attitudes of primary care physicians (PCPs) towards lung cancer screening guidelines in asymptomatic patients.

Methods: We conducted a cross-sectional descriptive study at the NGHA primary care centres, using a validated questionnaire, with 11 questions, developed by the National Cancer Institute, USA, and customized to our medical settings.

Results: 37% of family physicians requested chest X-rays; 1.9%, sputum cytology; and 3.7%, low-dose spiral chest computed tomography (CT) scans. Of the internists, 42.1% requested low-dose CT scans; 62.5%, chest X-rays; and 5%, sputum cytology. The mean knowledge scores for family physicians and internists regarding screening guidelines were 2.3 and 1.5, respectively, and those for their belief in the effectiveness of screening modalities were 7.6 and 6.6, respectively.

Keywords: *lung cancer, physicians, belief and recommendations of PCPs, screening guidelines, family medicine, internal medicine, asymptomatic.*

GJMR-F Classification: NLMC Code: QZ 206



Strictly as per the compliance and regulations of:



Lung Cancer Screening: Beliefs and Recommendations of Primary Care Physicians at the National Guard Hospital (NGHA)

Roaa R Amer ^a, Adel F Yasky ^a & Alia H Zawawi ^b

Abstract Background: Early detection of lung cancers via screening may aid in decreasing the associated mortality; however, optimal screening methods have not yet been established.

Objectives: We aimed to explore the beliefs and attitudes of primary care physicians (PCPs) towards lung cancer screening guidelines in asymptomatic patients.

Methods: We conducted a cross-sectional descriptive study at the NGHA primary care centres, using a validated questionnaire, with 11 questions, developed by the National Cancer Institute, USA, and customized to our medical settings.

Results: 37% of family physicians requested chest X-rays; 1.9%, sputum cytology; and 3.7%, low-dose spiral chest computed tomography (CT) scans. Of the internists, 42.1% requested low-dose CT scans; 62.5%, chest X-rays; and 5%, sputum cytology. The mean knowledge scores for family physicians and internists regarding screening guidelines were 2.3 and 1.5, respectively, and those for their belief in the effectiveness of screening modalities were 7.6 and 6.6, respectively.

Conclusion: To harmonize lung cancer screening guidelines with clinical practice, further research on factors influencing the perceptions and responses of PCPs to screening guidelines needs to be conducted.

Keywords: lung cancer, physicians, belief and recommendations of PCPs, screening guidelines, family medicine, internal medicine, asymptomatic.

I. INTRODUCTION

Worldwide, lung cancer is considered fatal. Early detection via screening may aid in the decrease of cancer-related mortality rate. Till date, the optimal method for lung cancer screening is controversial (Nanavaty et al, 2014). However, regardless of these controversies, due to the lack of sufficient evidence, major medical experts and recent guidelines do not recommend screening in asymptomatic patients, even in those have histories of heavy or long-term smoking (Lung Cancer: Screening - US Preventive Services Task Force, 2013; NCCN Guidelines for Patients® | Lung Cancer Screening, 2016; Lung Cancer Screening Guidelines, Cancer.org, 2016; Care, 2015). Due to the increasing incidence of

lung cancer in Saudi Arabia, primary care physicians (PCPs) should have significant roles in preventing lung cancers and identifying those who are at risk; the choice of appropriate tools and candidates for screening is very crucial. The aim of this study is to explore the beliefs and attitudes of PCPs, towards lung cancer screening guidelines, in asymptomatic patients.

II. METHODS

A cross-sectional descriptive study was conducted at the National Guard Hospital (NGHA), Riyadh between January February 2017, using the validated lung cancer screening questionnaire developed by the National Cancer Institute (NCI), USA, in collaboration with the Agency for Healthcare Research and Quality, and the Centers for Disease Control and Prevention, USA. The questionnaire was edited and customized, by adding and eliminating questions, to be compatible with our medical setting.

All 146 PCPs, including family physicians and internists, were included in the study, without sampling. A pilot study was conducted on 10 physicians to ensure full comprehension of the questionnaire; this resulted in some modifications in vocabulary and format to avoid ambiguity. The King Abdullah International Medical Research Center (KAIMRC) also reviewed the survey tool. This contains questions related to the knowledge, attitudes, and demographics of the physicians, and takes approximately 10 minutes to complete.

Data management and statistical analysis were performed using the Statistical Package for Social Sciences (SPSS) software version 20.0. Frequencies and percentages were utilized to represent categorical variables, and the Chi-square test was used to investigate the relationship between variables. The knowledge scores were marked as follows: correct answers were marked with 1, and wrong answers, with 0. The sum of all knowledge questions was calculated for each participant. The knowledge scores were computed based on 11 questions from the questionnaire, and the answers were evaluated according to the guidelines mentioned in Figure 1.

The attitude scores were marked as follows: answers with positive attitude were marked with 1, and negative attitude, with 0. The sum of all attitude

Author a: College of Medicine, King Saud Bin Abdulaziz University for Health Sciences. e-mails: roaa1414@hotmail.com, Amer541@ksau-hs.edu.sa



questions was calculated for each respondent. Attitude scores were computed based on 8 questions. The means of these scores were compared between groups, using the Student's t-test. P-values of 0.05 or less were considered significant. Permission for conducting the study was obtained from KAIMRC in Riyadh. The cover sheet of the questionnaire explained that the physicians participated voluntarily in the study, and this was considered as consent. All data was treated anonymously.

51% and internal medicine (response rate, 48.7%) departments, participated in the study. The mean knowledge scores for the internists and family medicine physicians, regarding their belief in the effectiveness of the different screening modalities in reducing the lung cancer-related mortality in asymptomatic patients were 6.6 and 7.6 (P-value=0.54), respectively. The results of the first question on the questionnaire are shown in table-1.

III. RESULTS

A total of 74 PCPs (total response rate, 50.68%), including those from family medicine (response rate,

Table 1: Responses of family physicians and internists on the modality of choice for lung cancer screening per case

		Never smoked	Former smoker	Current smoker		
Very Effective Modality						
Chest X-ray	IM	5.26%	P-Value	10.4%	P-Value	21.05%
	FM	7.27%	0.7635	9.09%	0.8538	27.27%
Sputum Cytology	IM	0%	P-Value	5.26%	P-Value	21.05%
	FM	0%	-	1.1%	0.4247	14.5%
Low-dose CT	IM	26.31%	P-Value	57.89%	P-Value	57.89%
	FM	14.4%	0.2451	30.9%	0.0245	55.96%
Somewhat Effective modality						
Chest X-ray	IM	36.84%	P-Value	42.1%	P-Value	42.1%
	FM	18.18%	0.0955	36.36%	0.6564	32.27%
Sputum Cytology	IM	42.10%	P-Value	47.36%	P-Value	36.84%
	FM	18.18%	0.0361	31.1%	0.1951	27.25%
Low-dose CT	IM	15.8%	P-Value	5.26%	P-Value	5.26%
	FM	12.72%	0.8955	23.63%	0.0779	7.19%
Not Effective Modality						
Chest X-ray	IM	52.63%	P-Value	42.1%	P-Value	36.85%
	FM	72.72%	0.1067	47.45%	0.6968	36.63%
Sputum Cytology	IM	52.63%	P-Value	42.1%	P-Value	36.84%
	FM	70.82%	0.1464	51.7%	0.508	49.17%
Low-dose CT	IM	47.36%	P-Value	36.85%	P-Value	36.85%
	FM	65.65%	0.2131	38.2%	0.8079	35.25%

Do not Know the Effectiveness of Modality							
Chest X-ray	IM	5.27%	P-Value	5.4%	P-Value	0%	P-Value
	FM	1.83%	0.4247	7.1%	0.7635	36.63%	<0.001
Sputum Cytology	IM	5.27%	P-Value	5.28%	P-Value	5.27%	P-Value
	FM	1.83%	0.4685	16.1%	0.2223	9.08%	0.5982
Low-dose CT	IM	10.53%	P-Value	0%	P-Value	0%	P-Value
	FM	7.23%	0.6542	7.27%	<0.001	1.6%	<0.001

IM: Internal Medicine, FM: Family medicine, CT: computed tomography. P-value<0.05 is considered significant.

The mean knowledge scores for family and internal medicine physicians regarding lung cancer screening guidelines were 2.3 and 1.5 (P-value=0.48), respectively. The knowledge of the physicians,

regarding the various lung cancer screening guidelines for asymptomatic patients with histories of smoking less than 30 packs per year, or for those who have never smoked is illustrated in figure-1.




Figure 1: Physicians knowledge regarding lung cancer screening guideline and recommendation

The second question included various scenarios where the physicians were asked to choose the best lung screening modality, with the assumptions that the patients had not been previously screened, did not have any symptoms of lung cancer, did not express any preferences for lung screening, either in general or using a specific modality, and had no occupational exposure to known or suspected lung carcinogens. The responses to the second question are demonstrated in figure-2.




Figure 2: Responses by the physicians to question 2, with the assumptions that these patients had not been previously screened for lung cancer, did not have symptoms of lung cancer, did not express any preference for screening, either in general or with a specific modality, and had no occupational exposure to known or suspected lung carcinogens

According to their practice, 37% of family physicians requested chest X-rays, 1.9%, sputum cytology, and 3.7%, low-dose spiral chest computed tomography (CT) scans for lung cancer screening of asymptomatic patients in the past 12 months (figure-3).

Additionally, of 19 internists, 62.5% requested chest X-rays, 5%, sputum cytology, and 42.1%, low-dose spiral CT scans for lung cancer screening of asymptomatic patients in the past 12 months (figure-3).




Figure 3: The practice of physicians toward clinical screening tools for lung cancer

The average percentage of time spent by family physicians in providing medical care was 81.73%; in research, 8.18%; teaching, 9%; and others, 1.09% (administration, higher education, etc.). The average percentage of time spent by internal medicine physicians in providing medical care was 80.52%; research, 12.1%; and teaching, 7.38%. The average number of patients treated during a typical week by our

family or internal medicine physicians is demonstrated in figure-4, and the mean age groups of patients seen per specialty are shown in figure-5.




Figure 4: Percentage of patients seen by physicians per week




Figure 5: Average of patients' age according to specialty

IV. DISCUSSION

Worldwide, lung cancer is considered fatal, with a 5-year survival rate of 16% in the United States (Dela Cruz et al, 2011; Alamoudi, 2010). During 2013, 212,584 new cases of lung cancer were diagnosed in the United States, and over 156,176 patients died (Fact Sheets by Cancer. Globocan.iarc.fr. 2016). In the United Kingdom, 34,000 new cases are documented annually (Spiro et al, 2005). According to the latest Saudi National Cancer Registry, 397 cases were diagnosed in 2010, accounting for 4% of cancers (King Faisal Specialist Hospital & Research Centre - Centers of Excellence - Cancer Centre. Telemedicine.kfshrc.edu.sa. 2016).

Early detection of lung cancer via screening is greatly expected to aid in decreasing the mortality rate. The optimal method for lung cancer screening has not

yet been established (Nanavaty et al, 2014). However, regardless of these controversies, due to lack of sufficient evidence, major medical experts and recent guidelines do not recommend screening in asymptomatic patients, even if they have histories of heavy or long-term smoking (Lung Cancer: Screening - US Preventive Services Task Force, 2013; NCCN Guidelines for Patients® | Lung Cancer Screening, 2016; Lung Cancer Screening Guidelines, Cancer.org, 2016; Care C, 2015).

Furthermore, a similar study conducted in 2012 in the United States aimed to explore the beliefs and attitudes of PCPs regarding lung cancer screening. The results showed that one-fourth of all PCPs recommended screening in asymptomatic patients, and one-half requested chest radiographs (Klabunde et al, 2012). In 2010, another study showed similar results; it

was found that the beliefs of many PCPs regarding lung cancer were inconsistent with the current guidelines and recommendations (Klabunde et al, 2010).

The findings of this study were consistent with existing research and theoretical evidence, which suggests a progressive increase in the incidence of lung cancer in Saudi Arabia, and indicates the challenges encountered in the timely recognition of lung cancer. The research indicated that the average knowledge of family physicians regarding lung cancer screening guidelines was minimal, although they reported a score of 6.6 with respect to their beliefs about the importance of lung cancer screening, and between practice and beliefs with a score of 2.2. This trend continues, despite recommendations by numerous organizations for lung cancer screening.

Previous research has determined that the recommendations by physicians are important predictors of health-seeking behaviour in patients (Dela Cruz et al, 2011). Limited knowledge of the guideline recommendations is likely to inhibit physicians from facilitating collective decision-making conversations, concerning the possible advantages, uncertainties, and disadvantages of lung cancer detections, when interacting with their patients (Spiro et al, 2005, National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcaredelivery.cancer.gov. 2016).

The results indicated that internal medicine physicians had knowledge scores of 1.5, regarding lung cancer screening guidelines. The findings of this investigation are in agreement with a previous survey in 2011 that examined lung cancer screening practices (Lung Cancer Screening (PDQ®)-Health Professional Version. National Cancer Institute. 2016). The survey illustrated that family physicians have higher preferences for requesting chest radiographs in asymptomatic patients with lung cancer, compared to internists.

The research findings showed that the lung cancer screening recommendations and beliefs of many PCPs were inconsistent with current evidence and guidelines. The study considered the key modalities that have been utilized in lung cancer screening, which include chest X-rays, low-dose CT scans, and sputum cytology. Most internists believe that low-dose CT is efficient in decreasing lung cancer mortalities among current smokers; this concurs with the assertion of previous researchers (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcaredelivery.cancer.gov. 2016; Lung Cancer Screening (PDQ®)-Health Professional Version. National Cancer Institute. 2016). Those physicians who had completed medical school more than 15 years ago were more likely to request for chest radio graphs for lung cancer detection. These results concur with physician beliefs concerning screening. Senior

physicians appear to request the specific lung cancer detection method that is compatible with their prior medical training (Humphrey et al, 2013).

The results show that the demands of patients are related to the physician reports regarding requests for lung cancer detection, and concur with previous studies demonstrating that the requests of patients affect the physician ordering a test for cancer susceptibility. Physician evaluations linked with lung cancer detection might aid in targeting the involvement of physicians that are in dire need of information and evidence regarding lung cancer detection guidelines (Mazzone et al, 2015). Most PCPs (89.09%) acknowledged that they recommended against screening in patients who were over 50 years of age, who had never smoked, or who did not have substantial exposure to passive smoking, but only 36.36% of PCPs recommended against screening for lung cancer inpatients who were either former or current smokers, including those exposed to passive smoking.

We analysed physician preferences for the best screening modalities for patients that have not been previously screened, have no symptoms of lung cancer, have not expressed a preference for lung cancer screening, either in general or with a specific modality, and have not had any prior exposure to known or suspected lung carcinogens. The results demonstrated that 37% of family physicians requested chest X-rays; 1.9%, sputum cytology; and 3.7%, low-dose spiral chest CT scans, for lung cancer detection. However, among the internists, 42.1% requested low-dose CT scans for lung cancer detection; 62.5%, chest X-rays; and 5%, sputum cytology. From the above results, it appears that primary care physicians in the King Abdulaziz Medical City (KAMC) primary care centre have not decreased the practice of requesting chest X-rays to detect lung cancers in asymptomatic individuals.

Nonetheless, among PCPs who recommend the screening of patients for lung cancer, 63.15% of internists and 36.36% of family physicians recommend the use of chest X-rays, which is not a recommended test (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcaredelivery.cancer.gov. 2016). This result is consistent with the previous understanding of the national provider, to examine lung cancer screening practices before launching the NCI guidelines. About 26.31% of internal medicine physicians viewed low-dose CT as an effective screening modality, compared to 14.4% of family physicians. The propensity of PCPs to suggest a particular screening technique increases with the exposure of the patient to smoking.

The knowledge of guidelines was not associated with the utilization of low-dose CT; surprisingly, despite only 31.5% of internists knowing the NCI, compared to 40% of family physicians, the use of

low-dose CT scans was the highest among the internists. The use of chest-X-rays is partly accredited to concerns about financial costs, the unavailability of other screening modalities, and lack of insurance coverage (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcare delivery.cancer.gov. 2016). The small percentage (13.01%) of all PCPs who possessed lung screening programs or aids in their work setting may be attributed to the slow uptake of low-dose CT, and the increased usage of chest X-rays becoming more common due to extensive coverage. Nevertheless, the National Lung Screening Trial (NLST) presents evidence that lung cancer detection with low-dose CT is more efficient than with other commonly performed screening interventions, such as sputum cytology and chest X-rays (Lung Cancer Screening (PDQ®)-Health Professional Version. National Cancer Institute. 2016).

The research findings have shown the beliefs of PCPs concerning practice guidelines, test effectiveness, and tendency to intensify for any cancer that is highly related to the lung cancer screening recommendations. They substantiate an earlier, but much smaller study, which suggests that aggressive cancer screening by family physicians is related to their beliefs (Alamoudi, 2010).

The requests by PCPs for unverified lung cancer screening techniques have various implications. One such implication is the potential psychological harm that results from false-positive or false-negative results (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcare delivery.cancer.gov. 2016). Physical damage can also result from unwarranted invasive procedures that are undertaken as follow-up for false-positive screening (Klabunde, 2012). The use of unrecommended lung cancer technologies will eventually drive up health care costs. According to the data from the NLST, an average of 30% of patients who undergo low-dose CT scanning as a detection procedure will have at least one false-positive screening (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcare delivery.cancer.gov. 2016). Moreover, three of every 1,000 persons screened are estimated to develop major complications associated with the procedure, and three-to-five people may be over-diagnosed with lung cancer (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcare delivery.cancer.gov. 2016). There is a need for current and future refinements in screening; for instance, raising the threshold for a "positive" result of screening in the national guidelines, and adopting calculators to predict

the likelihood of lung cancer, which will help lower the false-positives rate (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcare delivery.cancer.gov. 2016). The results from this study showed that PCPs had glaring gaps in their knowledge of lung cancer screening; this screening tends to happen opportunistically rather than through well-organized programs (Lung Cancer Screening (PDQ®)-Health Professional Version. National Cancer Institute. 2016; Klabunde, 2010). This, in turn, may result in a considerable percentage of incorrectly screened adults, unless intensive efforts are made to notify PCPs of the proper explanation of the NLST results, and better screening approaches in clinical practice.

The study suggests that only 35% of PCPs promote lung cancer screening by initiating conversations with the patients regarding the advantages and risks of undertaking such screening; this limited number results from the lack of familiarity with the clinical practice guidelines for lung cancer detection. The existing evidence does not support screening for asymptomatic patients, not withstanding their exposure to smoking (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcare delivery.cancer.gov. 2016). This may be because being attuned to the current practice guidelines can be a daunting task for physicians (Klabunde, 2010). The proliferation of several guidelines may negatively affect the ability of PCPs to adhere to them. The use of an academic detailing approach may encourage supportive attitudes and beliefs towards lung cancer screening, as well as, inspire disease advocacy groups, and encourage the availability of technology that facilitates screening (Nanavaty et al, 2016).

Strengths and limitations: A major strength of this study was the extent of evidence available in literature regarding the research objectives. The results reflect the views of both inexperienced and experienced providers with diverse clinical understanding, from the two predominant fields of primary care services, namely, family medicine and internal medicine.

A limitation of the study is based on the attitudes, recommendations, and practices of physicians; these were obtained through a self-reported questionnaire that was not verified using any other sources, such as medical claims or reports. To reduce the workload of the respondents, the survey questionnaire on lung cancer detection was comparatively short, and it did not have the capacity to extract details about specific features of the patients for whom the PCPs had requested lung cancer screenings, and their extent and type of smoking exposure. Moreover, the study relied on PCP accounts of screening behaviour, which are subject to recall bias or

social desirability. Lastly, the low response rate (51%) did not allow generalization of the results of the study to other primary care centres in Riyadh and Saudi Arabia.

V. CONCLUSION

Additional research is warranted to educate PCPs and health care consumers, concerning the need, evidence base, guidelines, cost, and potential risks, of lung cancer screening guidelines. The public has an exaggeratedly positive view of cancer detection, albeit with an inadequate understanding of the potential damage. The utilization of CT scans is rising rapidly in KAMC primary care centres. There is a need to address current barriers, such as, insurance coverage, financial cost, frequency of false-positive results, and associated complications with screening.

These initiatives will be essential in providing PCPs with the necessary knowledge to make decisions regarding lung cancer screening (National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcaredelivery.cancer.gov. 2016). These pertinent concerns, in conjunction with study findings and developing an evidence base, highlight the significance of continuously monitoring the knowledge, practices, and beliefs of PCPs, as these are interconnected with lung cancer detection. Further research is required to enumerate the factors influencing the perceptions and responses of PCPs, regarding lung cancer screening guidelines, to enhance the current understanding of these guidelines.

The present study uncovers the disconnect that exists amid evidence and practice in lung cancer detection, and explores critical background for reflection on the results of the significant and extremely publicized NSLT (Lung Cancer Screening (PDQ®)-Health Professional Version. National Cancer Institute. 2016). Most of the guidelines are very specific to people aged between 50–75 years, who are either current smokers or were heavy smokers with histories of smoking a minimum of 30packs per year(National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Healthcaredelivery.cancer.gov. 2016). Therefore, any alteration in the screening commendations would only apply to this select and high-risk population. Lung cancer screening is established concept in Saudi Arabia, and the information regarding the usage of lung cancer screening in Saudi Arabia is quite limited and is confined to the national data sources, which provide population estimates of cancer screening. There is a need to harmonize lung cancer screening guidelines and clinical practice.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Alamoudi OS. 2010 Lung cancer at a university hospital in Saudi Arabia: A four-year prospective study of clinical, pathological, radiological, bronchoscopic, and biochemical parameters. *Annals of Thoracic Medicine*.vol 5 no. 1 pp 30-36. doi:10.4103/1817-1737.58957.
2. Care C. Screening for Lung Cancer (2015) [Internet]. Canadian Task Force on Preventive Health Care. 2016 [cited 4 November 2016]. Available from: <http://www.cmaj.ca/content/188/6/425>
3. Dela Cruz CS, Tanoue LT, Matthay RA. Lung Cancer 2011 Epidemiology, Etiology, and Prevention. *Clinical Chest Medicine*.vol 32 no. 4 pp 605-44. doi: 10.1016/j.ccm.2011.09.001. 2016.
4. Fact Sheets by Cancer [Internet]. Globocan.iarc.fr. 2016 [cited 4 November 2016]. Available from: http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx
5. Final Update Summary: Lung Cancer: Screening - US Preventive Services Task Force, 2013 [Internet]. Uspreventiveservicestaskforce.org. 2016 [cited 4 November 2016]. Available from: <https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/lung-cancer-screening>
6. Humphrey LL, Deffebach M, Pappas M, Baumann C, Artis K, Mitchell JP, et al. 2013. Screening for lung cancer with low-dose computed tomography: a systematic review to update the US Preventive services task force recommendation. *Annals of Internal Medicine*. vol 159 no. 6 pp 411-20. doi: 10.7326/0003-4819-159-6-201309170-00690. PubMed - NCBI [Internet]. Ncbi.nlm.nih.gov. 2016 [cited 4 November 2016]. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23897166>
7. King Faisal Specialist Hospital & Research Centre - Centers of Excellence - Cancer Centre [Internet]. Telemedicine.kfshrc.edu.sa. 2016 [cited 4 November 2016]. Available from: https://telemedicine.kfshrc.edu.sa/wps/portal/En/ut/p/c0/04_SB8K8xLLM9MSSzPy8xBz9CP0os_jQEH9nSydDRwMLQwsDA09DEycfQw83lwMnE_2CbEdFAAMjHY4!/?WCM_PORTLET=PC_7_UTOC9B1A081800I14BL1HF20R2_WCM&WCM_GLOBAL_CONTEXT=/wp/s/wcm/connect/k
8. Klabunde CN, Marcus PM, Han PK, Richards TB, Vernon SW, Yuan G, Silvestri GA 2012. Lung cancer screening practices of primary care physicians: results from a national survey. *Annals of Family Medicine*.vol 10, no. 2 pp 102-10. doi:10.1370/afm.1340. - PubMed - NCBI [Internet]. Ncbi.nlm.nih.gov. 2016 [cited 4 November 2016]. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22412001>
9. Klabunde CN, Marcus PM, Silvestri GA, Han PK, Richards TB, Yuan G, et al. 2010. U.S. primary care

physicians' lung cancer screening beliefs and recommendations. American Journal of Preventive Medicine.vol 39 no. 5 pp 411-20. doi: 10.1016/j.amepre.2010.07.004.- PubMed - NCBI [Internet]. Ncbi.nlm.nih.gov. 2016 [cited 4 November 2016]. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20965378>

10. Lewis J, Petty W, Tooze J, Miller D, Chiles C, Miller A, et al. 2015. Low-Dose CT Lung Cancer Screening Practices and Attitudes among Primary Care Providers at an Academic Medical Center. Cancer Epidemiology Biomarkers & Prevention.vol 24 no. 4 pp 664-70.
11. Lung Cancer Screening (PDQ®)-Health Professional Version [Internet]. National Cancer Institute. 2016 [cited 4 November 2016]. Available from: <https://www.cancer.gov/types/lung/hp/lung-screening-pdq>
12. Lung Cancer Screening Guidelines [Internet]. Cancer.org. 2016 [cited 4 November 2016]. Available from: <http://www.cancer.org/healthy/informationforhealthcareprofessionals/acsguidelines/lungcancerscreeningguidelines/index>
13. Mazzone P, Powell C, Arenberg D, Bach P, Detterbeck F, Gould M, et al. 2015 Components Necessary for High-Quality Lung Cancer Screening: American College of Chest Physicians and American Thoracic Society Policy Statement Chest. Vol 147 no. 2 pp 295-303. doi:10.1378/chest.14-2500. [Internet]. Journal.publications.chestnet.org. 2016 [cited 4 November 2016]. Available from: <http://journal.publications.chestnet.org/article.aspx?articleid=1921603>
14. Did you mean, "National Cancer Institute. National Survey of Primary Care Physicians' Recommendations & Practice for Breast, Cervical, Colorectal, & Lung Cancer Screening. Bethesda, MD: National Cancer Institute; 2008 [updated 2008 Oct 8; cited 2008 Dec 20]. Available from: http://healthservices.cancer.gov/surveys/screening_rp/
15. NCCN Guidelines for Patients® | Lung Cancer Screening [Internet]. Nccn.org. 2016 [cited 4 November 2016]. Available from: https://www.nccn.org/patients/guidelines/lung_screening/files/assets/basic-html/page-1.html#
16. Ries LA, Eisner MP, Kosary CL, Hankey BF, Miller BA. SEER Cancer Statistics Review, 1975-2003, National Cancer Institute. Available from: https://seer.cancer.gov/archive/csr/1975_2003/results_merged/sect_15_lung_bronchus.pdf
17. Spiro SG, Silvestri GA. 2005 One Hundred Years of Lung Cancer. American Journal of Respiratory and Critical Care Medicine.vol 172 no. 5 pp 523-529.

