Artificial Intelligence formulated this projection for compatibility purposes from the original article published at Global Journals. However, this technology is currently in beta. *Therefore, kindly ignore odd layouts, missed formulae, text, tables, or figures.* 

# Stress Test Performance by Preterm-Born Females in Adolescence and Adulthood

Katarzyna Horosz

Received: 15 December 2016 Accepted: 3 January 2017 Published: 15 January 2017

#### 6 Abstract

1

2

3

Preterm birth often entails developmental disorders, with the circulatory and respiratory 7 systems are particularly at risk, yet few studies have examined long-term effects. To evaluate 8 the long-term impact of preterm birth on stress test performance in women born preterm, we 9 administered the same stress test to the same set of preterm-born subjects, first at the age of 10 puberty, then again in adulthood, comparing the results. The strength of the relationship 11 between anthropometric parameters and physical fitness, as well as estimated oxygen uptake 12 were also analyzed. A total of 70 girls (aged  $12.2\pm1.5$ ) who had been born preterm ( $34.7\pm1.86$ 13 weeks) were tested in 1997. Of those, after a gap of 18 years, a group of 13 as successfully 14 re-contacted and participated in the 2015 examination as adults (then aged  $27.6\pm2.60$  years, 15 born preterm at  $34.5\pm2.0$  weeks). Each time, an indirect HR test was performed (W150 test) 16 while striving to maintain HR=150 bpm; the level of physical load (W) therefore depended 17 only on physical fitness and exercise tolerance. In the group of adolescent girls, the rate of 18 stress oxygen uptake (O 2) and the metabolic stress response were measured by using the 19 Jaeger ErgoOxyscreen apparatus. In the adult group, the VO 2 max rate was measured. 20

22 Index terms—

21

# 23 1 Introduction

reterm birth, defined as birth prior to 37 completed weeks of gestation and generally associated with low birth weight, is a widespread phenomenon, accounting for 5% to 9% of all births in Europe [1], and more than 12% of all births in the United States [2], with similar high rates in Africa and southern Asia [3].

Although morbidity is still significant among very low birth weight infants, numerous technological advances, 27 collaborative efforts between obstetricians and neonatologists, widespread use of antenatal corticosteroids, 28 surfactant therapy, and high-frequency ventilation have all helped improve survival rates over the past two 29 decades [4][5][6]. Since individuals who were born preterm are now more likely to survive into adulthood than 30 before, a comprehensive understanding of the impact of premature birth in adulthood is therefore needed to 31 enable earlier prevention, detection, and treatment of its long-term health sequelae. However, such information 32 is largely lacking from the literature; most studies of the impact of premature birth have examined subjects only 33 in childhood. 34

35 The literature on exercise tolerance focuses mainly on preterm-born children in the early years of their lives, 36 generally not proceeding beyond age ten. As such, it has yet to be determined whether premature birth is also 37 associated with reduced physical fitness A total of 70 girls (aged  $12.2\pm1.5$ ) who had been born preterm ( $34.7\pm1.86$ 38 weeks) were tested in 1997. Of those, after a gap of 18 years, a group of 13 as successfully re-contacted and participated in the 2015 examination as adults (then aged  $27.6\pm2.60$  years, born preterm at  $34.5\pm2.0$  weeks). 39 Each time, an indirect HR test was performed (W150 test) while striving to maintain HR=150 bpm; the level of 40 physical load (W) therefore depended only on physical fitness and exercise tolerance. In the group of adolescent 41 girls, the rate of stress oxygen uptake (O 2) and the metabolic stress response were measured by using the Jaeger 42

43 ErgoOxyscreen apparatus. In the adult group, the VO 2 max rate was measured.

In adolescence, no difference was found between the study group and the reference group. In adulthood, however, the prematurely born adults demonstrated significantly lower values of VO 2 max and total work performed in relation to the reference group, pointing to a decline in physical fitness in the study group.

47 Physical fitness is always conditioned by numerous factors, and this even more so in the case of developmental 48 disorders (such as those resulting from premature birth). For instance, respiratory and circulatory capacities depend on the maximum oxygen uptake (VO 2 max) and, as such, full lung maturation is of paramount importance 49 in determining physical fitness [7]. Premature infants generally have difficulty breathing due to a reduced number 50 of alveoli and underdeveloped pulmonary capillaries [8]. Such pathological conditions at an early age may result 51 in diminished lung function later in life, including in terms of lung volume, ventilation homogeneity, and the 52 mechanics of the respiratory system, which will in turn detract from physical fitness [9][10][11][12]. Alongside the 53 respiratory system, the circulatory system also plays a significant role in the development of the child's physical 54 fitness. Both systems are involved in the process of gas exchange through the blood. In premature babies, 55 the circulatory system may experience a delayed closure of the ductus arteriosus and the pulmonary vascular 56 resistance may be diminished after birth; this condition (patent ductus arteriosus or PDA) significantly impairs 57

58 the respiratory function and hemodynamic conditions of premature infants.

The aim of the present study, therefore, was to explore the long-term effect of prematurity by examining the stress-test performance of subjects at age 12, then to re-examine a portion of the same subjects at the age of 30, using same examination setup, and to compare the results against those of a reference group. The strength of the relationship between anthropometric parameters (height and weight, BMI) and physical fitness, as well as estimated oxygen uptake were also analyzed.

# 64 **2** II.

# 65 **3** Material and Methods

## 66 4 a) Participants

Individuals participating in the study group for both stages of our study were females who had been registered at the Premature Birth Clinic, Rehabilitation Department, Institute of Mother and Child in Warsaw, as having been born with low birth weight, i.e. below 2,500 g, or born preterm, i.e. prior to 37th week of pregnancy (though with normal birth weight). The total work performed by the subjects on an exercise test was evaluated twice: once in 1997 (puberty) and again in 2015 (adulthood). An additional criterion for inclusion of preterm-born individuals in both states of the study was a normal resting ECG.

In the first stage, in 1997, total of 70 pretermborn girls aged 10-14 years (12.22±1.52 years) took part in the examinations (the results of which have not been previously published). The mean birth weight of among this the group was 1,865.8±566.3 g (min. 1,040 g, max. 2,580 g), and they were born in the 34.5±1.92 week (min. 32 weeks, max. 36 weeks). All of the subjects who ended up selected based on the above criteria, therefore, were all under 2,500 g birthweight and all born prior to 37th week of pregnancy. Most of the study group (85%) had been fed on formulas as infants, whereas two had been fed on a combination of infant formula and breast milk. On the basis of their health records, none of the subjects had experienced BPD.

Eighteen years later, we attempted to reestablish contact with all of the 1997 participants by twice sending 80 out a request letter to their previously recorded residence. Only 13 of the original participants responded and 81 agreed to undergo reexamination in 2015. The study group participants in 2015 had a mean birth weight of 82  $1,864.6\pm533.7$  g (min. 1,040 g, max. 2,580 g) and were born at  $34.5\pm2.0$  weeks (min. 32, max. 36) of pregnancy. 83 84 Note that, in fact, both boys and girls had been examined in the 1997 stage of the study, for which all 85 prematurely born and/or low-birth-weight children registered with the above-mentioned clinic who met the criteria for the study had been recruited. However, the 13 participants who ultimately agreed to participate 86 in the follow-up stage later turned out to be exclusively female; as such, we were able to analyze and report 87 long-term results exclusively for the female group. This sample size of 13 participants for the latter, follow-up 88 stage, while admittedly quite small in absolute terms, still represents a considerable achievement given the long 89 time-frame involved and the relative scarcity of such long-term data in the literature. 90

This study group was in each case compared against a reference group of their peers, consisting of 48 girls ( $12.4 \pm 1.52$  years) in the first stage, and of 27 women ( $28.3 \pm 2.16$  years) in the second, made up of girls and women from the province including Warsaw and its environs (the Mazowsze Voivodship), born at term, at ages corresponding to the test group. To compensate for social variables, the reference group was in each case recruited by asking the study participants (or their guardians) to invite their peers born at term and weighing more than 2,500g to take part in the study.

The basic parameters of the two groups are presented in Table 1. All participants were informed about the conditions and course of the study, and written informed consent was expressed for participation in the research (either from the legal guardians of the children or by the adult participants themselves). The study received the approval of the Ethics Committee at the Józef Pi?sudski University of Physical Education in Warsaw (decision no. SKE 01-47/2012). later in life. However, despite consistent findings concerning poor lung function, few studies have addressed exercise tolerance in low birth weight populations [13][14][15][16]. Authors have reported decreased [13,14] or relatively normal [15,16] peak VO 2 values in young prematurely born children. Contrary to these finding, in another study we previously reported finding an improvement in exercise tolerance in cohorts of preterm born children in adolescence [17]. Data concerning adults who were born prematurely are scarce, as there have been few longitudinal or follow-up studies extending into adulthood. Clemm et al. [18] compared the exercise capacity of adults born at term and EP, addressing the developmental patterns from adolescence to adulthood (18 to 25 years old) and reporting that exercise capacity was slightly lower in EPborn adults; however, those values were within a normal range, positively associated with self-reported physical activity, and unrelated to neonatal factors and current airway obstruction.

# <sup>111</sup> 5 Volume XVII Issue III Version I

# 112 6 b) Study Methods

Two stress-test examinations were carried out at an interval of 18 years, the first stage in 1997 and the second in 113 2015. All measurements were performed in the early hours of the morning at a diagnostic facility of the Central 114 Laboratory at the University of Physical Education in Warsaw. Tests were performed at least two hours after the 115 last meal. In each case indirect HR test (W150 Test) was performed using an ERGOTEST device, based on the 116 original W150 program [19]. Physical load values were determined by the computer using feedback (load/HR); 117 this results in the induction of cyclometer resistance at a certain level, forcing the achievement and maintenance 118 of physiological balance at HR=150 bpm (control level) to evaluate aerobic metabolism. The test uses a linear 119 relationship between the frequency of the heart rate and oxygen uptake. The effort was dispensed on a bicycle 120 ergometer Pro Med Medical Systems with a foot drive, heart rate were recorded and measured electronically 121 using precordial electrodes (transmitter BMI-USBD1-EU-S/N 054BM/S300). One of the criteria for qualifying 122 was a normal resting ECG. As the test was underway, for the study group individuals (preterm-born) their ECG 123 was monitored by a physician (Nõrava S-stress, Oxford Poland). 124

Before undergoing the exercise test, participants were subjected to a preliminary assessment of exercise capacity using the Ruffier test, which included measuring their blood pressure and heart rate (Dura Schock Welch Allyn sphygmomanometer). Warmup was performed on an ergometer driven by lower limbs for 10 minutes, with a load of 1W/kg of the subject's body weight. Heart rate during warm-up did not exceed HR=130 bpm. After the warm-up stage and recovery period, the proper testing began.

Tests were carried out in the presence of a physician, and were explained to the participants before they 130 began. The period of exercise lasted 10 minutes, followed by a recovery period lasting until the participant's HR 131 returned to pre-test level. The physical challenge method applied (also used in the 1997) assumes the maintenance 132 of the heart rate on control level of HR=150 bpm; load regulation is based on the individual reactions of the 133 cardiovascular system, with the load level (W) being adjusted as necessary during the exercise and with pedal 134 cadence ranging from 50 to 60/min. The load level therefore depended only on the subject's physical fitness and 135 exercise tolerance. During the exercise test the following parameters were recorded: HR, total work performed 136 during the test, power necessary to achieve the control level (HR=150 bpm), average power for the control period 137 138 (measured from the moment of attaining HR=150 bpm until the end of the test), and the time it took to achieve 139 the control level (HR=150 bpm). At the completion of the test in the adult group the VO 2 max rate was 140 automatically calculated. In the group of adolescent girls, the rate of oxygen uptake and the metabolic stress response were measured using the Jaeger ErgoOxyscreen apparatus. 141

The test was halted for in the event of any adverse cardio-respiratory reactions or signs of exercise intolerance, such as dizziness, shortness of breath, pallor, chest pain or imbalance.

# 144 7 Statistical analysis

The data was processed and all statistical calculations were made using STATISTICA 12.0 by StatSoft, according to the software instructions. The results were presented as arithmetic means  $\pm$  standard deviation. The Shapiro-Wilk test was used to evaluate the normality of distributions, and the Pearson correlation coefficient was used to measure the dependency between the parameters.

The variables were compared between the groups using a non-parametric test (the Mann-Whitney U test 149 adjusted for continuity). The ANCOVA test (from the General Linear Model) was used to exclude a covariate 150 (weight) in group comparisons. Because the distribution was not found to be normal, the Mann-Whitney U test 151 was used. The threshold for statistical significance was taken to be ?=0.05. We considered that the total work 152 153 value could be dependent on body weight, and so performed an ANCOVA analysis with covariate. The results 154 indicated that the body weight did not affect the significant differences in the total work performed between the groups (F = 8.97; p = 0.004749; ? = 0.1869). The test power was 0.83. In this study, we sought to trace 155 the long-term impact of preterm birth on physical fitness into puberty and later into adult life. Such long-term 156 consequences of preterm birth are complex and notoriously problematic to study, particularly in maintaining 157 the same group of subjects over a prolonged period of time. A major strength of our study, therefore, was its 158 longitudinal design. 159

## 160 8 III.

161 9 Results

## 162 10 Factor

#### 163 11 Correlations

164 However, maintaining the participation of test subjects from childhood through adulthood is indeed challenging, 165 and we were only able to recruit relatively few participants (13 out of 70, or 19%) to persuade them to take part 166 in the follow-up stage of the study. The original participants who did not participate in the followup examination may have done so for any of a variety of reasons, including change of residence in the interim, Among the variables 167 considered in the first study (adolescent period), no significant differences were found between the study group 168 and the reference group in terms of the parameters indicative of exercise tolerance at the peak of oxygen uptake 169 -i.e. O 2 [L/min] at HR=150 bpm and the total work performed W [kJ]. The total work performed, however, 170 was found to differ significantly between the study and reference groups of adults (p < 0.001). Similar differences 171 172 were observed in the VO 2 max rate, with the adult reference group presenting significantly higher levels of VO 173 2 max than their peers born prematurely (p<0.001). This indicates that adult women born prematurely do have 174 a lower exercise tolerance. No strong correlation (understood as p? 0.5) was found between the total work and body weight, between the total work and body height, or between the total work and BMI. The correlation 175 176 between the total work and VO 2 max, on the other hand, at p? 0.5, is consistent with the exercise response of 177 healthy individuals (the greater the work performed, the higher the VO 2 max rate).

## 178 12 Global Journal of

179 Medical Research

## 180 **13** IV.

#### $_{181}$ 14 Discussion

failure to receive the request letter, unwillingness or inability to participate, lack of interest, etc. Even so, given the relative scarcity of longitudinal data of this sort, we still do consider it a success to have managed to reexamine this sizeable of a share of the original group after such a time gap, and the proportion of reparticipation we achieved is nevertheless comparable to those reported in most similar studies [13,15,20]. Moreover, the magnitude of the effect we found was large (eta2>0.18), which ensured sufficient observed strength of the test (Power>0.80). This power analysis indicates that the sample size was sufficient to evaluate the phenomenon studied.

188 Most previous studies of children and adolescents, by contrast, have been cross-sectional [14][15][16]21]. 189 Vrijlandt et al. [16] found a mildly lower exercise capacity in a group of preterm individuals occurring in young adulthood, as compared to healthy, reference-group subjects. Moreover, their study group showed significantly 190 lower anaerobic threshold than their healthy peers, and tended to have lower work efficiency. However, this 191 might stem from less intensive participation in sporting activities, rather than impaired lung function or 192 limited ventilation. Kriemler et al. [15] reported that children with and without respiratory abnormalities 193 have some degree of pulmonary dysfunction at rest and following exercise, and a higher prevalence of exercise-194 induced bronchoconstriction with no reduction in maximal aerobic exercise performance. The higher oxygen 195 uptake seen at a given mechanical power in the preterm group may cause early fatigability during prolonged 196 exercise, even when aerobic performance is normal. Rogers et al. [14] reported significant differences between 197 198 low birth weight adolescents and term-born reference subjects in terms of lower aerobic capacity, strength, endurance, flexibility, and activity level. This again could be the result of less intensive participation in 199 physical activity and consequential physical limitations, such as neurodevelopment, sensory (visual and auditory 200 deficits), respiratory and cardiovascular impairments, reduced muscle strength, coordination difficulties and social 201 acceptance. motivating them to engage in this type of test, maximal exercise loads were not used. 202

In comparing our results for the study group at adulthood to those obtained during at puberty, we can conclude 203 that the anticipated deficits in their level of exercise tolerance in comparison to healthy peers nevertheless did 204 manifest themselves in adulthood, despite having been compensated for and even improved upon during puberty 205 (when preterm-born subjects fared 5% better than their healthy peers). Like other authors [18], we should 206 note that this could be attributable to a failure to maintain normal levels of physical activity after puberty due 207 208 to insufficient medical supervision and reduced parental control, but at the same time we cannot rule out the 209 late consequences of abnormal morphological and functional development in childhood due to premature birth. 210 Overall, premature birth contributes to increased risk of cognitive, sensory, neuromotor and coordination deficits, 211 which can interfere with the development of exercise tolerance. Less intensive physical activity may also result 212 from real or perceived limitations and weaknesses, leading to the reinforcement of negative attitudes towards exercise, and this in turn may have adverse effects on exercise tolerance [16,22,23]. 213

Baraldi and Filippone [24], however, disagree with this reasoning, arguing that early lung damage may itself have long-term consequences. Chronic lung disease cannot be considered solely as a pediatric illness because it is likely to continue into adulthood and increase the risk of acquiring chronic obstructive pulmonary disease

(COPD), as well as have a negative On the other hand, in our the second, follow-up testing of part of the same 217 preterm-born subjects in adult age, we found that, to the contrary, participants performed significantly worse in 218 terms of exercise tolerance than their peers from the reference group. This finding suggests that the evolution 219 of physical fitness and the consequences of prematurity need to be closely monitored into later stages of life. To 220 our knowledge, there is only one publication that addresses the developmental trajectories of exercise tolerance 221 from adolescence to adulthood: Clemm et al. [18]. In their study of 25-year-old adults born EP, they found that 222 exercise tolerance was 10% lower than in a reference group born at term, albeit still within a range considered 223 normal. A decreasing trend in exercise tolerance from 18 to 25 years of age was similar in both groups, and the 224 VO 2 peak obtained at age 18 strongly predicted the VO 2 peak at age 25. Exercise tolerance was unrelated 225 to neonatal factors and to current airway obstruction, but was positively associated with self-reported physical 226 activity. Our results similarly show that participants in the study group performed 22% less work than the 227 reference group. The physical load causing the same physiological response (maintaining HR at 150 bpm) was 228 significantly lower in the case of the study group. 229

Our results, on the other hand, differed from those reported in these studies: in our study group, preterm-230 born subjects at puberty actually achieved better results of total work and peak oxygen uptake with heart rate 231 maintained at 150 bpm than their peers born at term. These results indicate that, when it comes to submaximal 232 233 exercise, the physical functions of the preterm children we studied were not impacted by developmental limitations 234 in the first years of life. Of course, it could have been of more significance to study the exercise VO 2 max rate 235 in order to assess the children's physical capacity, but due to the state of health of preterm-born children and the difficulty in impact on the results of pulmonary function testing. Walter et al. [25] draw similar conclusions, 236 having found a higher risk of lung dysfunction in a group of adults born prematurely with low birth weight 237 in relation to their peers born at term. There are few studies in the literature on the effects of cardiovascular 238 disorders resulting from premature birth on exercise tolerance. Lewandowski et al. [26] performed an MRI on 234 239 individuals, and found increased left ventricular mass in young adults born prematurely, compared to the reference 240 group. Changes in the geometry of the heart were accompanied by abnormal hemodynamic parameters. The 241 authors argue that these disorders may consequently lead to reduced exercise tolerance, and may be associated 242 with an increased risk of cardiac events. 243

In general, the long-term consequences of premature birth remain not well documented, but it is becoming clearer that its complications do not only affect individuals in childhood. As the results of our study indicate, despite the fact that subjects born prematurely had much better performance results in adolescence than their peers in the reference group, later in life their exercise tolerance decreased and the same individuals fared worse as adults than those in the reference group. Therefore, the exercise tolerance preterm-born individuals needs to be measured at different stages of life, not only during puberty. Further research in this direction may help us better understand the complex, longer-term developmental trajectories of preterm-born individuals.

# **251 15 Conclusions**

1. During adolescence, prematurely born subjects displayed better aerobic metabolism and higher rates of total

work performed on a stress test than their peers born at term. 2. A subset of the same group of subjects
 nevertheless showed a significant decrease in the level of exercise tolerance in adulthood, as compared to their peers born at term.



Figure 1: F

1

1998	Reference $(n=48)$ Mean $\pm$ SD	Preterm (n=70) Mean $\pm$ SD
Body weight (kg)	$42.09 \pm 9.82$	$42.16 \pm 11.80$
Body height (cm)	$149.84 {\pm} 9.56$	$149.31 \pm 9.47$
2015	Reference (n=29) Mean $\pm$ SD	Preterm (n=13) Mean $\pm$ SD
Body weight (kg)	$62.7 \pm 9.08$	$60.08 \pm 14.66$
Body height (cm)	$167.8 {\pm} 5.6$	$163.00{\pm}10.08$
BMI	$22.23 \pm 2.48$	$22.55 \pm 4.53$

Figure 2: Table 1 :

Figure 3: Table 2 :

 $<sup>^{1}</sup>$ © 2017 Global Journals Inc. (US)

<sup>&</sup>lt;sup>2</sup>Stress Test Performance by Preterm-Born Females in Adolescence and Adulthood © 2017 Global Journals Inc. (US) <sup>3</sup>Stress Test Performance by Preterm-Born Females in Adolescence and Adulthood © 2017 Global Journals

<sup>&</sup>lt;sup>o</sup>Stress Test Performance by Preterm-Born Females in Adolescence and Adulthood © 2017 Global Journals Inc. (US)V.

# 255 .1 Funding

- 256 This study was supported by the Polish National Science Centre Project number 2012/07/D/NZ7/03265 Conflict 257 of interest statement
- 258 The authors declare no conflicts of interest in preparing this article.
- 259 [Peristats et al.], Peristats, March, Dimes. http://www.marchofdimes.com/Peristats/2011
- 260 [] , 10.1371/journal.pone.0080869. 8 p. e80869.
- [Kriemler et al. ()] 'Aerobic and lung performance in premature children with and without chronic lung disease of prematurity' S Kriemler, H Keller, S Saigal, O Bar-Or. 16162994. *Clin J Sport Med* 2005. 15 (5) p. .

[Clemm et al. ()] 'Aerobic Capacity and Exercise Performance in Young People Born Extremely Preterm'. H
 Clemm, O Roksund, E Thorsen, G E Eide, T Markestad, T Halvorsen. 10.1542/peds.2011-0326. *Pediatrics* 2012. 129 (1) p. .

- 266 [Rogers et al. ()] 'Aerobic capacity, strength, flexibility, and activity level in unimpaired extremely low birth
- weight (or=800 g) survivors at 17 years of age compared with term-born control subjects'. M Rogers , T B
  Fay , M F Whitfield , J Tomlinson , R E Grunau . 10.1542/peds.2004-1603. *Pediatrics* 2005. 116 (1) p. .
- [Saigal and Doyle ()] 'An overview of mortality and sequelae of preterm birth from infancy to adulthood'. S
   Saigal , L W Doyle . 10.1016/S0140-6736(08)60136-1. Lancet 2008. 19 (371) p. .
- [Baraldi and Filippone ()] 'Chronic Lung Disease after Premature Birth'. E Baraldi , M Filippone . doi: 10.1056/
   NEJMra067279. New England Journal of Medicine 2007. 357 (19) p. .
- [Magiera ()] Determining exercise capacity with indirect tests using computer-aided engineering (CAE), A
   Magiera . 2000. 35 p. . (Sport Wyczynowy)
- [Goldenberg et al. ()] 'Epidemiology and causes of preterm birth'. R L Goldenberg , J F Culhane , J D Iams , R
   Romero . 10.1016/S0140-6736(08). Lancet 2008. 371 p. .
- 277 [Clemm et al. ()] 'Exercise capacity after extremely preterm birth. Development from adolescence to adulthood'.
- H Clemm, M Vollsaeter, O D Røksund, G E Eide, T Markestad, T Halvorsen. 10.1513/AnnalsATS.201309311OC. Annals of the American Thoracic Society 2014. 11 (4) p. .
- [Narang et al. ()] 'Gas transfer and pulmonary blood flow at rest and during exercise in adults 21 years after
   preterm birth'. I Narang , A Bush , M Rosenthal . Am J Respir Crit Care Med 2009. 180 p. .
- [Svedenkrans et al. ()] 'Long-Term Impact of Preterm Birth on Exercise Capacity in Healthy Young Men: A
   National Population-Based Cohort Study'. J Svedenkrans, E Henckel, J Kowalski, M Norman, K Bohlin.
   *PLoS One* 2013.
- [Walter et al. ()] 'Low Birth Weight and Respiratory Disease in Adulthood. A Population-based Case-Control
   Study'. E C Walter , J Ehlenbach , D L Hotchkin , J W Chien , T D Koepsell . 10.1164/rccm.200901-0046OC.
   *American Journal of Respiratory and Critical Care Medicine* 2009. 180 (2) p. .
- [Vrijlandt et al. ()] 'Lung Function and Exercise Capacity in Young Adults Born Prematurely'. E J Vrijlandt ,
   J Gerritsen , H M Boezen , R G Grevink , E J Duiverman . doi: 10.1164/ rccm.200507-1140OC. American
   Journal Respiratory and Critical Care Medicine 2006. 173 (8) p. .
- [Hulskamp et al. ()] 'Lung function tests in neonates and infants with chronic lung disease of infancy: functional
   residual capacity'. G Hulskamp , J J Pillow , J Dinger , J Stocks . 10.1002/ppul.20318. *Pediatr Pulmonol* 2006. 41 (1) p. .
- [Pillow et al. ()] 'Lung function tests in neonates and infants with chronic lung disease: global and regional
  ventilation in homogeneity'. J J Pillow , I Frerichs , J Stocks . 10.1002/ppul.20319. Pediatr Pulmonol 2006.
  41 (2) p. .
- [Gappa et al. ()] 'Lung function tests in neonates and infants with chronic lung disease: lung and chest-wall
   mechanics'. M Gappa , J J Pillow , Allen J Mayer , O Stocks , J . 10.1002/ppul.20380. Pediatr Pulmonol
   2006. 41 (4) p. .
- [Baldwin et al. ()] 'Lungfunction tests in neonates and infants with chronic lung disease: tidal breathing and
   respiratory control'. D N Baldwin , J J Pillow , J Stocks , U Frey . 10.1002/ppul.20400. Pediatr Pulmonol
   2006. 41 (5) p. .
- [De Kleine et al. ()] Outcome of perinatal care for very preterm infants at 5 years of age: a comparison between *1983 and 1993. Paediatric and Perinatal Epidemiology*, M J De Kleine, A L Den Ouden, L A Kollée, A
  Van Baar, M W Nijhuis-Van Der Sanden, A Ilsen. 10.1111/j.1365-3016.2007.00781.x. 2007. 21 p. .
- [Kotecha and Allen ()] 'Oxygen therapy for infants with chronic lung disease'. S Kotecha , J Allen .
   10.1136/fn.87.1.F11. Arch Dis Child Fetal Neonatal Ed 2002. 87 p. .
- [Tsimis et al. ()] 'Prematurity: present and future'. M E Tsimis , Abu Al-Hamayel , A Germaine , H .
   PMC4323881. Minerva Ginecologica 2015. 67 (1) p. .

- [Lewandowski et al. ()] 'Preterm heart in adult life: cardiovascular magnetic resonance reveals distinct differences
   in left ventricular mass, geometry, and function'. A J Lewandowski , D Augustine , P Lamata , E F Davis ,
- M Lazdam , J Francis . *Circulation* 2013.
- [Kilbride et al. ()] 'Pulmonary function and exercise capacity for ELBW survivors in preadolescence: effect of
   neonatal chronic lung disease'. H W Kilbride , M C Gelatt , R J Sabath . 10.1067/S0022-3476(03)00413-X. J
   *Pediatr* 2003. 143 p. .
- <sup>316</sup> [Welsh et al. ()] 'The EPICure study: maximal exercise and physical activity in school children born extremely
- preterm'. L Welsh , J Kirkby , S Lum , D Odendaal , N Marlow , G Derrick . 10.1136/thx.2008.107474.
   Thorax 2010. 65 p. .
- Beck et al. ()] 'The worldwide incidence of preterm birth: a systematic review of maternal mortality and morbidity'. S Beck , D Wojdyla , L Say , A P Betran , M Merialdi , J H Requejo . 10.2471/BLT.08.062554.
- $_{\rm 321}$   $\qquad Bull World Health Organ 2010. 8 (1) p. .$
- [Tibboel and Jobe ()] 'Update in Pediatric Lung Disease'. D Tibboel , A H Jobe . 10.1164/rccm.201001-0117UP.
   American Journal of Respiratory and Critical Care Medicine 2009. 2010. 181 p. .
- 324 [Magiera et al. ()] 'Zmiany tolerancji wysi?ku u wcze?niaków w okresie pokwitania'. A Magiera , A Ronikier , L
- 325 Kluba , P Zap?atka . Post?py Rehabilitacji 2000. XIV (1) p. .