

Dental Caries Experience in 2-15-Year-Olds Living with HIV in Nairobi and Mombasa "Kenya"

Mariam A. Hussein¹, Amani Wanjala² and Gladys N. Opin³

¹ University of Nairobi

Received: 9 December 2016 Accepted: 31 December 2016 Published: 15 January 2017

Abstract

Title of the article: Dental caries experience in 2-15 year old children living with HIV in Nairobi and Mombasa "Kenya". Background: Dental caries affects children both healthy and those living with HIV. Aims: To determine the dental caries prevalence and association with dental parameters, HIV related oral lesions, oral health practices, immune status and use of ARV's. Methods: This was a cross sectional descriptive study with a sample size of 237 children aged 2-15 years. A specially structured questionnaire captured demographic data, oral health practices, oral complaints. Plaque scores, gingivitis and dental caries were recorded 'in a modified WHO form. Data on ARV use and immune suppression state were obtained from the medical records. Data was analysed using the SPSS software version 17. Results: 237 children confirmed with HIV aged 2-15 years (mean age=7.5 years) were assessed. The deciduous dentition caries prevalence was 84.4

Index terms— children HIV, caries experience prevalence, dmft, DMFT.

1 Introduction

here is minimal literature on the oral health of children with HIV including both dental parameters and oral lesions within the country and our neighbours despite more than 90% of HIV infected children living in Sub Saharan Africa 1,2 . Elsewhere, although there are many studies evaluating oral soft tissue manifestations of HIV/AIDS in children, there are relatively few clinical investigations into the prevalence and incidence of dental caries in the primary and permanent dentitions. [3][4][5] Some authorities in pediatric HIV/AIDS have indicated that there is no substantial difference between pediatric HIV/AIDS children and the general child population; while others have noted increased caries susceptibility. Among the studies conducted, a wide variation in the prevalence of dental caries has been noted. [6][7][8][9][10][11][12] The high prevalence has been associated with nutritional supplements, Cariogenic medications, xerostomia, presence of painful oral lesions. 6,7 II.

2 Materials and Methods

The study was conducted in two private homes for children living with HIV (Nyumbani and New life childrens homes) and two referral hospital outpatient comprehensive care centers (Kenyatta National Hospital and Coast Province General hospital) running HIV programmes for children, located in Nairobi and Mombasa respectively. Since this was a descriptive cross sectional study the sample size was determined using the formula: $n = \frac{Z^2 \cdot P(1-P)}{d^2}$ with P at 85% as the prevalence of gingivitis in HIV infected children 7 CI-Confidence Interval 95%; d-absolute precision-5% [?]; Z 1-? standard error of mean: 1.96 A minimum of 237 children were recruited via convenience sampling technique.

The study was conducted over a period of three months and approved by the joint ethical research committee of KNH and University of Nairobi.

The subjects were categorized into three baseline age groups: 2-5 year olds, 6-11 year olds, and 12-15 year olds based on the dentition primary, mixed and permanent dentition stages respectively.

3 RESULTS

Plaque scores (WHO Plaque index 1999), Gingival scores (Loe and Sillness 1963), Dental caries status (Klein, Palmer and Knutson 1938) and specific HIV related oral manifestations were recorded in a modified WHO Form 13 A specially structured questionnaire was used to assess the oral health practices of each child, complaints of oral pain and challenges in feeding and maintaining oral hygiene. Data on ARV'S and the immune suppression state based on the CD4counts/% were obtained from the records. Data was analysed using S.P.S.S version 17.0 T III.

3 Results

Among the 237children, Six (2.5%) were from NewLife childrens home, 77(32.5%) Nyumbani childrens home, 39(16.5%) from the Comprehensive Care Centre at KNH while 115(48.5%) were from the C.C.C at the CPGH. Children from the homes comprised a total of 83(35%) while those from the outpatient centres completed the remaining study population (154, 65%) as depicted (Fig1).

Among the population examined 112 (47.3%) were males and 115(52.7%) females.

The mean and modal ages were 7.5 and 9 years respectively and age cohorts as shown (Table 1). The mean dmft was 6.38+ SD 5.45, 84.4% had caries while only 15.6% were caries free, 2% had fillings and 7.5% of the children had teeth missing due to caries. The mean DMFT was 3.35 + SD3.55, 78.3% had decayed teeth, 3.9% missing teeth and only 6.2% had fillings (Table 2).

The mean dmft /DMFT of children from the homes ($4.77 \pm SD 4.10$ / $2.95 \pm SD 3.51$) was significantly lower than that of the C.C.C outpatient centres ($7.39 \pm SD 6.04$ / $3.67 \pm SD 3.57$): Tables 3 and 4.

Males had a dmft of 6.49+SD 5.84 and DMFT of 3.03+

The mean dmft of children in the 2-5 yrs , 6-11 yrs and 12-15 year age groups were 7.12 +SD 6.79, 6.57+SD 4.88 and 3.4 ± 1.844 respectively, though not statistically significant.(Table 3). SD 3.13 while females had a dmft of 6.55+SD 5.39 and DMFT of 3.65+SD3.89.The gender differences were not statistically significant: Tables 3 and 4.

Differences in the mean DMFT scores in the 6-11 year ($2.41 \pm SD 2.944$) and 12-15 year ($4.98 \pm SD 3.964$) age groups were found to be statistically significant (Table 4).

Children who reported not brushing (32) had mean dmft of 6.66 ± 6.225 , while those who brushed once (90) had a mean dmft score of 7.84+SD 5.98 and those who brushed twice regularly (77) had the lowest mean dmft of 4.92+SD 4.4. (Table ?? and 6).

Children who complained of pain in the mouth had higher dmft and DMFT's (7.90 ± 6.08 and 4.29 ± 4.21) than those who did not (dmft 5.53 ± 5.04 and DMFT 2.97 ± 3.00) Tables ?? and 6.

Higher dmft and DMFT scores (dmft $8.05 \pm SD 6.03$, DMFT $4.3 \pm SD 4.26$) were recorded in children who reported difficulty in maintaining oral hygiene as opposed to those who did not (dmft $5.45 \pm SD 5.04$, DMFT $2.82 \pm SD 2.97$); Tables ?? and 6.

A statistically significant difference was also noted in the caries status of children who reported of difficulty in feeding due to pain in the oral soft tissues and the teeth and those who did not. Children who did not complain of feeding difficulties had dmft scores of $5.47 \pm SD 5.041$ and DMFT of $2.94 \pm SD 3.099$ whereas children who reported feeding difficulty had dmft scores of $8.05 \pm SD 6.048$ and DMFT $4.13 \pm SD 4.26$. Children who were on ARV therapy (n=109, N=75) had a mean dmft of 7.03 ± 5.742 and DMFT 3.41 ± 3.271 . Children who were not on ARV's (n=90, N=54) had a dmft of 5.91 ± 5.4 and DMFT of 3.28 ± 3.931 respectively. However, statistical tests were not significant as shown below: SD4.182. , Tables ?? and 6.

DMFT: Mann U, (Z=0.483); $p > 0.05$ (0.629). dmft: Mann U(Z=1.385); $p > 0.05$ (0.166) Children with no evidence of immune suppression (n=75, N=50) had a dmft of $5.39 \pm SD 4.426$ and DMFT of $2.22 \pm SD 2.179$ while the group with moderate immunosuppression (n=55, N=41) had a dmft of $4.84 \pm SD 4.574$ and DMFT $4.76 \pm SD 4.939$. Amongst the severely immunosuppressed (n=56, N=30), the mean dmft was $10.04 \pm SD 6.734$ and mean DMFT was $3.4 \pm SD 2.737$ respectively. Statistical tests were highly significant as shown:

DMFT: Kruskal-Wallis, $X^2 = 8.505$; 2df; $p < 0.05$ (0.014) dmft: Kruskal-Wallis, $X^2 = 22.213$; 2df; $p < 0.05$ (0.000) Children with good oral hygiene scores had a lower mean dmft of $4.29 \pm SD 5.22$, and highest dmft scores of $9.43 \pm SD 7.32$ were seen in patients with poor oral hygiene. (Table 7).

Children with good OH scores had a mean DMFT of $3.38 \pm SD 3.18$, those with fair OH scores had a mean DMFT of $3.07 \pm SD 3.29$ while those with poor OH scores presented with the highest scores of $4.35 \pm SD 4.55$. However no significant differences were noted, Kruskal-Wallis test $\{X^2 = 0.593$; 2df; $p = 0.101\}$.

Children who were free of gingivitis had mean dmft of $2.46 \pm SD 3.75$ and a mean DMFT of $3.42 \pm SD 4.36$ and those with mild gingivitis presented with a mean dmft of $6.37 \pm SD 4.67$ and mean DMFT of $3.36 \pm SD 3.55$, while those who had moderate gingivitis presented with a mean dmft of $8.09 \pm SD 6.41$ and mean DMFT of $3.33 \pm SD 3.42$ respectively.

Kruskal-Wallis test was statistically significant $\{X^2 = 24.065$; 2df; $p < 0.05$ (0.000)} :Table 7, however, it was not statistically significant in comparison for the mean DMFT scores $\{X^2 = 0.784$; 2df; $p > 0.05$ (0.676) } .

The mean DMFT of children who did not present with any HIV related oral manifestation (82) was $3.1 \pm SD 3.343$, while those with one manifestation (38) had a mean DMFT of $4.08 \pm SD 4.207$, and those with two or more oral lesions (10) had a mean DMFT of $2.7 \pm SD 1.889$, however these differences were not statistically significant, Kruskal-Wallis $\{X^2 = 2.900$; 2df; $p > 0.05$ (0.235)}.

The mean dmft in children who did not present with any HIV related oral manifestations (104) was $5.16 \pm$

SD4.616 and those who showed presence of one manifestation (81) had mean dmft score of $7.33 \pm SD 5.922$, while those with more than one manifestation (14) had dmft scores of $11.31 \pm SD 6.408$. Kruskal-Wallis was significant $\{X^2 = 16.539, 2df; p < 0.05(0.000)\}$.

IV.

4 Discussion

The overall caries experience in both the deciduous and permanent dentitions was high. The mean dmft was $6.38 \pm SD 5.45$ and 84.4% of the children while only 2% had fillings indicating most of the caries were untreated. The mean DMFT was $3.35 \pm SD 3.55$ and 78.3% of the children had caries while only 6.2% had fillings.

Our study reports dmft and DMFT which are 3.5 and 12.4 times respectively higher than those reported for healthy Kenyan children ??4 .

The caries prevalence is higher than other studies which report prevalences of 60% and 63.8% respectively. 6,7 .

The trend toward increased caries susceptibility in HIV infected children noted may be due to several different factors.

Many of the children experience failure to thrive hence more frequent feedings with carbohydrate and sucrose-rich foods may be necessary to maintain body weight. [3][4][5][6][7][8] It has been shown that increased frequency of carbohydrate intake in HIV infection is associated with increased caries prevalence and cariogenic microbes. In addition to the increased frequency of carbohydrates, many of the medications essential for antiretroviral therapy and prophylaxis against opportunistic infections contain relatively high sweetener levels or are sucrose based for compliance purposes. [3][4][5][6][7][8][9][10][11] The fact that these medications need to be taken on a frequent basis increases the child's exposure to cariogenic substances ??5 .

Lack of effective oral hygiene practices and their late commencement, inadequate oral health education, lack of an adequate referral system resulting in poor dental care for these children may also be factors to consider. Ignorance and fear of the serostatus also prevents these children from seeking dental attention in addition to the high cost of dental treatment which inevitably causes the parents/guardians to prioritize other health needs 6,7 .

The adolescent group also had high mean DMFT of 4.92 ± 3.95 ($p = 0.000$), a finding close to what Meless et al. reported (294/407) fifteen-year-olds had a DMFT 4.1 ± 3.0 16 . This may be attributed to the duration of time the permanent teeth were in the oral cavity. During this period the teeth the erupted permanent teeth are exposed to a cariogenic diet, plaque, gingivitis, and microorganisms or cumulative neglected decay.

In the present study, no record was made of the dietary habits of the children and consumption of snacks.

Significant differences were noted in the mean dmft and DMFT scores among children from the homes ($p = 0.006$) and the C.C.C's ($p = 0.033$). This could be attributed to better and supervised oral hygiene practices incorporated in their care .

The mean dmft scores were also lower in children with good and fair oral hygiene ($p = 0.03$) and mild gingivitis ($p = 0.008$); these findings are in agreement with other studies ??0-11. Among the oral health practices, increased frequency of brushing was significantly associated with lower dmft scores ($p = 0.007$) showing that improved oral hygiene could help slow the decay processes, a finding in agreement with other studies [10][11] .

Children with positive complaints of pain and difficulty eating presented with higher dmft scores ($7.90 \pm SD 6.08$) and DMFT scores (4.29 ± 4.21) as opposed to those with no such complaints (dmft: 5.38 ± 5.04 , DMFT 2.97 ± 3.00) once again reflecting on the vicious cycle of oral disease-painful oral lesions-painful dental disease ??8 . Painful gums and pain from oral lesions create a vicious circle where a child who does not brush their teeth because they are in pain, results in plaque accumulation, followed by increased severity of gingivitis and then caries experience resulting in more pain 11, ??1, ??2 . During these periods there is deteriorating health along with the influence of increased use of medications most of which are likely to be sucrose based to combat various opportunistic infections ??2, ??3 .

The mean dmft and DMFT did not vary significantly among children who were on ARV's and those who were not on therapy. These findings are in agreement with other studies which have not associated ARV's with increased caries prevalence. However, it is known that frequent use of medications most of which are sucrose based for compliance purposes may be responsible for the high caries prevalence ??5 .

Further research may be required to establish any association between duration of ARV therapy, the type of medication, its sucrose content and the caries experience.

Both dmft and DMFT scores were notably higher in children with evidence of moderate and severe immunosuppression ($p = 0.000, 0.014$), a finding in agreement with various studies in the literature to assess the relation between caries and CD4 status ??1, 19, .

The mean dmft scores were lower in children with good and fair oral hygiene ($p = 0.03$) and mild gingivitis ($p = 0.008$), a finding in line with, Hicks et al 11 , Riberio et al 12 and Chen et al ??0 .

High dmft scores were noted among children with HIV related oral manifestations ($p = 0.000$). Reports have cited it as the next common manifestations following candidiasis and concluded an increase in caries prevalence among the pediatric HIV population 2,3,4 .(D D D D) K

Chen et al reported fungal infections were associated with an increased caries rate. ??0 The high prevalence of dental caries observed in the children in this study may be associated with increased levels of streptococcus mutans in the saliva of children living with HIV ??1. Children suffering from HIV infections have been reported

to have xerostomia. Combined with pain, on chewing food salivary flow would be reduced hence the cleansing lubricating effect from the saliva reduces resulting in dryness of the oral cavity leading to increased severity of gingivitis and decay. ??1 .

Low et al 22 studied the effect of severe caries on the quality of life in children .Tooth-associated pain, difficulty with nutrition, hampered growth as noted by body height and weight, and sleep patterns are adversely altered by severe dental caries in children. HIV infected children do have growth retardation and failure to thrive. This may be further complicated by the presence of severe dental caries.

V.

5 Conclusion

This study had a high prevalence of dental caries in both primary and permanent dentition with high mean dmft /DMFT scores. The oral health of HIVinfected children and a reduction in the caries experience are quite important. In pediatric HIV patients, many factors play a role in caries development and attention should be directed toward establishment of a caries prevention regimen.

It is advisable for oral health care to be integrated in the primary care for these children in order to alleviate their pain and improve their nutritional status and quality of life.

6 Challenges and limitations of the study

This study was institution based and conducted over a short period. Hence, the results may not be a representative of the caries experience and prevalence of children living with HIV in Kenya as a whole.

Without the aid of radiographs, only 35% of the decay may have been reported.

7 Volume XVII Issue VII Version I

1

Year 2017

Figure 1: Table 1 :

2

Descriptive	Mean	SD	%
Decayed deciduous	6.38	5.45	84.4%
Missing deciduos	0.15	0.59	7.5%
Filled deciduous	0.05	0.289	2%
Dmft (n=199)	6.38	5.45	
Decayed perm	3.19	3.44	78.3%
Missing perm	0.05	0.29	3.9%
Filled perm	0.16	0.69	6.2%
DMFT (n=130)	3.35	3.55	

Figure 2: Table 2 :

1 2

¹Volume XVII Issue VII Version I © 2017 Global Journals Inc. (US) Year 2017

²© 2017 Global Journals Inc. (US)

3

Variable		N	dmft $\pm$ SD		Statistical test
Study Center					
?	Children homes	66	4.77	4.10	Mann U(Z-.761);p=0.006
?	Outpatient centre	133	7.39	6.04	
Age					
?	2-5yrs	76	7.12	6.79	Kruskal-Wallis Chi-square 4.502;2df;p=0.105
?	6-11yrs	107	6.57	4.88	
?	12-15yrs	15	3.40	1.84	
Gender					
?	Male	99	6.49	5.84	Mann U(Z-.394);p=0.693
?	Female	100	6.55	5.39	

[Note: dmft d-decayed m-missing f-filled in deciduous dentition; $\pm$ -plus or minus, SD-standard deviation © 2017 Global Journals Inc. (US) K]

Figure 3: Table 3 :

4

Variable		N	DMFT $\pm$ SD		Statistical test
Study Center					
?	Children homes	57	2.95	3.51	Mann U (Z-2.132) P=0.033
?	Outpatient Centres	73	3.67	3.57	

Figure 4: Table 4 :

7

Variable		N	dmft $\pm$ SD		Statistical Test
Oral Hygiene					
?	Good	24	4.29	5.22	Kruskal-Wallis Chi-square 11.382;2df;P=0.03
?	Fair	131	5.95	4.60	
?	Poor	44	9.43	7.32	
Gingivitis					
?	No	30	2.46	3.75	Kruskal-Wallis Chi-square 24.065;2df;P=0.000
?	Mild	99	6.37	4.67	
?	Moderate	77	8.09	6.41	

Figure 5: Table 7 :

.1 Acknowledgements

We are grateful to the administrations of New Life and Nyumbani Childrens homes, Comprehensive Care Centers at the Kenyatta National Hospital and Coast Province General Hospital for granting us permission to conduct this study. We appreciate the contribution of Dr Abdul Halim Hussein during the research process, support from staff members Department of Pediatric Dentistry and Orthodontics and School Of Dental Sciences during research presentations and Miss Alice Lakati for assisting with the biostatistics. Our gratitude to our family members for their encouragement.

This study had Ethical clearance of the Kenyatta National Hospital Ethical Research Committee(Ref no: KNH-ERC/01/2407).

[Gelbier et al. ()] ‘A preliminary investigation of dental disease in children with HIV infection’. Marylin Gelbier , . S Victoria , Nike E Lucas , Zerou , J Graham , Vas Roberts , Novelli . *International Journal of Pediatric Dentistry* 2000. 10 p. .

[Fine et al. ()] ‘Clinical applications of the oral manifestations of HIV infection in children’. D H Fine , N Tofsky , E M Nelson , D Schoen , A Barasch . *Dent Clin North Am* 2003. 47 p. .

[Tofsky et al. ()] ‘Dental caries in HIV-infected children versus household peers: 2-year findings’. N Tofsky , E N Nelson , R N Lopez , F A Catalanotto , D H Fine , R V Katz . *Paediatr Dent* 2000. (22) p. .

[Hicks et al. ()] ‘Dental Caries in HIV-infected children: a longitudinal study’. M J Hicks , C M Flaitz , A B Carter , S G Cron , S N Rossmann , S L Simon , G J Demmler , M W Kline . *Paediatr Dent* 2000. (22) p. .

[Eldrige and Gallagher ()] ‘Dental caries prevalence and dental health behaviour in HIV infected children’. K Eldrige , J E Gallagher . *Int J of Paediatr Dent* 2000. 10 p. .

[Rovaris et al. ()] ‘Oral Health Condition of Children Living with HIV Dent’. N S Rovaris , L G Martins , J Traebert . *J* 2014. 2 p. .

[Pongsiriwet et al. (200313)] ‘Oral Lesions and Dental caries status in perinatally HIV-infected children in Northern Thailand’. S Pongsiriwet , A Iamaroon , S Kanjanavanit , K Pattanaporn , Krisana Prakornkit , S . *Int. J Paediatr Dent* 200313. p. .

[Shetty (2006)] ‘Oral lesions commonly associated with pediatric HIV infection-presentation, management and review of literature’. Kishore Shetty . *General Dentistry* 2006. July-August: 284-287.

[Ramos-Gomez et al. ()] ‘Oral manifestations and dental status in pediatric HIV infection’. F J Ramos-Gomez , A Petru , HiltonJ F , CancholaA J , Wara D Greenspan , JS . *International Journal of Pediatric Dentistry* 2000. (10) p. .

[Rwenyonyi et al. (2011)] ‘Oral manifestations in HIV/AIDS-infected children’. C M Rwenyonyi , A Kutesa , L Muwazi , I Okullo , A Kasangaki , A Kekitinwa . *Eur J Dent* 2011 Jul. 5 (3) p. .

[Naidoo and Chikte ()] ‘Orofacial manifestations in pediatric HIV:a comparative study of institutionalized and hospital outpatients’. S Naidoo , U Chikte . *Oral Dis*2004. 10 p. .

[Riberio Ade et al. ()] ‘Relation between Biofilm, caries activity and gingivitis in HIV+ children’. A Riberio Ade , M Portella , I P Souza . *Pesqui Odontol Bras* 2002. 16 p. .

[WHO Oral health Surveys, Basic methods ()] *WHO Oral health Surveys, Basic methods*, 1997. (th Edition: 22-52)