

Nutritional and Microbial Evaluation of Commercial Apple Juices Available in Market of Peshawar City

Sana Irshad Khan¹, Zafar Iqbal² and Ubairah³

¹ Agriculture Research Institute Tarnab, Peshawar-Pakistan

Received: 8 December 2016 Accepted: 4 January 2017 Published: 15 January 2017

Abstract

The present study was aimed to carry out quality assessment of different commercial apple juices available in the market of Peshawar city. Physiochemical characteristics like moisture, total soluble solids (Brix°), pH, acidity (

Index terms— apple juices, quality assessment, SO₂, physico-chemical analysis, WHO standard.

1 Introduction

Apples, a highly nutritious fruit, found abundant of essential food elements including fats (11%), carbohydrates (14.9%), sugars (11%), proteins (0.4%) and balanced level of pectin, dietary fibers and potassium, vitamins (A, B and C) and different types of phenolics (1). Apple can be used raw or processed into juices, jellies, jams, cider and wine etc. (2). Apple juices act as nutritious beverages and are becoming an important part of the modern diet in many communities, essentially available in the same form almost anywhere in the world (3). Several epidemiological studies suggested the antioxidant and detoxification effect of apple juices in the human body and thus playing role to reduce the risk of chronic degenerative diseases (4,5,6).

Quality of fruit juices is strictly maintained in developed countries under some law and regulation but in many developing and under developed countries the manufacturer is not concerned about the microbiological safety, hygiene and nutritional importance of fruit juices because of unawareness and lack of legislation (3). Thus the transmission of some human diseases through juice and other drinks are considered a serious problem in these countries today (7).

From our local market survey, it was revealed that although a large number of fruit juices brands (bottles and tetra packs) are available. Some of these juice brands have been found nutritionally low in quality and synthetic. According FDA (2001) reports fruit juices contain water, sugar and natural fruit pulp that could support microbial growth. Several factors encourage, prevent or limit the growth of microorganisms in juices; the most important are pH, hygienic practice storage temperature and concentration of preservatives and water. Industries apply chemical preservatives (including Sulphur Dioxide (SO₂) and benzoate) that can inhibit all types of microbial growth (8). However these preservatives can significantly damage the vegetative cells also. In order to develop awareness among the people about commercial fruit juices nutritional quality and health hazard due to microbial contamination, this study was attempted to measure nutritional and microbiological quality and SO₂ level of industrially processed apple juices available in the local market of Peshawar city.

2 II.

3 Materials and Methods

4 a) Sample Collection

Different commercially available apple juice samples (Two multinational and four national) were collected from the local market of Peshawar city and were labeled with laboratory code Nos. i.e. I (multinational), II (national), III (national), IV (multinational), V (national) & VI (national) respectively. RTS apple juices were analyzed for the following parameters.

5 b) Physico-Chemical Analysis

Moisture contents of the commercial apple juice samples were determined by direct heating method (9). Total soluble solids of the commercial apple juice samples were recorded by digital Refractometer (Atago Rx-1000) and results were expressed as soluble solids in °Brix (10). Acidity was estimated by titrating diluted samples (1%) against 0.1% NaOH according to the method as described in AOAC (2012) method no. 94.15 (11). The pH was recorded by pH meter (HANNA, HI 2211, pH/ORP meter) by using standard method of AOAC (2012) method no. 2005.02 (12). Reducing and non-reducing sugar (%) were investigated by Lyan and Eynon methods as reported in AOAC (2012) method no. 920.183 & method no. 920.184 respectively (13). Ascorbic acid content (%) of the commercial juice samples was calculated by indophenols titrimetric method as described in AOAC (2012) (9). Total dissolve salts (TDS) in parts per trillion (ppt) of the juice samples was analyzed by digital TDS meter (HANNA, Dist 2 Hi98302) using standard method of AOAC (2012).

6 c) Sulfur dioxide (SO₂) Residue Determination (g/Kg)

Sulfur dioxide (SO₂) concentration (g/Kg) in apple juices was determined as described by Laboratory manual of agricultural Chemistry, The University of Agriculture, Peshawar (14).

7 d) Microbial Analysis (cfu/mL)

For microbial evaluation of commercial juice samples Total Viable Count was used, by pour plate method (7). Sample (1 mL) was taken from the three dilutions (10⁻¹, 10⁻² and 10⁻³) and was added to Petri dish. Then Plate Count Agar ((PCA) media was added to each Petri dish. After incubation at 35 °C for 48 hours, colonies were counted by colony counter and results were expressed as cfu/mL.

8 e) Statistical Analysis

Statistical analysis of results was carried out using CR design. Means of triplicate reading will be represented as mean ± Std.

9 III.

10 Result and Discussion

11 a) Physico-chemical Analysis

Physico-chemical properties of different commercial apple juices available in the market of Peshawar city were shown in Table 1. Maximum % moisture contents was observed in sample VI (88.25±1.88) and minimum % moisture content (82.67±1.71) was observed in sample III of commercial apple juice. Sample I, II, IV and V have moisture content (%) of 84.2±1.13, 86.01±2.01, 83.91±2.17 and 86.12±2.20 respectively. Moisture content affect shelf stability of food samples significantly. Apple juice sample having higher moisture content (%) have low storage life. Of all the samples, sample III has highest TSS 13.1±0.82 followed by sample I (12.23±0.27). In contrast lowest TSS was found for sample VI (1.17±0.15) followed by sample V (2.26±0.05). TSS for samples II and IV are 8.86±0.06 and 10.03±0.20 respectively. TSS for samples III and I fall within acceptable limit of ready to serve drinks (15). PH values for juice samples I to VI were 3.32±0.08, 3.08±0.01, 3.22±0.11, 3.31±0.10, 3.30±0.05 and 3.39±0.05. From pH of juices samples it was clear that sample VI has highest value of 3.39±0.05 and sample II has lowest pH value of 3.08±0.01. Total acidity (%) of the commercial apple juice samples was found in the range of 0.11-0.83. maximum acidity (%) was found for sample I (0.83±0.05) followed by sample III, II and IV (0.68±0.05), (0.24±0.07) and (0.23±0.03) and minimum acidity (%) was found for sample VI (0.11±0.03) followed by sample V (0.14±0.01) respectively. Vit-C contents of the commercial juice samples were ranged from 14% to 26%. Sample III has maximum Vit-C concentration (%) of 26.01±0.05 followed by samples I, II, IV, V and VI having % Vit-C concentration of 23.50±1.83, 22.00±1.02, 20.20±0.33, 20.00±0.02 and 14.00±0.10. Total dissolve solids (TDS) of the commercial apple juice samples (I-VI) were recorded in the range of 0.84 to 0.23. highest TDS was found for sample I (0.84±0.03) and lowest for sample II 0.23±0.01. Recorded TDS for samples III to VI was 0.67±0.11, 0.48±0.08, 0.56±0.10 and 0.48±0.22. Reducing sugar (%) was found highest in sample I (21.45±0.93) followed by sample III (20.20±0.51), IV (18.3±0.43) and II (8.46±0.55). in contrast, in juice samples V and VI no content of reducing sugar was found. Like reducing sugar (%), non-reducing sugar (%) contents were also found 0.00 in sample V and VI. Sample I contained highest concentration (2.1±0.74) of non-reducing sugar followed by sample III (1.81±0.15). While sample II and IV has 0.26±0.06% and 0.93±0.20 % of non reducing sugars.

12 b) Sulfer Dioxixe (SO₂) residue determination (g/Kg)

Sulfur dioxide concentration of juice samples (I to VI) were 0.01±0.02, 0.10±0.00, 0.01±0.01, 0.05±0.00, 0.10±0.02 and 0.17±0.09. According to WHO standard, MRL (maximum Residual Limit) for SO₂ in juices is 0.03 g/Kg. Of all the selected juice samples, only samples I (multinational) and III (national) showed SO₂ residues concentration less than 0.03 g/Kg (i.e. 0.01±0.02 and 0.01±0.00 respectively) and were found suitable for human consumption.

97 All other samples were proven toxic and need to neglect its consumption (Table 1). Consumption of SO₂ above
 98 standard limits

99 13 Conclusion

100 From this study it was observed that sample III (national) followed by sample I (multinational) of the selected
 101 apple juices is more suitable for human consumption. In contrast, human usage of samples VI and V is not good
 102 physic-chemically. Similarly all samples were found acceptable for human's consumption from microbial point of
 103 view and fall in the range of WHO standards.

104 14 Parameters

1

Parameters	Sample I	Sample II	SampleIII	Sample IV	Sample V	Sample VI
Moisture %	84.2±1.13	86.01±2.01	82.67±1.71	83.91±2.17	86.12±2.20	88.25±1.88
TSS Brix°	12.23±0.27	8.86±0.06	13.1±0.82	10.03±0.20	2.26±0.05	1.17±0.15
pH 32.4 °C	3.32±0.08	3.08±0.01	3.22±0.11	3.31±0.10	3.30±0.05	3.39±0.05
Acidity %	0.83±0.05	0.24±0.07	0.68±0.05	0.23±0.03	0.14±0.01	0.11±0.03
Vit-C %	23.50±1.83	22.00±1.02	26.01±0.05	20.20±0.33	20.00±0.02	14.00±0.10
TDS ppt	0.84±0.03	0.23±0.01	0.67±0.11	0.48±0.08	0.56±0.10	0.48±0.22
Red.sugar%	21.45±0.93	8.46±0.55	20.20±0.51	18.3±0.43	0.00±0.00	0.00±0.00
Non-reducing sugar %	2.1±0.74	0.26±0.06	1.81±0.15	0.93±0.20	0.00±0.00	0.00±0.00
SO ₂ (g/Kg)	0.01±0.02	0.10±0.00	0.01±0.01	0.05±0.00	0.10±0.02	0.17±0.09
c) Microbial Analy- sis						
Total viable count (CFU/mL) concentration was 30, 21						

Figure 1: Table 1 :

2

Figure 2: Table 2 :

105 1

-
- [Kurowska et al.] , E M Kurowska , J D Spence , J Jordan , S Wetmore , D J Freeman , L A Piché , & P Hdl .
 [Baron and Drilleau ()] , R C Baron , G S Drilleau . *Int. J. Biol. Macromol* 2001. 29 p. .
 [Deanna and Jeffrey ()] ‘Acid-alkaline balance: role in chronic disease and detoxification’. M M S B Deanna ,
 Jeffrey . *Alternative Therapies* 2007. 13 (4) p. 65.
 [AOAC (Association of official and analytical chemist) ()] *AOAC (Association of official and analytical chemist)*,
 2012. (Official method of analysis. 1st edition USA. Method no. 932.12)
 [AOAC (Association of official and analytical chemist) ()] *AOAC (Association of official and analytical chemist)*,
 2012. (Official method of analysis. 1st edition USA. Method no. 2005.02)
 [AOAC (Association of official and analytical chemist) ()] *AOAC (Association of official and analytical chemist)*,
 2012. (Official method of analysis. 1st edition USA. method no. 920.183 & method no. 920.184)
 [AOAC (Association of official and analytical chemist). 2012. Official ,method of analysis] *AOAC (Association
 of official and analytical chemist). 2012. Official ,method of analysis*, (1 st edition USA. method no. 942.15)
 [Franke et al. ()] ‘Bioavailability and antioxidant effects of orange juice components in humans’. A A Franke , R
 V Cooney , S M Henning , L J Custer . *J Agric Food Chem* 2005. 53 (13) p. 5170.
 [Serratore ()] ‘Cholesterolraising effect of orange juice in subjects with hypercholesterolemia’. Serratore . *Am J
 Clin Nutr* 2000. 72 p. .
 [Doyle et al. ()] *Food Microbiology*, M P Doyle , L R J Beuchat & T , Montville . 2001. Washington DC: ASM
 Press.
 [IPCS -International Programme on Chemical Safety World Health Organization ()] ‘IPCS -International Pro-
 gramme on Chemical Safety’. *World Health Organization* 1999.
 [Shah et al. ()] ‘Quality evaluation and preparation of apple and olive fruit blended jam’. W Shah , A Khan , A
 Zeb , M A Khan , F N Shah , N U Amin , M Ayub , S Wahab , A Muhammad , S H Khan . *Global Journal
 of Medical Research:L Nutrition and food Science* 2015. (1) p. 15.
 [Akhtar et al. ()] ‘Quality evaluation of different brands of grapes juice available in local markets of Peshawar
 city’. S J Akhtar , S Ali , S Hassan , B Abbas , F A Javid , M Khan , Siddique . *International Journal of
 Basic Medical Sciences and Pharmacy* 2014. 4 (1) p. .
 [Hossain et al. ()] ‘Quality of Industrially Processed Fruit Juices: An Assessment Using Multivariate Framework.
 Dhaka Univ’. A Hossain , M Rahman , Z Rahman Shabuz . *J. Sci* 2012. 60 (2) p. .
 [Cindric et al. ()] ‘Sample Preparation Methods for the Determination of the Antioxidative Capacity of Apple
 Juices’. I Cindric , J , M Kun?tic , M Zeiner , G Stingeder , G Rusak . *Croat. Chem. Acta* 2011. 84 (3) p. .
 [Bennett and Donovan ()] ‘The determination of sulphur dioxide in citrus juices’. A H Bennett , F K Donovan .
Analyst 1943. 68 p. .