



GLOBAL JOURNAL OF MEDICAL RESEARCH: B
PHARMA, DRUG DISCOVERY, TOXICOLOGY & MEDICINE
Volume 18 Issue 1 Version 1.0 Year 2018
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Studies on Majoon Aarad Khurma and its Granules Prepared with Stevia, with Reference to the Standardization and Toxicity Evaluation

By Mateen Ahmad Khan, Yasmeen Ansari, Roohi Zaman & Izharul Hasan

AIUMC Muzaffarnagar

Abstract- There is a huge treasure of Compound drugs described in various pharmacopoeias that have developed as a result of painstaking and cumulative efforts of elite scholars of Unani medicine. However, there has always been scope for inclusion of new compound drugs whose safety and efficacy has been proved scientifically.

Majoon Aarad Khurma which is widely used as an effective aphrodisiac is prepared with sugar as base. It is contraindicated in diabetic patients who are suffering from sexual dysfunctions. Preparation of medicines which are sugar free should be innovated or designed to meet the demand of the diabetic patients. Therefore the present study is aimed to develop granules of Majoon Aarad Khurma with natural sweetening agent *Stevia rebaudiana*. Granules are more convenient and comfortable in usage and dispensing. Granules uphold the same principles and maintain the same characteristics as traditional dosage forms, granules are safe, light, efficacious, stable and quality controlled.

Keywords: aphrodisiac, pharmacopoeias, diabetes, sexual dysfunction, granules, stevia rebaudiana, ccrum, sweetening agent.

GJMR-B Classification: NLMC Code: QV 4



Strictly as per the compliance and regulations of:



Studies on Majoon Aarad Khurma and its Granules Prepared with Stevia, with Reference to the Standardization and Toxicity Evaluation

Mateen Ahmad Khan ^α, Yasmeen Ansari ^σ, Roohi Zaman ^ρ & Izharul Hasan ^ω

Abstract- There is a huge treasure of Compound drugs described in various pharmacopoeias that have developed as a result of painstaking and cumulative efforts of elite scholars of Unani medicine. However, there has always been scope for inclusion of new compound drugs whose safety and efficacy has been proved scientifically.

Majoon Aarad Khurma which is widely used as an effective aphrodisiac is prepared with sugar as base. It is contraindicated in diabetic patients who are suffering from sexual dysfunctions. Preparation of medicines which are sugar free should be innovated or designed to meet the demand of the diabetic patients. Therefore the present study is aimed to develop granules of Majoon Aarad Khurma with natural sweetening agent *Stevia rebaudiana*. Granules are more convenient and comfortable in usage and dispensing. Granules uphold the same principles and maintain the same characteristics as traditional dosage forms, granules are safe, light, efficacious, stable and quality controlled.

In present study an important Unani formulation i.e. Majoon Aarad Khurma has been modified into granules using *Stevia rebaudiana* as a sweetener and the safety and toxicity of the granules of Majoon Aarad Khurma has also been evaluated. Majoon Aarad Khurma and Granules of Majoon Aarad Khurma were prepared and subjected to Physico-Chemical evaluation with reference to the standards mentioned in Physicochemical Standards of Unani formulations by CCRUM.

Keywords: aphrodisiac, pharmacopoeias, diabetes, sexual dysfunction, granules, stevia rebaudiana, ccrum, sweetening agent.

1. INTRODUCTION

According to the Unani system of medicine, the health is a state of body in which there is equilibrium in humors and functions of the body. To maintain the correct humoral balance there is a power of self preservation called "Quwwate Mudabbirahe Badan" (Immunity of body) in the body.

Therefore the aim of the Unani physician is to find out the cause of the underlying disruption of humors, so that it can be corrected and disease can be cured. The temperament of the person is identified and

diet/medicine/other recommendations are made that are most suitable for achieving and maintaining health of the particular person.

Sexual function is an important component of quality of life and subjective well being of humans. Human sexuality is a multidimensional phenomenon having biological, psychological, behavioral, clinical, moral and cultural aspects. It has been integral part of all cultures since time immemorial. But no single dimension of sexuality is universally dominant. Every person has sexual feelings, attitude and believes, but everyone's experience is unique because it proceeds through an intentionally personal prospective. The cardinal phases in sexual act in male are desire, erection, penetration and orgasm. The phases of sexual act in females are quite different from male. Usually in medicine and also in cultural aspects, sexuality has been mainly concerned with male sexual desire which increase in proportion to the level of secretion of the sex hormones. Sexual response is triggered by both psychological and physical stimuli. Sexual problems are widespread and adversely effects mood, well being, and inter-personal functioning.

Unani medicine treats sexual debility in its own way and proposes different methods of treatment Ilaj bil Ghiza (Dietotherapy), Ilaj bil Dawa (Pharmacotherapy) and Ilaj bit Tadbeer (Regimental therapy). Therapeutic use of the drugs is based on certain principles. The sexual problems are taken up in individualized way taking into account the entire personality of the patient ¹. Unani medicine has holistic approach towards diagnosis and treatment of sexual dysfunction that is not just confined to inability to perform the sex rather includes loss of libido, erectile dysfunction, ejaculatory insufficiency, an orgasmic state, excessive nocturnal emissions and even infertility in males, which may be due to Zoofe Bah (sexual dysfunction) or Nuqse Mani (seminal defects). It also distinguishes between sexual inadequacy and seminal inadequacy ²⁻⁴.

In present study an important Unani compound formulation Majoon Aarad Khurma has been modified into granular form using natural sweetening agent *Stevia rebaudiana* which has sweetening property as well as hypoglycemic activity, the granules of Majoon Aarad Khurma become palatable and will not cause any harm to diabetic patients who are suffering from sexual

Author α: Associate professor, Dept. of Ilmul Saidla, AIUMC Muzaffarnagar. e-mail: saurabh1468@gmail.com

Author σ: Lecturer, Dept. of Ilmul Qabalat-wa-Amraz-e-Niswan, AIUMC Muzaffarnagar.

Author ρ: Professor, Dept. of Ilmul Saidla, NIUM Bangalore.

Author ω: Consultant Unani, AWC, President Estate, Rashtrapati Bhavan, New Delhi.

dysfunction. This study also includes evaluation of physicochemical standards of Majoon Aarad Khurma and granules of Majoon Aarad Khurma and their safety and toxicity study.

II. MATERIAL AND METHODS

According to the protocol of the study "Studies on Majoon Aarad Khurma and its Granules prepared with Stevia, with reference to the Standardization and Toxicity Evaluation" Majoon Aarad Khurma and Granules of Majoon Aarad Khurma was prepared in the laboratory of Dept. of Ilmul Saidla, NIUM.

III. PREPARATION OF MAJOON AARAD KHURMA

All the required ingredients of Majoon Aarad Khurma and Granules of Majoon Aarad Khurma were

Table 1: Ingredients of Majoon Aarad Khurma

Sl. No.	UNANI NAME	BOTANICAL NAME	PART USED	QUANTITY
1	Khurma	<i>Phoenix dactylifera</i>	Fruit	200gm
2	Samagh arbi	<i>Acacia arabica</i>	Gum	200gm
3	Singhara khushk	<i>Trapa bispinosa</i>	Fruit	200gm
4	Satawar	<i>Asparagus rasemosus</i>	Root	50gm
5	Jaiphal	<i>Myristica fragrans</i>	Nutmeg	1.25gm
6	Javitri	<i>Myristica fragrans</i>	Mace	1.25gm
7	Qaranfal	<i>Myrtus caryophyllus</i>	Flower Buds	2.5gm
8	Maghaze Badam	<i>Prunus amygdalus</i>	Fruit	25gm
9	Maghaze Chilghoza	<i>Pinus gerardiana</i>	Fruit	25gm
10	Maghaze Fundaq	<i>Corylus avellana</i>	Fruit	25gm
11	Maghaze Pambadana	<i>Gossypium herbaceum</i>	Fruit	5gm
12	Qand safaid	Sugar	Sugarcane	1kg

The dried raw drugs from 2-7 mentioned in table were powdered in mixer and sieved in (sieve number 80), raw drugs from 8-11 were powdered separately and sieved in (sieve number 40), and dates were separately dried in a hot air oven at 100°C for 4 hours then powdered and passed through (sieve number 60). Qiwan was prepared according to method mentioned in Formulary with 1 kilo sugar and 600 ml water; the dried drugs were mixed one by one in the Qiwan and stirred slowly. Finally all the Maghazyat (Kernels) 8-11 were mixed gradually in the Qiwan. And

procured from the raw drug dealers under the supervision of the Guide, and all the raw drugs were identified and authenticated by the expert Dept. of Ilmul Advia, NIUM Bangalore, (Karnataka).

The Majoon Aarad Khurma was prepared as per the formulation mentioned in the National Formulary of Unani Medicine, Part-1, Govt. of India. The composition of Majoon Aarad Khurma is as given below:

stored in a container at room temperature for further study⁵.

IV. PREPARATION OF GRANULES OF MAJOON AARAD KHURMA

The granule of Majoon Aarad Khurma was prepared as per the formulation mentioned in the National Formulary of Unani Medicine, Part-1, Govt. of India, the composition of granules of Majoon Aarad Khurma is as given below:

Table 2: Ingredients of granules of Majoon Aarad Khurma

Sl. No.	UNANI NAME	BOTANICAL NAME	PART USED	QUANTITY
1	Khurma	<i>Phoenix dactylifera</i>	Fruit	200gm
2	Kamagh arbi	<i>Acacia arabica</i>	Gum	200gm
3	Singhara khushk	<i>Trapa bispinosa</i>	Fruit	200gm
4	Satawar	<i>Asparagus rasemosus</i>	Root	50gm
5	Jaiphal	<i>Myristica fragrans</i>	Nutmeg	1.25gm
6	Javitri	<i>Myristica fragrans</i>	Mace	1.25gm
7	Qaranfal	<i>Myrtus caryophyllus</i>	Fruit	2.5gm
8	Maghaze Badam	<i>Prunus amygdalus</i>	Fruit	25gm
9	Maghaze Chilghoza	<i>Pinus gerardiana</i>	Fruit	25gm
10	Maghaze Fundaq	<i>Corylus avellana</i>	Fruit	25gm
11	Maghaze Pambadana	<i>Gossypium herbaceum</i>	Fruit	5gm
12	Stevia plant powder	<i>Stevia rebaudiana</i>	leaves	3.50gm

All the dried ingredients were powdered and sieved in (sieve number 80). All the Maghaziyat (kernels) were powdered separately and sieved in (sieve number 40), and dates were separately dried in a hot air oven at 100 °C for 4 hours and then powdered and passed through sieve number 60. Stevia plant extract was prepared with 120 ml water at low temperature for 15 minutes, and sieved through muslin cloth, the total quantity of this extract obtained was 80 ml. All the dried drugs were mixed one by one in Stevia extract, and subjected into the granulator (sieve number 20) for formation of granules and then stored in container at room temperature for further study ⁵.

V. ACUTE TOXICITY STUDY

a) Experimental Animal

Swiss mice of both sexes, weighing 25-35 gm were used. The animals were procured from the, Sri Raghvendra Enterprises, Vijayanagar, Bangalore, Karnataka (India).

Prior to the experiment the animals were allowed to acclimatized for at least one week. They were maintained under standard laboratory conditions throughout the experimental period and were provided with standard diet and water *ad libitum* unless stated otherwise. They were housed in clean polypropylene cages at room temperature 25±2°C, humidity at 45-55% with 12 hours light: 12 hours dark cycle. The animal care procedures and experimental protocol were in according with the guidelines of CPCSEA.

b) Extractive Values ⁶

For the determination of extractive values in non-successive of GMAK was carried out in Soxhlet apparatus, with hydro-alcoholic solvents i.e. 50% distilled water and 50% ethanol (1:1) ratio.

c) Methodology for Acute Toxicity Study ^{7, 8 - 15}

Acute toxicity test was performed according to the World Health Organization (WHO) guideline (WHO 2000) and the Organization of Economic Co-operation and Development (OECD) guideline for testing of chemicals 420 (OECD 2001). Swiss mice of either sex

weighing 25-35 gram were randomly assigned to four groups (I, II, III, & IV,) of 7 mice each. Mice were fasted overnight (12 hrs) with free access to water prior to administration of single doses (0.398, 5.73, 9.73, & 16.69 g/kg b.wt.). The extract dissolved in distilled water and administered orally once a day. After the administration of the test drug all the animals were kept in polypropylene cages singly and were observed for Gross behaviour and mortality at 0 min, 30 min, 60 min, 120 min, 240 min and 24 hrs. The Gross behavioural changes such as piloerection, grooming, trembling, wriggling, diarrhoea, breathing difficulty, constant changing position, immobility, asthenia, anorexia, ataxia, urination and syncope were monitored continuously for any above abnormal changes.

VI. PHYSICO-CHEMICAL EVALUATION

The Physico-Chemical studies were carried out on Majoon Aarad Khurma and Granules of Majoon Aarad Khurma in the laboratory of Dept of Ilmu Saidla, NIUM, Bangalore. Majoon Aarad Khurma and Granules of Majoon Aarad Khurma were prepared and subjected to Physico-Chemical evaluation under the following parameters:

(1) Organoleptic properties such as the appearance, colour, smell, and taste (2) Alcohol soluble matter and Water soluble matter (3) Successive extractive values (4) PH value (5) Bulk density and Tapped density (6) Ash value (7) Volatile oil (8) Saponification value (9) Iodine value (10) Acid value (11) Estimation of total Alkaloids (12) Resin (13) Reducing and non-Reducing sugars (14) Crude fibers (15) Thin layer chromatography (TLC) was also conducted for identification of compounds.

VII. RESULTS AND OBSERVATION

Both the test drugs sample Majoon Aarad Khurma and granules of Majoon Aarad Khurma were evaluated for physico-chemical parameters as recommended and almost all the values of both the test drugs were found within the standard limits.

Table 3: Physico-Chemical comparative data of MAK and GMAK

Sl. No	Physico Chemical Properties	MAK	GMAK
1.	Organoleptic Properties Apearance Odour Smell Taste	Semi Solid Brownish Pleasant Sweet	Granules Brownish Pleasant Sweet
2.	Alcohol Soluble Matter	65.5%	24.6%
3.	Water Soluble Matter	46.5%	36.6%
4.	Successive Extractives Petroleum Ether Chloroform Ethyl Alcohol Aqueous	2.4% 0.4% 41.7% 35%	4.2% 0.6% 19.13% 37.2

5.	PH Value 1% 10%	4.90 4.32	5.82 5.27
6.	Ash Value Total Ash Acid Insoluble Ash Water Soluble Ash	0.66% 0.66% 1.33%	2.5% 0.66% 1.16%
7.	Bulk Density Tapped Density Carr's Index Hausner Ratio		0.6gm/ml 0.68gm/ml 12% 1.13
8.	Volatile oils	Traces	0.1%
9.	Saponification Value	680.13	77.07
10.	Iodine Value	4.29	1.14
11.	Acid Value	4.21	2.80
12.	Alkaloids Total	0.13%	3.52%
13.	Resin Estimation	15.1%	37.2%
14.	Determination of Resin	15.2%	37.2%
15.	Reducing sugar	9.2%	15.6%
16.	Non-Reducing sugar	32.11%	24.2%
17.	Crude Fibers	1.13%	2.62%
18.	TLC Rf Values	0.27 0.31 0.50	0.31 0.36 0.50 0.68 0.75

VIII. CHEMICAL EVALUATION

a) Thin Layer Chromatography: ¹⁶⁻²⁹

Thin layer chromatography was carried out on T.L.C. pre coated aluminium plates, silica gel 60 F 254 (layer thickness 0.25 mm) for ethanolic extract of both the test drug samples MAK and GMAK in various mobile phases, later sprayed by different spraying

reagents to visualise the spots. The R_f values of the spots were calculated for both the drugs by the following formula.

$$R_f \text{ Value} = \frac{\text{Distance travelled by Spot}}{\text{Distance travelled by Solvent}}$$

Table 4: TLC of Majoon Aarad Khurma and granules of Majoon Aarad Khurma

Extract	Solvent System	No. of Spots	R_f Value	Colours
MAK Ethanol	Toluene: Ethyl acetate (7:3, with 2 drop Sulphuric acid)	3	0.27	Green
			0.31	Yellow
			0.50	Pink
GMAK Ethanol	Toluene: Ethyl acetate(7:3, with 2 drop Sulphuric acid)	5	0.31	Green
			0.36	Brown
			0.50	Light Pink
			0.68	Pink
			0.75	Yellow




Fig. 1: TLC plates high lighting the spots in MAK and GMAK

b) Acute Toxicity Study

The acute toxicity study was done on Swiss mice of either sex using Hydro-Alcoholic extract of granules of Majoon Aarad Khurma orally, no behavioural changes and mortality was found during 24 hours observation period.

IX. DISCUSSION

Majoon Aarad Khurma is one such popular drug which is widely used as an effective aphrodisiac, which is prepared with sugar as base but as we know that the intake of sugar is not advisable in diabetic patients because the presence of sugar in large amount in blood may develop the complications of diabetes more rapidly so any preparation having sugar as a base or content may create such risk. So even after gaining such popularity as an aphrodisiac, Majoon Aarad Khurma cannot be given to diabetic patients who are suffering from erectile dysfunction. Hence sugar free an alternate formulation should be innovated or designed to meet the demand of the diabetic patients.

X. CONCLUSION

The Physicochemical standards for scientific evaluation of Majoon Aarad Khurma and granules of Majoon Aarad Khurma were estimated and the standards were evaluated as recommended by CCRUM.

Based on the finding it is concluded that

- Granules possessed the same principles and maintained same characteristics as traditional dosage form Majoon Aarad Khurma.
- The granules of Majoon Aarad Khurma were found to be more stable, convenient and comfortable in usage and dispensing, and also safe, light, efficacious, cost effective and quality controlled.
- Stevia a natural sweetening agent which was used as base for granules was evaluated for its toxicity in animal models and no toxicity was found, hence Stevia can be used as safe and efficacious sweetening agent in preparation of granules as well as in other Unani formulations.

REFERENCES RÉFÉRENCES REFERENCIAS

1. McCary JL and Mc Carry SP. *Aphrodisiac and Anaphrodisiacs in Mccary's Human Sexuality*. 4th ed. 1974: 227-33.
2. Basheer A. *Risalae Quwwate Bah*. Kanpur: Munshi Naval Kishore; 1886. pnm.
3. Khan MN. A scientific study of some Unani drugs used for improving the male sexual function. Aligarh- Thesis submitted to Dept. of Ilmul Advia, A.K.T.C, A.M.U; 1993:
4. Moben FAM. *Qanoone mubasharat*. Delhi: Jamia Barqi Press; 1934: pnm.

5. Said HM. *Hamdard Pharmacopoeia of Eastern Medicine*. New Delhi: Sri Sat Guru Publications; 1997: 9-10, 259, 353.
6. Mohammad TH. *Evaluation of lithotriptic activity of Tukhme Karafs (seeds of Apium graveolens Linn) in experimental animals*, Dept. of Ilmul Advia, NIUM, Bangalore, 2010.
7. *Acute and sub-chronic oral toxicity studies of an aqueous stem bark extract of Pterocarpus soyauxii Taub (Papilionaceae) in rodents*; Journal of Ethnopharmacology 133 (2011) 329–335.
8. Irshad S. Temperature standardization and comparative toxicity study of Kushta Sammul Far prepared by different methods, *Thesis*. Dept. of Ilmul Advia, NIUM, Bangalore, 2009.
9. Ibn Baitar. *Aljame al Mufradat al Advia wa al Aghzia*. Vol.1st. Urdu translation by CCRUM New Delhi: Dept of AYUSH, Ministry of H & FW. Govt. of India; 1999: 136, 349, 437.
10. Aulton ME, Wells TI. *Pharmaceutics: The Science of Dosage Form Design*. Churchill Livingstone London, England, 1988. Evaluation of tapioca sago starch as a binder in tablet formulation. *IJPI's Journal of Pharmacognosy and Herbal Formulations* 2010; 1(1).
11. Afaq SH, Tajuddin and Siddiqui MMH. *Standardization of herbal drugs*. Aligarh: Publication AMU Aligarh: 100.
12. Sutharsingh R, Kavimani S, Jayakar B, Uvarani M, Thangathirupathi A. Quantitative phytochemical estimation and antioxidant studies on aerial parts of *Naravelia zeylanica* dc. *International Journal of Pharmaceutical Studies and Research* 2011 April June; II (II): 52-56.
13. Harborne JB. *Phytochemical methods*, London: Chapman and Hall, Ltd. 1973; 49-188.
14. Vogel. *Elementary practice organic chemistry, qualitative organic analysis*. Part 3. Great Britain: Longman Group Limited; 1970: 743.
15. Hertz C, Gerstein. *Evidence-Based Diabetes Care*; BC Decker Inc Hamilton, London: 2001: 531.
16. Siddique AB. et al., *Chemical composition of essential oil by different extraction methods and fatty acid analysis of the leaves of Stevia Rebaudiana Bertoni*. *Arabian Journal of Chemistry* (2012), doi:10.1016/j.arabjc.2012.01.004.
17. Luo, H. et al., Comparison of effectiveness and safety between granules and decoction of Chinese herbal medicine. A systematic review of randomized clinical trials. *J. Ethnopharmacol.* (2012), doi: 10.1016/j.jep.2012.01.031.
18. Fan T.P. et al., Future development of global regulations of Chinese herbal products. *J. Ethnopharmacol.* (2012), doi:10.1016/j.jep.2012.02.029.
19. Niharika Sahoo et al., *Herbal drugs: Standards and regulation*. Department of Biotechnology, West Bengal, India. *Fitoterapia* 2010; 81: 462–471.
20. Jenkins GL, Knevel AM and Digangi FE. *Quantitative Pharmaceutical Chemistry*. 6th ed. New Delhi: CBS Publishers; 2008: 243-253, 379.
21. Goodman and Gilman's. *The Pharmacological Basis of Therapeutics*. 10th ed. U.S.A: Mc Graw-Hill, Medical publishing Division; 2001: 1687.
22. http://wiki.answers.com/Q/What_is_a_non-reducing_sugar [cited Dec. 2011: 15/122011].
23. Shamsi. Standardization and Pharmacological Evaluation of two Nervine Unani Formulations, *Thesis*. Aligarh: Dept. of Ilmul Advia, AMU; 2008: 85.
24. Kinsey et al. 1948 cited by Masters WH, Johnson VE, Kolodny RC. *Masters and Johnson on Sex and human loving*. Mumbai: Jaico publishing house; 2002: 572.
25. Kabiruddin HM. *Al-Akseer*. Vol. 2nd, New Delhi: Ajaz publication House, 2003: 1254-1284.
26. Siddique MSH, Ashraf SM and Siddiqui MMH. *Impotence: Unani concepts and its management*, The Journal of Research and Education in Indian Medicine. 1992: 14-17.
27. Sahib Singh R. *The Effect of Testosterone on the Cavernous Tissue and Erectile function*; World J. Urol 1997; 15, 21.
28. www.diabetes.niddk.nih.gov/dm/pubs/complications.
29. Rance J, Phillips C, Davies S, et al. *How much of a priority is treating erectile dysfunction: a study of patients' perceptions*. *Diabetic Med* 2003; 20: 205-9.