



GLOBAL JOURNAL OF MEDICAL RESEARCH: G
VETERINARY SCIENCE AND VETERINARY MEDICINE
Volume 18 Issue 1 Version 1.0 Year 2018
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Biochemical Identifiers of Postmortem Time Interval on Autopsy of Albino Rats versus Physiological One

By Mahmoud M. El Alfy & El-Hadidy G. Mona

Mansoura University

Abstract- Post mortem time interval is still controversial point as no definitive markers estimate successful time since death even there are many studies on human or experimental models investigate physical, biochemical and molecular marker for identification periods of time elapsed after death. After death, no system or organ work in integrity manner, body loss its immune defence and no elimination of breakdown of each cells which could be explain elevation of certain biochemical in serum of dead animal. In this study, we induced brain stem death induced in albino male rat and take heart blood at different time point for analysis of PH, Ast, Alt, lactate dehydrogenase, creatinine, s-fas and tnfa. It was found that PH reduced while serum lactate dehydrogenase was increased in time dependent manner. Notably, s-fas and tnfa increase in dependent manner until last time point. But AST, ALT, BUN and creatinine shown no change except little increase in last time point. On conclusion group of biochemical could be used in estimation of postmortem time interval for each time point.

Keywords: *post mortem time interval, serum biochemicals, LDH, s-fas and tnfa, albino rats.*

GJMR-G Classification: *NLMC Code: QW 70*



Strictly as per the compliance and regulations of:



Biochemical Identifiers of Postmortem Time Interval on Autopsy of Albino Rats versus Physiological One

Mahmoud M. El Alfy ^α & El-Hadidy G. Mona ^ο

Abstract- Post mortem time interval is still controversial point as no definitive markers estimate successful time since death even there are many studies on human or experimental models investigate physical, biochemical and molecular marker for identification periods of time elapsed after death. After death, no system or organ work in integrity manner, body loss its immune defence and no elimination of breakdown of each cells which could be explain elevation of certain biochemical in serum of dead animal. In this study, we induced brain stem death induced in albino male rat and take heart blood at different time point for analysis of PH, Ast, Alt, lactate dehydrogenase, creatinine, s-fas and tnfa. It was found that PH reduced while serum lactate dehydrogenase was increased in time dependent manner. Notably, s-fas and tnfa increase in dependent manner until last time point. But AST, ALT, BUN and creatinine shown no change except little increase in last time point. On conclusion group of biochemical could be used in estimation of postmortem time interval for each time point.

Keywords: post mortem time interval, serum biochemicals, LDH, s-fas and tnfa, albino rats.

I. INTRODUCTION

The accurate detection of time elapsed since death is one of the most challenges in forensic medicine. there are many methods used to detect the postmortem interval, including physical, chemical and genetic methods (1).

In forensic medicine, the estimation of the time elapsed since death is important due to its role in knowing possible criminal cases and help in apply justice and penalties.

The previously known different elements of the postmortem interval can be measurable from 1 day to many years after death. The determination of the postmortem interval is based on the different changes that a cadaver passes after death including physical, chemical, metabolic processes, bacterial processes and the effect of insect activity (2). The estimating the time since death can be detected quantitatively and qualitatively (3).

Determination of the time since death by chemical means based on systematic patho-

physiological alteration and are found to be more benefit since the effect of external conditions is less (4). The Concentrations of hypoxanthine, NADH, ammonia, and formic acid increased with time and these metabolites may be good markers for post-mortem interval (5). Moreover, in the post-mortem period vitreous humour Na⁺, K⁺ and Ca²⁺ concentration has been investigated for many years. Various authors have found the correlation between increasing these cations and post-mortem interval (6). Additionally, the correlation between total and direct bilirubin, uric acid, urea, transferrin, immunoglobulin M (IgM), creatine kinase (CK), aspartate transaminase (AST), iron and calcium increased significantly with the time of blood putrefaction (7).

The estimation of the time elapsed since death by molecular techniques reported that a time-dependent increase in the mRNA expression of mRNA expression of Fas Ligand (FasL) and phosphatase and tensin homologue deleted on chromosome 10 (PteN) by Quantitative-PCR. A direct linear correlation was found between the mRNA levels of both proteins up until 6 h after death, using a regression analysis (8). Additionally, postmortem serum levels of HMGB1 protein of 90 male Wistar rats preserved at 4, 14 and 24°C since death were estimated by enzyme-linked immunosorbent assay. The serum hmGB1 level showed a time-dependent increase. These observation provide that HMGB1 is related to the postmortem interval in rats up to seven days at 4°C. (9).

The rationale of these study to investigate different biochemicals in serum or liver tissue homogenate enhanced identification of postmortem interval on rats.

II. MATERIALS AND METHODS

a) Animal Protocol

Animal procedures were conducted with the approval of the animal care committee of the ethics Board of the faculty of veterinary medicine mansoura university (EGYPT).

25 male Albino rats (weight, 230-260 g; age were purchased from faculty of pharmacy, mansoura University, EGYPT). They were maintained on a 12-h light/dark cycle with free access to food and water. Brain

Author α: Department of Forensic Medicine and Toxicology, Faculty of Veterinary Medicine, Mansoura University, Egypt.

e-mail: mahmoudelalfy@mans.edu.eg

Author ο: Medical Physiological Department, Faculty of Medicine, Mansoura University, Egypt.

stem death was induced according (10). Dead rat were kept in fixed supine position on temperature for 30 hour post mortem (august 2017).

Blood samples of each 4 rat for each time point were collected from the heart and great vessels at autopsy at 0, 2, 4, 8, 16 and 30 hours postmortem. Samples were centrifuged immediately for 20 min at 5000 rpm. Additionally, blood samples collected from living rat from retroflexus serve as control physiological group and separated for 20 min at 5000 rpm. Serum samples was stored at - 70 liquid nitrogen until measurement.

b) Quantitative pH and biochemical activity detection

Blood pH was detected using pH meter (Sigma-Aldrich). Certified calibration buffer standards (4 or 8 pH) (Sigma-Aldrich) were used before each pH analysis. AST, ALT, BuN, creatinine and lactate dehydrogenase were analysed in serum samples.

c) Concentration of serum FAS ligand and TNF α were detected on by ELISA

The monoclonal antibodies of rat TNF α (sc52746) and Fas (sc-74540) used for ELISA were used as follows: ELISA plates were coated with rat TNF α and Fas antibody (5 μ g/ml) and then incubated overnight at 4°C. The plates were blocked with phosphate-buffered saline (PBS)-3% bovine serum albumin (BSA) for 1 h at room temperature and then incubated with the test

sample for 4 to 5 h at room temperature. The plates were washed and sequentially incubated with biotinylated secondary antibody, avidin-alkaline phosphatase, and substrate. The OD was read at 405 nm by using ELISA reader.

d) Statistical Analysis


Statistical analysis was carried out using the student's t-test. $P < 0.05$ was considered significant.

III. RESULTS

Post mortem time interval still controversial, to better understand the determinant biochemicals identify time elapsed since death, we induced death in male albino rats and measured different biochemicals in first 30 hours post death at defined times point.

AST, ALT, BUN and creatinine shown no change except little increase in last two successful time points see figure (a, b). Additionally, it was found that PH reduced as lactate escape from tissue while serum lactate dehydrogenase was increased in time dependent manner as leakage from tissue and may be a part of its increase due to hemolysis figure (1 c, d).

Notably, s-fas and tnfa increase in dependent manner until last time point see figure 1(d, e). On conclusion no group of biochemicals could be used in estimation of postmortem time interval for each time point.



Figure

IV. DISCUSSION

Post mortem time interval is still have interest for research from forensic point as no definitive markers estimate successful time since death even there are

many studies on human or experimental models investigate physical, biochemical and molecular marker for identification periods of time elapsed after death. After death, body loss its immune defence, aerobic oxidation and no elimination of breakdown of each cells

which could be explain elevation of certain biochemical parameters in serum of dead animal.

The aim of this study was to investigate metabolic changes that occur in blood post-mortem, with a view towards identifying biochemical markers that have potential for use in determining post-mortem interval. Blood pH and the concentrations of serum lactate dehydrogenase, AST, ALT, creatinine, BUN, TNFa and s-FAS changes post-mortem in rat corpses blood.

Blood is potential material as it remained liquid inside the rat corpses over the 96 hour of observation period due to release of active fibrinolysin enzymes from the vascular wall (11, 12 and 5). Post-mortem blood pH fell from 7.4 to 6.0 within the first 30 hours after death. Blood pH reduced post-mortem due to accumulation of lactic acid which resulted from anaerobic oxidation, blood stagnation in most dependant part and loss of buffer system (13 and 5).

Hepatocyte autolysis result in release of liver enzyme but this only happen when the putrefaction start in the corpse (5 and 7). While stop renal function retain creatinine but not occur immediately after death as Proteolysis did not occur uniformly throughout the body due to the resistance of some proteins such as collagen to breakdown (14). Blood does not contain substances that elevate the creatinine level Serum creatinine correlated significantly with lean mass (15) and so muscle creatinine may entered the blood and elevated the serum creatinine level.

Lactate dehydrogenase was decreased to 131 U/mg 24 hours post mortem on human dental pulp (16). Also a loss of lactate dehydrogenase activity was observed and distinguish between fresh and frozen-thawed fish fillets (17). Notably, lactate and malate dehydrogenases were detected in tissue extracts of human liver kept at 5 different temperatures until 35 days after death. The investigated activities of lactate dehydrogenase were reduced in proportion to time of storage which enabled detection of time elapsed after death (18). As in the current study, serum lactate dehydrogenase activity was increased by prolonged storage of dead rat due to leakage from tissue and also little hemolysis may be a pârt of increase level of lactate dehydrogenase.

In the present study, both sfas and tnfa were increased especially at 30 hours after death induction. Increased Fas and FasL immunoreactivity was seen in the rat cortex after brain injury site from 15 minutes to 72 hours after the trauma (19). Serum sFas, TNF- α levels can be useful as biochemical markers for early selection of patients at risk of deterioration after advanced degree of traumatic brain injury (20 and 21).

On conclusion, group of biochemical could be used in estimation of postmortem time interval for each time point.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Sara C. Zapico · Sofía T. Menéndez Paula Núñez. (2014) Cell death proteins as markers of early postmortem interval Cell. Mol. Life Sci71: 2957–296.
2. Amadasi A1, Cappella A2, Cattaneo C2, Cofrancesco P3, Cucca L3, Merli D4, Milanese C3, Pinto A3, Profumo A3, Scarpulla V2, Sguazza E2. Determination of the post mortem interval in skeletal remains by the comparative use of different physico-chemical methods: Are they reliable as an alternative to 14C? Homo. 2017 May; 68(3): 213-221.
3. Sharmaa Ruchi, Kumar Rakesh and Gaurb Garga J.R. (2015) Various methods for the estimation of the post mortem interval from Calliphoridae. Egyptian Journal of Forensic Sciences Volume 5, Issue 1, March, Pages 1-12.
4. Mathur Ashima and Agrawal Y. K. (2011) An overview of methods used for estimation of time since death journal of Australian Journal of Forensic Sciences_Volume 43, - Issue 4.
5. Donaldson AE, Lamont IL (2013) Biochemistry Changes That Occur after Death: Potential Markers for Determining Post-Mortem Interval. PLoS ONE 8(11): e8 2011.
6. Nidhi Sachdeva, **Yashoda Rani, ***Ritu Singh and Atul Murar Estimation of Post-Mortem Interval from the Changes in Vitreous Biochemistry J Indian Acad Forensic Med. April-June 2011, Vol. 33, No. 2.
7. Costa Isabel, Carvalho Félix, Magalhães Teresa, Pinho Paula Guedes de, Silvestre Ricardo and Dinis-Oliveira Ricardo Jorge (2015): Promising blood-derived biomarkers for estimation of the postmortem interval† Toxicol. Res., 4, 1443–1452 | 1443.
8. Zapico C S¹, Menéndez ST, Núñez P 2014 Cell death proteins as markers of early post mortem interval. Cell Mol Life Sci_ 2014 Aug; 71(15): 2957-62.
9. Kikuchi K¹, Kawahara KI, Biswas KK, Ito T, Tancharoen S, Shiomi N, Koda Y, Matsuda F, Morimoto Y, Oyama Y, Takenouchi K, Miura N, Arimura N, Nawa Y, Arimura S, Jie MX, Shrestha B, Iwata M, Mera K, Sameshima H, Ohno Y, Maenosono R, Tajima Y, Uchikado H, Kuramoto T, Nakayama K, Shigemori M, Yoshida Y, Hashiguchi T, Maruyama I 2010, HMGB1: A new marker for estimation of the postmortem interval. Exp Ther Med. 2010 Jan; 1(1): 109-111. Epub Jan 1.
10. Zhang S1, Cao S, Wang T, Yan B, Lu Y, Zhao Y. Modified brain death model for rats. Exp Clin Transplant. 2014 Oct; 12(5): 469-73.
11. Mole RH (1948) Fibrinolysin and the fluidity of the blood post mortem. J Pathol Bacteriol 60: 413-427.10.1002/path.1700600308 [Ref list].

12. Takeichi S, Wakasugi C, Shikata I. Fluidity of cadaveric blood after sudden death: Part II. Mechanism of release of plasminogen activator from blood vessels. *Am J Forensic Med Pathol.* 1985 Mar; 6(1): 25-9.
13. Sawyer WR, Steup DR, Martin BS, Forney RB (1988) Cardiac blood pH as a possible indicator of postmortem interval. *J Forensic Sci* 33: 1439-1444.
14. Dent, B.B., Forbes, S.L., Stuart, B.H. (2004, October 15). Review of Human Decomposition Processes in Soil. In *Environmental Geology*, 45, 576-585.
15. Alessandra Calábria Baxmann, Marion Souza Ahmed, Natália Cristina Marques, Viviane Barcellos Menon, *Aparecido Bernardo Pereira, Gianna Mastroianni Kirsztajn, and Ita Pfeferman Heilberg (2008): Influence of Muscle Mass and Physical Activity on Serum and Urinary Creatinine and Serum Cystatin C *Clin J Am Soc Nephrol.* Mar; 3(2): 348-354.
16. Caviedes-Bucheli J1, Avendaño N, Gutierrez R, Hernández S, Moreno GC, Romero MC, Muñoz HR. Quantification of lactate-dehydrogenase and cell viability in postmortem human dental pulp. *J Endod.* 2006 Mar; 32(3): 183-5.
17. Diop M1, Watier D2, Masson PY3, Diouf A4, Amara R5, Grard T6, Lencel P7F Assessment of freshness and freeze-thawing of sea bream fillets (*Sparus aurata*) by a cytosolic enzyme: Lactate dehydrogenase. *Chem.* 2016 Nov 1; 210: 428-34. Doi.
18. Gos T1, Raszeja S. Postmortem activity of lactate and malate dehydrogenase in human liver in relation to time after death. *Int J Legal Med.* 1993; 106(1): 25-9.
19. Beer Ronny, Franz Gerhard, Schopf Marion, Reindl Markus, Zelger Bernhard, Schmutzhard Erich, Poewe Werner, and Kampfl Andreas (2000): Expression of Fas and Fas Ligand After Experimental Traumatic Brain Injury in the Rat *Journal of Cerebral Blood Flow and Metabolism* 20: 669-677.
20. Qiu J1, Whalen MJ, Lowenstein P, Fiskum G, Fahy B, Darwish R, Aarabi B, Yuan J, Moskowitz MA. Upregulation of the Fas receptor death-inducing signaling complex after traumatic brain injury in mice and humans. *J Neurosci.* 2002 May 1; 22(9): 3504-11.
21. Badr Rawia, Badr Hatem, Aly Nabil and Ibrahim Abdelwahab (2013): Serum sFas, TNF- α and Bcl2 expression following Traumatic Brain Injury *egyptian Journal of Medical Microbiology*, April Vol. 22, No. 2 pp1-11.