



GLOBAL JOURNAL OF MEDICAL RESEARCH: J  
DENTISTRY & OTOLARYNGOLOGY  
Volume 18 Issue 1 Version 1.0 Year 2018  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## A Clinico-Pathological Presentation of Solitary Thyroid Nodule

By Dr. Ishank Jaiswal, Dr. Nicola C. Lyngdoh, Vidhiyaveeran, Rituparnasaha,  
Roshan Ali, Markhia Bhuphang & Pulokkutum

*Regional Institute of Medical Sciences*

**Abstract- Background:** Thyroid nodules represent a difficult diagnostic problem. They are quite common in the population, while clinically apparent thyroid cancer the principal concern of both patient and physician is comparatively rare.

**Materials and method:** We prospectively evaluated the clinical and pathological profile of 50 consecutive patients presenting with solitary thyroid nodule.

**Results:** Most common age group was 30-39 years (4th decade) with a female: male ratio=6:1. Overall malignancy rate was 32%. The diagnostic assessment by Fine needle aspiration cytology in this study produces sensitivity of 92.71%, specificity of 78.26%, positive predictive value of 83.33% and negative predictive value of 90% thereby having an overall accuracy of 86%. Histopathological examination remains the gold standard for final accurate diagnosis.

**Keywords:** solitary thyroid nodule, fine needle aspiration cytology (FNAC), histopathological examination (HPE).

**GJMR-J Classification:** NLMC Code: WK 200



*Strictly as per the compliance and regulations of:*



# A Clinico-Pathological Presentation of Solitary Thyroid Nodule

Dr. Ishank Jaiswal <sup>α</sup>, Dr. Nicola C. Lyngdoh <sup>σ</sup>, Vidhiyaveeran <sup>ρ</sup>, Rituparnasaha <sup>ω</sup>, Roshan Ali <sup>ξ</sup>,  
Markhia Bhuphang <sup>§</sup> & Pulokkutum <sup>x</sup>

**Abstract- Background:** Thyroid nodules represent a difficult diagnostic problem. They are quite common in the population, while clinically apparent thyroid cancer the principal concern of both patient and physician is comparatively rare.

**Materials and method:** We prospectively evaluated the clinical and pathological profile of 50 consecutive patients presenting with solitary thyroid nodule.

**Results:** Most common age group was 30-39 years (4<sup>th</sup> decade) with a female: male ratio=6:1. Overall malignancy rate was 32%. The diagnostic assessment by Fine needle aspiration cytology in this study produces sensitivity of 92.71%, specificity of 78.26%, positive predictive value of 83.33% and negative predictive value of 90% thereby having an overall accuracy of 86%. Histopathological examination remains the gold standard for final accurate diagnosis.

**Keywords:** solitary thyroid nodule, fine needle aspiration cytology (FNAC), histopathological examination (HPE).

## I. INTRODUCTION

A solitary thyroid nodule is a palpable discrete swelling within an otherwise apparently normal thyroid gland. Though it is a common disorder of the thyroid gland, it is less prevalent among children. Childhood thyroid nodules need special attention due to higher incidence of malignancy as compared to adults. Differential diagnosis of a thyroid nodule is crucial as malignancy necessitates surgery; while strict follow up is necessary in benign cases.<sup>1</sup> Fine needle aspiration cytology (FNAC) is a cost effective procedure that provides specific diagnosis rapidly with minimal complications. Our objectives were to study the demographic descriptions, clinical, cytological, histopathological profile and correlation of cytological finding with that of histological, of patients with solitary thyroid nodule attending our tertiary referral centre.

## II. MATERIALS AND METHODS

Fifty consecutive patients irrespective of age, sex, religion and socio-economic status with solitary thyroid nodule admitted in the department of Otorhinolaryngology, Regional Institute of Medical

Sciences, Imphal between August 2015 to September 2017 were prospectively studied.

Every patient was meticulously worked up with a thorough history taking and clinical examination followed by routine laboratory investigations and investigations specific to thyroid like thyroid hormone assay, Fine needle aspiration cytology (FNAC) and Ultrasonography (USG) of the thyroid. Post operatively the thyroidectomy specimens were sent for Histopathological Examination (HPE).

## III. RESULTS AND OBSERVATION

The age of the patients ranged from 15 to 70 years. The youngest patient was a 15 year old male while the oldest was a 70 year male. The female to male ratio was 6.1:1.25; (50%) cases presented with swelling with duration of less than 1 year while 3(6%) cases had duration of 9-10 year. The shortest duration of swelling was 2 months which was diagnosed as follicular adenoma, while the longest was for 10 years which was diagnosed as colloid goiter.

All the 50 patients came with chief complaint of swelling in front of the neck of whom 46(92%) had a gradual increase in size while in 4(8%) cases there was rapid increase in size. There was Dysphagia in 2(4%) cases. Associated pain, difficulty in breathing, hoarseness, decreased appetite, weight loss and cervical lymphadenopathy were seen in a 70 year old male patient who was diagnosed as papillary carcinoma. Firm consistency was observed in 38(78%) cases of which maximum were of colloid goiter while other varieties of consistency (soft/cystic) in follicular adenoma, papillary carcinoma, hashimoto's thyroiditis and Hurthle's neoplasm were also seen. One case had hard consistency with metastatic lymphadenopathy.

In FNAC 30 (60%) cases were found to be colloid goiter while papillary carcinoma was seen in 9 (18%) cases. There were 7 (14%) cases with follicular neoplasm, 2 (4%) cases of Hurthle's neoplasm and 1 (2%) each of Hashimoto's and Lymphocytic thyroiditis. On Ultrasonography 30 (60%) cases were of cystic nature while 12 (24%) were solid and 8 (16%) cases were of suspicious malignancy. On Histopathological examination of the operated tissues 27 (54%) cases were colloid in nature, 11 (22%) cases were papillary carcinoma while 3 (6%) cases were of papillary-follicular variant. There was also a case of squamous carcinoma

**Author α:** Post Graduate Trainee, Department of Otorhinolaryngology, Regional Institute of Medical Sciences, Imphal, Manipur, India.  
e-mail: jaiswalishank@gmail.com

**Author σ:** Associate Professor, Department of Otorhinolaryngology, Regional Institute of Medical Sciences, Imphal, Manipur, India.  
e-mail: nclyngdoh@gmail.com

in a female patient of 26 years who was diagnosed as colloid nodule on FNAC. Most of the solid findings and suspicious cases of malignancy on ultrasonography

were found to be papillary carcinoma and follicular adenoma on HPE. Majority of cystic findings on Ultrasonography were colloid goiter in nature.

*Table 1:* Characteristics of the patients presented with clinical solitary thyroid nodule

Age Distribution		
Age in years	No. of cases	Percentages (%)
0-9	0	0%
10-19	7	14%
20-29	8	16%
30-39	17	34%
40-49	8	16%
50-59	4	8%
60-69	5	10%
70 & above	1	2%
Sex Distribution		
Sex	No. of cases	Percentage (%)
Male	7	14%
Female	43	86%
Duration of Swelling		
Duration (months)	No. of cases	Percentage (%)
0-12	25	50%
13-24	10	20%
25-36	6	12%
37-48	4	8%
49-60	1	2%
61-72	1	2%
73-108	0	0%
109-120	3	6%
Symptoms of Patient		
Symptoms	No. of cases	Percentage (%)
Swelling	50	100%
Gradual progression	46	92%
Rapid progression	4	8%
Associated pain	1	2%
Dysphagia	2	4%
Difficulty in breathing	1	2%
Hoarseness	1	2%
Decrease appetite	1	2%
Weight loss	1	2%
Lymphadenopathy	1	2%
Location of Swelling		
Location	No. of cases	Percentage (%)
Right lobe	22	44%
Left lobe	18	36%
Isthmus	10	20%
Consistency of Swelling		
Consistency	No. of cases	Percentage (%)
Cystic	3	6%
Soft	8	16%
Firm	38	76%
Hard	1	2%

Table 2: Investigations Reports

FNAC Results		
Cytology	No. of cases	Percentage (%)
Colloid Goiter	30	60%
Papillary Carcinoma	9	18%
Follicular Carcinoma	7	14%
Hashimoto's Thyroiditis	1	2%
Hurthle's Neoplasm	2	4%
Lymphocytic Thyroiditis	1	2%
Ultrasound Results		
Ultrasound	No. of cases	Percentage (%)
Cystic	30	60%
Solid	12	24%
Suspected Malignancy	8	16%
Histopathological Results (HPE)		
HPE	No. of cases	Percentage (%)
Colloid Goitre	27	54%
Papillary Carcinoma	14	28%
Follicular Adenoma	6	12%
Follicular Carcinoma	0	0
Hashimoto's Thyroiditis	1	2%
Hurthle's Carcinoma	1	2%
Squamous Cell Carcinoma	1	2%

Out of 30 patients of colloid nodule on FNAC, 27 cases were found to be colloid goiter on HPE. The remaining turned out to be papillary carcinoma and squamous cell carcinoma in 1 case. All cases of papillary carcinoma on FNAC were diagnosed to be the same on HPE.

Hashimoto's and Hurthle's cases were almost same in both examinations. All except one case of


follicular neoplasm on FNAC turned out to be follicular adenoma on HPE. There was significantly more occurrence of malignancy at both the extremes of age, a rate of 42.85% below 20 years and 33.33% above 60 years. The youngest patient was a 15 year old male and the oldest 70 year old patient and both were diagnosed as papillary carcinoma.

Table 3: Comparison of FNAC/HPE reports


Type	FNAC (%)	Subdivision	HPE	%
Colloid Goitre	30 (60%)		27	54%
Papillary Carcinoma	9 (18%)		14	28%
Follicular Neoplasm	7 (14%)	Follicular Adenoma	6	12%
		Follicular Carcinoma	0	
Hashimoto's Thyroiditis	1 (2%)		1	2%
Hurthle's Neoplasm	2 (4%)	Hurthle's Carcinoma	1	2%
Lymphocytic Thyroiditis	1 (2%)			
Squamous Carcinoma			1	2%

Table 4: Carcinoma occurrence in Male/Female with correlation between age and malignancy

Female	No. of cases	Percentage %	Male	No. of cases	Percentage %
Malignant	13	30.2%	Malignant	3	42.8%
Benign	30	69.8%	Benign	4	57.2%
Age group		Total cases	Malignancy		Percentage
< 20 years		7	3		42.85%
> 60 years		6	2		33.33%
20-60 years		37	6		16.21%



*Fig. 1:* HPE under high magnification (40x) view shows cellular smear with cells arranged in papillary architecture having enlarged round to ovoid nuclei with nuclear overcrowding, overlapping and having fine granular powdery chromatin; suggestive of a papillary carcinoma



*Fig. 2:* FNAC smear under high magnification (40x) shows benign follicular cells arranged in clusters in a background of thin and thick colloid; suggestive of colloid goiter

#### IV. DISCUSSION

In the present study, the maximum number of cases occur in the age group of 30-39 years, i.e. 4<sup>th</sup> decade of life. Other studies also reported a higher incidence in the 3<sup>rd</sup> and 4<sup>th</sup> decade age group.<sup>2,3,4</sup>

Several studies have highlighted the greater risk of malignancy in thyroid nodules in younger age group

and older age-group people.<sup>5,6</sup> Several studies have highlighted the greater risk of malignancy in thyroid nodules in younger age group and older age-group people. Similar findings noted in our study. Female: Male ratio was 6.1:1, comparable to other literature.<sup>2,3,7</sup> The overall malignancy rate was 32%, similar to older literature.<sup>8,9</sup>



The most common presenting feature was swelling in the neck seen in all 50 cases (100%). This correlates with the observance made by other authors.<sup>4,7,10</sup> There was a gradual increase in size of the solitary thyroid nodule in 46 (92%) cases, while 4 (8%) cases had rapid progression. Dysphagia noted in 2 cases of papillary carcinoma and follicular adenoma. Associated pain with hoarseness, weight loss, difficulty in breathing, decreased appetite, and cervical lymphadenopathy was seen in 70-year-old patient with papillary carcinoma. All these clinical indicators suggestive of malignancy reported in other studies.<sup>8,11</sup>

The presence of nodule on the right lobe was in 22 (44%) cases, 18 (36%) in the left lobe and 10 (20%) in the isthmus. Similar findings reported by another author.<sup>10</sup> Firm consistency felt in 38 (76%) cases, cystic in 3 (6%) cases and soft in 8 (16%) cases. An elderly patient with hard nodule diagnosed as papillary carcinoma. Similar observations found in other studies.<sup>4,6</sup>

The diagnostic assessment by FNAC in this study produces a sensitivity of 92.71%, specificity of 78.26%, a positive predictive value of 83.33% and negative predictive value of 90%. The accuracy of FNAC found to be 86%. This finding was comparable to other studies.<sup>6,12</sup>

Histopathological examination (HPE) revealed a colloid occurrence in 54%, papillary carcinoma in 28%, follicular adenoma in 12%, 2% each in Hashimoto's and Hurthle's. These findings tally with that of other studies. The percentage of malignancy in our study was 29% whereas other series reported 8-37%.<sup>12,13</sup> The most common finding in FNAC in our study is colloid goiter which accounts for almost 60%, which is consistent with the other studies.<sup>14</sup> Out of total 14 cases of papillary carcinoma cases diagnosed from HPE, almost 9 (64.3%) cases were identified in FNAC also. This shows there is a chance of false negativity for papillary carcinoma, should always be kept in mind. It is consistent with the study conducted by Sukumaran et al.<sup>15</sup>

## V. CONCLUSION

The incidence is highest in the 4<sup>th</sup> decade of life. The rates of malignancy in extremes of age group are significantly higher than other general population. The incidence of a solitary thyroid nodule is much higher in female (F: M-6.1:1).

The most common mode of presentation is swelling in the neck with a majority of (92%) patients giving a history of gradual progression of size. Location of the nodule was slightly more on the right (44%) as compared to the left lobe and isthmus and most nodules presented with a firm consistency.

FNAC was found to be not only an easy and inexpensive mode of diagnosis but, also a highly accurate means of investigation with an accuracy of

86% in this study. Histopathological examination remains the gold standard for the final correct diagnosis of a solitary thyroid nodule.

## REFERENCES RÉFÉRENCES REFERENCIAS

1. Stergiois AP, Marina K, Auraam A. Thyroid nodule-stepwise diagnosis and management. *Hormones* 2007; 6(2): 101-119.
2. Kumar A, Ahuja MMS, Chattopadhy TK, Padhya AK, Gupta AK, Kapila K, Goel AK and Karmakar MG. Fine needle aspiration cytology, sonography and radionuclide scanning in solitary thyroid nodule. *JAssocPhysicians*1992; 40: 302-7.
3. Mandrekar SRS, Nadkarni NS, Pinto RGW, Menezee S. Role of fine needle aspiration cytology as the initial modality in the investigation of thyroid nodules. *Acta Cytol*1995; 39(5): 898-904.
4. Ananthakrishnan N, Madhava Rao K, Narashima R, Veliath AJ. Problems and limitatins with FNAC of solitary thyroid nodules. *ANZ J Surg* 2008; 60(1): 35-9.
5. George LAF, Victor GL. Thyroid disease. In: Stephen A, Falk editor, *Endocrinology, Surgery, Nuclear medicine and Radiotherapy*. Philadelphia: Lippincott Raven publishers; 1997.p. 411-429.
6. Paul RM, Malcolm HW. Approach to thyroid nodules. In; Clark-Duh-Kelsebew editor, *Textbook of Endocrine surgery*. Philadelphia: Elsevier Saunders Publishers; 1996. p. 85-91.
7. Bouvet M, Feldman JI, Gill GN, Dillman WH, Nahum AM, Russack V, Robbins KT. Surgical management of the thyroid nodule, patient selection based on the results of fine needle aspiration cytology. *Laryngoscope* 1992; 102: 1353-6.
8. Ashok RS. Controversies in management of thyroid nodule. *Laryngoscope* 2000; 110: 183-5.
9. Kamaljit K, Nishi S, Bapna AS, Pradeep M. A comparative study of FNAC, USG and Radio nucleotide scan in the management of solitary thyroid nodule: A prospective analysis of 50 cases. *Indian JOtolaryngolHead Neck Surg* 2002; 54(2): 96.
10. Amjad I, Muhammed I, Nisar A, Javed A and Javed Z: Management of solitary thyroid nodule. *Postgrad MedJ* 2005; 19(1): 1-2.
11. Barroeta JE, Wang H, Shiina N, Gupta PK, Livolsi VA and Baloch ZW. Fine needle aspiration of multiple thyroid nodule justified? *EndocrPathol* 2006; 17(1): 61-6.
12. Cusick EL, Mac-Intosh CA, Zygmunt HK, Williams VM, Matheson NA. Management of Isolated thyroid swelling: a perspective six year study of fine needle aspiration cytology in diagnosis. *Brit MedJ* 1990; 301: 318-321.
13. Ali Z. New insight into the diagnosis and management of thyroid nodules. *ShirazEMedJ* 2005; 6(1, 2): 1-2.

15. Sukumaran R, Kattor J, Pillai KR, Ramadas PT, Nayak N, Somanathan T, George NA, Sebastian P. Fine Needle Aspiration Cytology of Thyroid Lesions and its Correlation with histopathology in a series of 248 patients. Indian J SurgOncol 2014; 5(3): 237-241.

