

Immediate and Short Term Effect of Combination Physical Therapy and Manual Therapy in Post ACL Reconstruction Surgery Displaced Mal-Alignment of the Joint. An Invivo Clinical Trial

Suraksha Shanbag

Received: 7 December 2017 Accepted: 4 January 2018 Published: 15 January 2018

Abstract

Post ACL Reconstruction surgery displaced mal-alignment of the joint due to re-injury is rare and difficult to treat the condition, especially in young adult and old age when they develop kinesiophobia (fear of pain due to movement) and pain. During this time if re-surgery is not appreciated physical therapy has to play a very important role in the rehabilitation. Combination of different physiotherapy protocols has to be taken into consideration by the well-experienced therapist to show the immediate effect on the pain and other symptoms to build the confidence of the patient during the therapy. In the present study we used the combination of soft tissue myofascial release technique with MT2 blade, mobilization with belt and manual followed by Kinesio tapping to see the immediate effect. Active exercise therapy was included to carry-on the strengthening and to see short-term effect of the entire Physiotherapy protocol. The present study suggested beneficial effects in combination of physiotherapy treatment protocols.

Index terms—

Abstract-Post ACL Reconstruction surgery displaced malalignment of the joint due to re-injury is rare and difficult to treat the condition, especially in young adult and old age when they develop kinesiophobia (fear of pain due to movement) and pain. During this time if re-surgery is not appreciated physical therapy has to play a very important role in the rehabilitation. Combination of different physiotherapy protocols has to be taken into consideration by the wellexperienced therapist to show the immediate effect on the pain and other symptoms to build the confidence of the patient during the therapy. In the present study we used the combination of soft tissue myofascial release technique with MT2 blade, mobilization with belt and manual followed by Kinesio tapping to see the immediate effect. Active exercise therapy was included to carry-on the strengthening and to see short-term effect of the entire Physiotherapy protocol. The present study suggested beneficial effects in combination of physiotherapy treatment protocols.

1 I. Introduction

anterior cruciate ligament is known to be one of the key structures in the knee and one of the commonest injured or sprained ligament in the body. Not only in sports but it has the highest prevalence in any other normal human being. Most common risk factors seen are due to sudden changing of direction, stopping suddenly, and landing from height or incorrectly jumping and direct contact or collision during any fall 1 . A study done by Hideyuki Koga et al.

proposed a new mechanism of injury that believes ACL ruptures take place within the first 40 minutes after initial ground contact. Due to the unchanged hip joint angle at an initial position lateral knee compression causes valgus loading. The anterior force vector causes quadriceps contraction. Displacement of the femur relative to

the tibia takes place and the lateral femoral condyle shifts posteriorly due to the joint geometry. Tibia translates anteriorly and rotates internally, thereby resulting in ACL rupture 2 .

Though there have been many recent advances in ACL Reconstruction surgery's and rehabilitation there is less explanation of the same during the re-injure phase and kinesiophobia. A study finding suggested that young patients are at high risk of sustaining a second ACL injury within 24 months after ACLR 3 . Reasons may be many of which age, sex, knee-related confidence, and performance on the triple hop for distance were mentioned as the primary predictors of a second ACL injury 4 . Another study concludes that there is a high prevalence (61.69%) of kinesiophobia in Young adults post ACL reconstruction for 4 to 8 weeks 5 . The second injury after the primary ACL reconstruction and rehabilitation goes under different positioning of the underline structures. Also, it is difficult to reconstruct and rehabilitate again due to more kinesiophobia and pain. Another study suggests diminished knee related quality of life and social reasons or psychological hindrances such as fear of re-injury may influence return to sports or other daily activities 6,7 . In the present study to present the pre influence of kinesiophobia related to re-injury of the knee was recorded by The Tampa Scale of Kinesiophobia (TSK). The same scale was used to see the post influence during and after the treatment. This scale measures patients' fear of movement 8,9 .

Combinations of physical therapies have always shown good results than selecting and deciding to go with a one-line treatment. It is the crucial point where the therapist should be highly experienced to detect which structures are injured and need treatment along with how to carry and which all techniques to be include in the therapy with the concern of the patient convenience and financial status. In the present case, patient was also had kinesiophobia along with nerve entrapment which made the decision difficult to build his confidence to cooperate with active and resisted treatments such as weight-bearing exercise and gait training. We so decided to go with Instrument-assisted soft tissue mobilization (IASTM) for stiffness and pain followed by cryotherapy. We used the M2T blade which is a newly designed Instrument-assisted soft tissue mobilization (IASTM) that is three dimensional analytic and treatment tool, enables clinicians to assess, detect, and treat individuals diagnosed with scar tissue, adhesions, and soft tissue dysfunction 10,11 . Knee mobilization was included with and without mobilization belt as to free the joint range and nerve entrapment. Many studies give different finding why mobilization is so-important in postoperative cases of any joints especially knee joint. One of most is that Joint mobilization may assist in reducing pain and increasing joint range by passive oscillatory movements of small or large amplitude and sustained stretching within the anatomical limits and at different directions 12 . Based on the different literature we also used Kinesio tapping in the study 13 . Applications of Kinesio tapes were to facilitate contraction and increase muscle strength and to maintain the joint space. After the above treatment protocol conservative treatments like stretching, strengthening and biofeedback for gait training were done.

2 II. Case Report

A 30-year-old male with secondary injury and malalignment of the knee joint structure attended our clinic with primary complain of pain and buckling of the knee. The patient also gave the history of burning sensation in lateral aspect of the knee till the mid-upper superior part of the lower leg. On assessment, patient was having limping gait and decreased strength in the affected extremity. The patient had no external injury or exposed area. On special test, patient was positive for Lachman Test, Positive Lateral Pivot Shift and Positive Anterior Drawer test. Patients explained numbness in lateral aspect of the knee till the mid-upper superior part of the lower leg. The X-Ray also revealed slight malposition of the primary ACLR screw implant laterally. The slight malposition of the primary ACLR screw implant laterally was causing compression to the common peroneal nerve at the sural nerve branch originating at the lateral end of lateral condyle and meniscus over lateral aspect of tibia head leading to numbness, sometime burning sensation and paraesthesia at in lateral aspect of the knee till the mid-upper superior part of the lower leg. The patient also complained of the gradual increase in the weakness in the affected leg during walking (mild circumduction gait during walking were noticed).

After informed consent was explained to the patient in his vernacular language and signed by the patient, he was taken into present study. Pre-data were recorded before any enhancement of the treatment. The patient was instructed to thoroughly follow the exercise protocol advised by the therapist and not to undergo any other treatment or medication till study sessions get completed. In the present study, we used outcome measure as visual Analog Scale, pre-treatment video on Gait analysis, areas of diminished sensation and the borders of distribution, lower limb strength.

3 III. Discussion

The present case is rare and difficult to treat the case as due to the secondary injury to the knee joint there is increase in kinesiophobia and pain along with nerve interruption. A young patient is always scared of any treatment and little cooperative to the active physical therapy. In such cases, patient should be given the confidence to practice independent activity sessions. As we discussed, we found that more the pain more uncomfortable the patient will be at the therapy sessions. Hence the goals of the study were to decrease the pain first along with numbness and then to increase the strength and mobility of the lower limbs followed by gait training.

In the present study, we used a combination of the different physical therapy techniques to encounter the pain, numbness and joint mobility at the same site as they were related to each other. We used soft tissue release technique with the help of MT2 blade. MT2 blade uses the principal of myofascial release wherein a stretch is applied on the tight fascia that is maintained for 90-120 seconds prove to lengthen the tight fascia. Other studies on MT2 blade have shown beneficial effects at the cellular, level reduction in scar and increase vascular response 14,15 . MT2 blade therapy also releases fascia leading to pain relief. These effects of MT2 were beneficial in our case as to release the surrounding structures of the knee joint before performing joint mobilization. This therapy gave pain relief and loosened hold on the knee joint and other surrounding structures around it. After MT2 Cryo-therapy was used to reduce if-any soreness caused due to instrument assisted soft tissue muscle relies. We used manual mobilization 16,17 and belt mobilization in combination with knee glides to decrease the compression of the lateral-end of lateral-condyle and meniscus over lateral-aspect of tibia head that was leading to compression of the sural nerve. Immediately after the soft-tissue release patient was asked to seat at the bed end and distraction in downwardly med-line position was applied. This created the space between the lateral end of lateral condyle and meniscus over lateral aspect of tibia head reducing the compression force. A posterior-medial glide in the midplane of the knee joint pushed the lower meniscus position with antero-posterior glide to tibio-femoral joint in such a position that it released the pressure from the common peroneal nerve at the origin of sural nerve. This relieved the nerve compression with the reduction in pain and burning sensation immediately. In the same position, we applied Kinesio tape to the knee joint to hold the position in place and for other beneficial effects on ACL reconstruction rehabilitation which are widely known 18,19,20 .

The above combination of three therapies gave immediate effect on pain reduction and reduction in burning sensation and numbness. The patient was asked to perform some of the active exercises after the manual session for the strengthening of the lower limb. Manual mobilization techniques were continued for 15 sessions alternatively. Exercises were performed on daily bases under the observation of the therapist. Gait training was also started with visual feedback from the 8th day of the session.

4 IV. Results

Tampa Scale-11 (TSK-11) showed significant changes from pre-assessment to day 1, 2, 6 and 12 depicted in fig 3 ?? The patient was comfortable to walk with the slight pain on day 1 post-treatment as pain reduced from VAS 8 to VAS 5. On the next session, the patient was feeling comfortable and pain-free to do free and light weighted exercises. As the session was regular on daily bases at the 6 th session patient was able to walk with a great improvement in his gait pattern. At the 12 th session, patient showed significant improvement in walking with no pain and fear of the reoccurrence of reinjures. Burning sensation and paraesthesia at in lateral aspect of the knee till the mid-upper superior part of the lower leg also showed a great reduction on the first-day post session. The joint space was maintained by mobilization in an uncompressed position of the sural nerve and prolonged support by Kinesio taping in same position. Relief in nerve entrapment further reduced nerve inflammation. The patient reported less frequent episodes of burning and paraesthesia in from day 2 to day 6 and nil to 1 episode only if done heavy stepping movements. Active exercises with theraband also showed significant improvement in increasing the lower limb strength.

5 V. Conclusion

A case presented with anatomical, physical as well as psychological conditioning is very-difficult to treat in any type of rehabilitation protocol. If the patient is kinesiophobic or having any other psychological fear it is very difficult to treat, hence careful assessment and gaining of the patient confidence in physical activity or exercise and towards the therapist is of outmost important. In the present case, we decided to relieve the stiffness and pain of the superficial area and then to do mobilization to relieve inner compression and stiffness of the joint. This gives confidence to the patient to move the joint and that the therapy will not cause further injury to him. Kinesio taping was also impotent to support the joint during exercises in correct and uninterrupted pain free position of sural nerve and other structures of the knee joint. Theraband exercises increased the strength in lower limb and biofeedback during gait training was also beneficial.

6 Conflicts of Interest:

All authors declare no conflicts of interest. ¹ ²

¹© 2018 Global Journals

²© 2018 Global Journals 1



1

Figure 1: Fig. 1 :



2

Figure 2: Fig. 2 :

Figure 3:

[Arthroscopy: The Journal of Arthroscopic Related Surgery (2018)] , *Arthroscopy: The Journal of Arthroscopic & Related Surgery* 2018 May 1. (5) p. .

[ACL Injury Mechanisms: Lessons Learned from Video Analysis] *ACL Injury Mechanisms: Lessons Learned from Video Analysis*,

[Lentz et al. (2012)] *Analysis of shortened versions of the Tampa Scale for Kinesiophobia and Pain Catastrophizing Scale for patients following anterior cruciate ligament reconstruction. The Clinical journal of pain*, George S Z Lentz , T A Zeppieri Jr , G , Lee D Chmielewski , TL . 2012 Jan: 28. p. 73.

[Anterior Cruciate Ligament (ACL) Injuries (2018)] <https://orthoinfo.aaos.org/en/diseases--conditions/anterior-cruciate-ligament-acl-injuries/viewedon21> *Anterior Cruciate Ligament (ACL) Injuries*, may 2018.

[Clinical Factors That Predict a Second ACL Injury After ACL Reconstruction and Return to Sport 5. Prevalence of Kinesiophobia] *Clinical Factors That Predict a Second ACL Injury After ACL Reconstruction and Return to Sport 5. Prevalence of Kinesiophobia in Young Adults Post ACL Reconstruction*,

[Clinical Factors That Predict a Second ACL Injury After ACL Reconstruction and Return to Sport: Preliminary Development of a Clinical Decision Algorithm] *Clinical Factors That Predict a Second ACL Injury After ACL Reconstruction and Return to Sport: Preliminary Development of a Clinical Decision Algorithm*,

[Calloway et al.] *Clinical Outcomes After Arthroscopic Release of Patellofemoral Arthro-fibrosis in Patients With Prior Anterior Cruciate Ligament Reconstruction*, S P Calloway , C J Soppe , B R Mandelbaum .

[Brown et al. (2016)] *Decreased range of motion after total knee arthroplasty is predicted by the Tampa Scale of Kinesiophobia. The Journal of arthroplasty*, M L Brown , J F Plate , Von Thae , S , Fino N F , Smith B P Seyler , TM , Lang J E . 2016 Apr 1. p. .

[Nadali et al. ()] 'Effect of kinesio taping on vertical jump after ACL reconstruction'. S Nadali , M A Mahdi Amel Khabazan , A S Aryamanesh , A Hoseleh , M H Mohamad Hosein Khabaz , A Bakhshizadeh . *Intl J Sport Std* 2014. 4 (6) p. .

[Begalle et al. (2017)] 'Fear of Re-Injury Remains High in Adolescent Female Athletes Following Anterior Cruciate Ligament Reconstruction and Full Return to Sport'. R L Begalle , J T Blackburn , D A Padua . *Journal of Athletic Training* 2017 Jun 1. (6) p. S43.

[Kvist et al. (2005)] *Fear of reinjury: a hindrance for returning to sports after anterior cruciate ligament reconstruction. Knee surgery, sports traumatology, arthroscopy*, J Kvist , A Ek , K Sporrstedt , L Good . 2005 Jul 1. p. .

[Gehlsen and Helfst ()] 'Fibroblast responses to variation in fckLR soft tissue mobilization pressure'. G M Gehlsen , Ganion L R Helfst , RH . *Med Sci Sports Exerc fckLR* 1999. 31 p. .

[Oliveira et al. ()] 'Immediate effects of Kinesio Taping® on neuromuscular performance of quadriceps and balance in individuals submitted to anterior cruciate ligament reconstruction: a randomized clinical trial'. A K Oliveira , D T Borges , C A Lins , R L Cavalcanti , L B Macedo , J S Brasileiro . PP. 2-6 PII: S1440-2440. *J Sci Med Sport* 2014. 19 (1) p. .

[Oliveira et al. (2016)] *Immediate effects of Kinesio Taping® on neuromuscular performance of quadriceps and balance in individuals submitted to anterior cruciate ligament reconstruction: a randomized clinical trial*, A K Oliveira , D T Borges , C A Lins , R L Cavalcanti , L B Macedo , J Brasileiro . 2016 Jan 1 : 19. p. . (Journal of science and medicine in sport)

[Naik et al. ()] 'Immediate Effects of M2T Blade on Pain and Range of Motion Inrecreational Badminton Shoulder Pain Subjects: A Pilot Study'. D Naik , R Shaikh , S Koyande . *Journal of Medical Science And clinical Research* 2016. (10) p. .

[Fernandez-De-Las-Penas ()] 'Joint mobilization enhances mechanisms of conditioned pain modulation in individuals with osteoarthritis of the knee'. Courtney C A , Steffen A D Fernandez-De-Las-Penas , C . *J Orthop Sports Phys Ther* 2016. 46 p. .

[Wanivenhaus et al. (2016)] 'Novel protocol for knee mobilization under femoral and sciatic nerve blocks for postoperative knee stiffness'. F Wanivenhaus , P M Tscholl , Fucentese S F Aguirre , J A Giger , R . *Orthopedics* 2016 Jul 25. 39 (4) p. .

[The basic science of myofascial release: morphologic changes in connective tissue M. F. Barnes. Journal of bodywork and movement therapies] *The basic science of myofascial release: morphologic changes in connective tissue M. F. Barnes. Journal of bodywork and movement therapies*, 1997.

[Donec and Krisciunas ()] 'The effectiveness of Kinesio Taping® after total knee replacement in early postoperative rehabilitation period. A randomized controlled trial'. V Donec , A Krisciunas . *Eur J Phys Rehab Med* 2014. 50 (4) p. .

[Tan et al.] *Theory of Blade Design for Large Deflections: Part II-Annular Cascades CS*, W R Tan , J E Hawthorne , C McCune , Wang .