

GLOBAL JOURNAL OF MEDICAL RESEARCH: G VETERINARY SCIENCE AND VETERINARY MEDICINE Volume 18 Issue 2 Version 1.0 Year 2018 Type: Double Blind Peer Reviewed International Research Journal Publisher: Global Journals Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## Survey on Pathological Lesion and its Financial Losses in Ovine Slaughtered at Jimma Municipal Abattoir, Jimma, Ethiopia

By Mohammed Abatemam, Endegena Taye, Dinaol Belina & Abu Urji Haramaya University

Abstract- A cross-sectional study was conducted from October 2016 to July 2017 on ovine slaughtered at at Jimma municipal abattoir with the aim of identifying main pathological lesions causing organs and carcass condemnation, and associated direct financial losses. In this, 384 sheep were recruited to the study using systematic random sampling and standard antemortem (AM), and postmortem inspection (PMI) procedures were employed. Nasal discharge, tick infestation, coughing, lameness, emaciation, depression and salivation are recorded as the major AM findings of the current study. Accordingly, 47 (12.2%) sheep showed signs of diseases and abnormalities; of which 23 (6%) were conditionally approved whereas 11 (2.9%) sheep were unfit and judged to be detained and rejected. In the present study age, body conditions and geographic origin of the animals were considered as study variables, and the results showed BCS and age groups had statistically higher ( $p \le 0.05$ ) rejection probabilities.

Keywords: abattoir, financial loss, lesion, organ condemnation, PMI, sheep, jimma.

GJMR-G Classification: NLMC Code: WB 141



Strictly as per the compliance and regulations of:



© 2018 Mohammed Abatemam, Endegena Taye, Dinaol Belina & Abu Urji. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Survey on Pathological Lesion and its Financial Losses in Ovine Slaughtered at Jimma Municipal Abattoir, Jimma, Ethiopia

Mohammed Abatemam <sup>a</sup>, Endegena Taye <sup>o</sup>, Dinaol Belina <sup>o</sup> & Abu Urji <sup>a</sup>

Abstract- A cross-sectional study was conducted from October 2016 to July 2017 on ovine slaughtered at Jimma municipal abattoir with the aim of identifying main pathological lesions causing organs and carcass condemnation, and associated direct financial losses. In this, 384 sheep were recruited to the study using systematic random sampling and standard antemortem (AM), and postmortem inspection (PMI) procedures were employed. Nasal discharge, tick infestation, coughing, lameness, emaciation, depression and salivation are recorded as the major AM findings of the current study. Accordingly, 47 (12.2%) sheep showed signs of diseases and abnormalities; of which 23 (6%) were conditionally approved whereas 11 (2.9%) sheep were unfit and judged to be detained and rejected. In the present study age, body conditions and geographic origin of the animals were considered as study variables, and the results showed BCS and age groups had statistically higher ( $p \le 0.05$ ) rejection probabilities. PM examination findings indicated a total of 192 lesions were encountered, of which 48.9%, 29.2%, 6.8%, and 5.7% lesions were recorded from livers, lungs, GIT, and hearts and kidneys, respectively. During the PM inspection, both total and partial condemnation judgments were passed on organs and carcass appeared with a sign of abnormality. C. teniculosis 58 (30.2%), calcification 36(18.7%), abscess 31(16.2%), hydatid cyst 23 (12%), hepatitis 10 (5.2%), pericarditis 4(2%), bruise 3(1.6%) and nephritis 2(1%) were found to be the major pathological lesions recorded. In two sheep all organs and carcass were totally condemned as their entire bodies appeared yellowish. The direct loss due to the condemnation of organs and carcasses at Jimma municipal abattoir was also investigated in slaughtered sheep, and there were about 56,576 USD losses per year. In conclusion, this study has identified the pathological lesions affecting edible organs and meat, and then rendering them unfit for human consumption. The study also estimated pathological lesions associated direct financial losses at Jimma abattoir. Therefore, further studies focusing on the primary causes of the abnormalities were recommended in the study area. Keywords: abattoir. financial loss. lesion. oraan

condemnation, PMI, sheep, jimma.

#### I. INTRODUCTION

he livestock sector globally is highly dynamic, contributes 40% of the global value of agricultural output and support the livelihoods and food security of almost a billion people (Thornton, 2010). Within African society, small ruminant comprises a greater proportion of the total wealth of the rural families, because of the low input requirements such as low initial capital, fewer resources and maintenance cost. They are also able to produce milk and meat in readily usable quantities using marginal lands and poor pasture and crop residues. Furthermore, their production cycle makes them need only short periods to reconstitute flocks after a disaster and respond quickly to the demand (Genet by, 1991). Ethiopia is the leading African country in livestock population, having around 34-40 million TLU (Tropical livestock unit) out of which 17% and 12% cattle and small ruminants, respectively, are found in Ethiopia (Ministry of information (MOI), 2005). According to (Development, ppStatis, 2009), the population of sheep and goats in Ethiopia is estimated to be 26.1 and 21.7 million respectively. It was the third largest number of sheep and goat among African nations and rank eighth in the world (Alemu and Merkel, 2008).

They generate cash income from export of meat. edible organs, skins and live animals (Ibrahim, 1998). There is also a high domestic meat demand from these animals, particularly during religious festivals. Even though this sub-sector contributes much to the national economy, its development is hampered by various constraints. These include endemic animal diseases, insufficient nutrition, poor husbandry, and lack of sufficient infrastructure, trained labor and government policies (PACE, 2003). Each year a large loss results from the death of animals and weight loss during transportation; and condemnation of edible organs and carcasses at slaughter.

Abattoir meat inspection is essential to remove gross abnormalities from meat and its products, to prevent the distribution of contaminated meat and to assist detecting and eradication of certain livestock diseases. More specifically, ante mortem inspection attempts to avoid introduction of clinically diseased animals into slaughter house and also serves to obtain information that will be useful in making sound post mortem inspection. Likewise, postmortem inspection is the center around which meat hygiene revolves since it provides information essential for evaluation of clinical signs and pathological process that affect the wholesomeness of meat (Herenda, *et al.*, 1994).

Author α σ ρ Ω: College of Veterinary Medicine, Haramaya University, P.O. Box 138, Dire Dawa, Ethiopia. e-mails: belina.timketa@gmail.com, dinaol.belina@haramaya.edu.et

As the meat is the sources of protein to a human being, it should be clean and free from diseases of particular importance to the public such as tuberculosis and cysticercosis. Meat is also condemned at slaughterhouse to break the chain of some zoo noses which are not transmitted to man directly via meat like hydatidosis and other important diseases of animals such as fasciolosis (Arbabi and Hooshyr, 2006; Fufaet *al.*, 2010).

Each year a significant economic loss results from mortality, poor weight gain, condemnation of edible organs and carcasses at slaughter. This production loss in the livestock industry is estimated at more than 900 million USD annually (Jacob, 1979; Abebe, 1995; Jobreet al., 1996). The major causes of pathological lesion during PMI of slaughtered ovine at abattoir are the disease caused by parasites, bacterial and other abnormalities. The final judgment as to action to be taken with an organ, the carcass or part of a carcass is based on the total evidence produced by the visual observation, palpation and incision (Teka, 1997).

Abattoir data is an important option for observing the diseases of both economic and public health importance (Arbabi and Hooshyr, 2006; Fufa, et al., 2010). Nowadays, several modern abattoirs like: HELMEX, ELFORA, Metehara, Modjo and Luna are established in Ethiopia. This increase in a number of slaughterhouse shows that increase in demand for meat supply, but the provisions have been challenging due to diseases, production problems and other factors. Given this, proper evaluation of financial losses due to organ condemnation resulting from various diseases at abattoirs is needed (Ezana, 2007). It is necessary to have enough information on a pathological lesion that causes organs and carcass condemnation at the abattoir. Hence, having information on where and how to reduce the losses that may be caused by the various abnormalities (lesions/pathology). Various studies (Jembere, 2002; Yimam, 2003; Aseffa, 2005; Get chew, 2008; Regessaet al., 2013) were carried out in the country in this regard to know the causes and losses associated. However, in Jimma there are no recorded studies conducted on major causes and financial losses associated with organs and carcass condemnation along with survey on pathological lesions. Therefore, the objectives of this study were to:

- Identify major pathological lesions causing organs and carcass condemnation in slaughtered sheep at Jimma municipal abattoir and,
- Estimate the direct financial losses attributed to condemned organs and carcass in sheep.

## II. MATERIALS AND METHODS

a) Study Area

The study was conducted from November 2016 to July 2017 at Jimma municipal abattoir in Jimma zone.

Jimma two is found in Oromia region south-western part of Ethiopia at a distance of 346 km away from Addis Ababa and lies between 36°50'E longitude and 7°40'N latitude atan average elevation of 1750 meter above sea level. Jimma is the largest city in south-western Ethiopia. It is special zone of the Oromia Regional state and is surrounded by different Jimma woreda. The climate of the area is characterized by humid tropical with bimodal heavy rainfall which is uniform in amount and distribution, ranging from 1200 to 2800 mm per year, with short and main seasons occurring from mid-February to May and June to September, respectively. The rainy season extends from mid-February to early October. Temperatures at Jimma are in a comfortable range, with the daily mean staying between 20°C and 25°C year-round. The total human populations of Jimma town was about 174, 446(88, 766 males and 85, 680 females). The livestock population of the area was reported to be about 2, 016, 823cattle, 942, 908 sheep, 288, 411 goats, 74, 574 horses, 49, 489 donkey, 28, 371 mules, 1, 139, 735 poultry and 418, 831bee hives (GOR, 2006).

#### b) Study Population

The study animals were sheep brought Jimma municipal abattoir and destined for slaughter. All animals were male and belonged to indigenous breeds kept under extensive management system. Sheep destined for slaughter had come from different parts of the weredas in the Jimma zone such as Dedo, Serbo, Saqa and Bilida inspected by standard AM, and PMI.

#### c) Study Design

A cross-sectional study using systemic random sampling technique was conducted from December 2016 to April 2017 to determine the pathological lesion that causes organs and carcass condemnation and to estimate the magnitude of direct financial loss attributed in sheep slaughtered at Jimma abattoir.

#### d) Sampling Method

## i. Sample size determination

In this study, systematic random sampling method was applied to include study animals, and study animals were grouped into young (under 1 years and three months) and adult above this based on the eruption of one or more incisor teeth according to Vattaet al. (2005).Since there was no published work on lesion survey from Jimma abattoir, 50% expected prevalence is considered to calculate the total sample size with 95% Cl, 5% level of precision (Thrush field, 2007). The sample size was 384 and determined using the formula given by

$$N = \frac{1.96^{2}P_{exp} (1-P_{exp})}{d^{2}}$$

Where N= required sample size,  $P_{exp}$ = expected prevalence and <sup>d</sup> is desired absolute

precision. Accordingly, the total sheep included in the study were 384.

#### e) Abattoir Survey

In the cross-sectional study of active abattoir survey, both AM and PMI were carried out in accordance by the procedures of Ethiopian Meat Inspection Regulation (Belina and Melese, 2017).

#### i. Ante-mortem Inspection (AMI)

The AMI (pre-slaughter examinations) of ovine was conducted at lairage both in motion and at rest and information related to study variables such as the behavior of an animal, age, BCS and origin and were recorded. At the same time, various signs of diseases and abnormalities were inspected with physical animal examination and its judgment were approved, conditionally approved, detained and rejected. Study animals were grouped into young and adult age groups according to standard dentation method (Vattaet *al.*, 2005).

#### ii. Post Mortem inspection (PMI)

During PMI all internal organs (liver, lungs, heart, kidney, gastro-intestinal tract), and carcasses were thoroughly inspected by visualization, palpation and making systematic incisions for the presence of cysts, parasites and other abnormalities. Pathological lesions were differentiated and judged according to guidelines on meat inspection for developing countries as totally fit for human consumption, and conditionally approved, and totally or partially condemned when unfit for human consumption (FAO, 1993).

#### f) Assessment of direct Financial Loss

In the current study, the total financial loss due to organs and carcass condemnation was computed by considering the condemnation rate or percentage of each edible organ and carcass, average number of animals slaughtered in the abattoir per year from retrospective data of the abattoir. The average weight of each organ and carcasses in kg, average current local market price of major organs and carcass, and each condemned organ was counted to estimate the financial loss. The average current local market price of each organ and mutton was collected by questionnaire from the butcheries in Jimma town for ease of computing the loss.

The retail average market prices obtained from butcher shops found in Jimma town in ETB were: Liver=30, lung=20, kidney=15, heart=18, GIT=90 ,whole carcass=4000 and mutton =150ETB/kg. In the case when there was whole carcass plus organs (whole body) rejection at PM, the average price of sheep came for slaughter was considered (4,000 ETB). The direct loss is calculated according to the procedures described by Ogurinade and Ogunrinade (1980), and the formula:

$$Los = MAK * PL\sum_{i=1}^{i=n} (PiCi)$$
, where

LOS is direct annual financial loss due to organs and carcass loss, MAK is annual average number of sheep slaughtered at Jimma abattoir, PL is overall prevalence of lesion, Pi is prevalence of each organ and carcass condemned, C<sub>i</sub> is average market price of each organ and 1kg mutton at butcher shops of the Jimma town.

The direct financial loss was expressed in US Dollar (\$) based on the current currency exchange rate of 1 USD = 22.5 Ethiopian Birr (ETB).

## g) Data Analysis

The active abattoir data, and questionnaire survey were entered into Microsoft Excel- 2016 spread sheet and the process of coding, cleaning and validating was done on this sheet and analyzed using SPSS version 20. For the data from PMI, descriptive statistics were used to determine organ and carcass condemnation rates, defined as the proportion of organs and carcasses condemned to the total number of organs and carcasses examined. Each financial loss was also calculated. Possible variation between rejection rates of specific organs, age groups and origin, were taken into consideration.

## III. Results

#### a) Abattoir Survey

## i. Ante mortem Inspection (AMI)

Detail AMI was conducted on a total of 384 sheep destined for slaughter at Jimma municipal abattoir and 47 (12.2%) of ovine were found to have different abnormalities. Nasal discharge, coughing, tick infestation, depression, emaciation, and lameness were those frequently observed among signs of diseases encountered in both age groups. The result also showed 6% (23/384) animals were conditionally passed for slaughter because of abnormalities such as lameness, respiratory problem, and their collection with tick infestation. On the other hand2% (8/384) sheep were unfit for human consumption and rejected during AMI. Since, they showed two and more signs of diseases such as emaciation with nasal discharge and depression1 %( 4/384), salivation and salivation with coughing 1 %( 4/384) were the major cause of rejection (Table 1).

The AMI result also depicted of 28 sheep with poor body conditions 46.43% were found to have one or more sign/s of illness whereas in those with good BCS

only 7% of the sheep were showed sign/s of diseases (Graph. 1)

In the current study, four different decisions were passed as AM judgments where apparently healthy sheep were passed for slaughter (91%), and others showed mild signoff illness and conditionally approved (6%) whereas 2.9% of sheep were detained and rejected as unfit for slaughter. Rejection rate was significantly higher ( $p \le 0.05$ ) in young animals with poor BCS than in adult animals with good and medium BCS (Table 2).

Table 1: Abnormalities encountered during AM inspection within Age groups and Origin of the animals

Ante-mortem	Age (%)		Origin (%)					
Finding	Adult	Young	Sub Total	Dedo	Bilida	Seka	Serbo	Sub Total
Apparently healthy	171(85.9)	166(89.7)	337(87.8)	127(88)	65(84)	99(89)	46(88.5)	337(87.8)
Depression	2(1)	1(0.5)	3(0.8)	2(1.4)	1(1.3)	0(0)	0(0)	3(0.78)
Salivation	1(0.5)	1(0.5)	2(0.5)	0(0)	1(1.3)	1(0.9)	0(0)	2(0.5)
Emaciation and nasal discharge	1(0.5)	1(0.5)	2(0.5)	1(0.7)	0(0)	0(0)	1(2)	2(0.5)
Tick infestation and coughing	2(1)	0(0)	2(0.5)	1(0.7)	0(0)	1(0.9)	0(0)	2(0.5)
Coughing	3(1.5)	2(1.1)	5(1.3)	0(0)	1(1.3)	4(3.6)	0(0)	5(1.3)
Lameness	3(1.5)	1(0.5)	4(1)	0(0)	2(2.6)	1(0.9)	0(0)	4(1)
Coughing and salivation	0(0)	2(1.1)	2(0.5)	0(0)	1(1.3)	1(0.9)	0(0)	2(0.5)
Tick infestation	5(2.5)	5(2.7)	10(2.6)	2(1.4)	3(4)	1(0.9)	3(5.8)	10(2.6)
Emaciation and depression	1(0.5)	1(0.5)	2(0.5)	2(1.4)	0(0)	0(0)	0(0)	2(0.5)
Nasal discharge	10(5)	4(2.2)	14(3.6)	8(5.6)	2(2.6)	2(1.8)	2(3.8)	14(3.6)
Tick infestation and lameness	0(0)	1(0.5)	1(0.3)	1(0.7)	0(0)	1(0.9)	0(0)	1(0.26)
Total	199	185	384	144	77	111	52	384



Graph 1: Proportion of ante-mortem finding by animals' boay condition score

Table 2 <sup>·</sup> Pro	portion of animals	rejected and pa	assed the AMI i	judaments by a	age aroup and F	SCS at the abattoirs
10010 2.110	portion of animula	rojooloa ana pa		adginorito by a	igo group ana L	oo al lito aballono

AMI	Age (%)		Body	Subtotal (%)		
judgment	Young	adult	Good	Medium	Poor	
Rejected	6(3.2)	2(1.0)	2(1.0)	0	6(21.4)	8(2)
Detained	0	4(2.0)	0	1(0.64)	3(10.7)	4(1)
Cond. Approved	8(4.3)	13(6.5)	10(5)	9(5.8)	4(14.3)	23(6)
Approved	171(92.4)	178(89.5)	188(94)	146(93.6)	15(53.6)	349(91)
Total	185	199	200	156	28	384
X(P valu	e)	0.05			0.00	

#### ii. Postmortem Inspection (PMI)

Among animals that had been examined during AMI 373 were slaughtered and subjected to through PMI following standard postmortem procedure and a total of192gross pathological lesion leading to partial and total condemnation of organs and carcasses were recorded. Among these abnormalities, lesions were frequently encountered from liver; and of which Cyst cercus teniculosis (34%) and calcification accounted 25.5%. These followed by abscess (13.8%), hepatitis (10.6%), cirrhosis (7.4%) and hydatid cyst (7.4%), fasciolosis (5.3%) and stelezia hepatica (3.2%).

A total of 56 lungs were also condemned as they were affected by teniculosis (35.7%), Hydatid cyst (21.4%), marbling lesion (17.8%), Emphysema and calcification (16%), pneumonia (12.5%), abscess (3.6%) and 5% with unidentified lesions. In this study abscessation was also inspected in other organs like heart, kidney and GIT, and carcass (Table 3). Out of a total of 11 hearts condemned (Table 4), hydatid cyst and pericarditis recorded as major causes contributing 45.5% and 36.4% which followed by abscess (18.2%) and discoloration (0.9%). Renal problems were observed in 11 kidney examined and 54.5% found to be caused by abscess whereas 27.3% and 18.2% were due to calcification and nephritis and other unidentified causes, respectively.13 GITs were also encounter as with abnormalities like cyst cercus teniculosis and abscess and foreign bodies, and they were subjected to total and partial condemnation accordingly.

The major pathological conditions for carcass rejection from local market were bruising accounting for 42.9%. Out of 7 rejections judgments2 were totaland the rest 5 were partial (Table: 3; and 4).

Table 3: Relative percentages of pathological lesions resulted in condemnations of organs and or carcasses
at the abattoir

Organ	Causes/ lesion		PMJ (%)			
affected		*AP	PC	TC	Total	
	Teniculosis	0	14(23)	2(1.8)	16(9)	
	Calcification	1(16.7)	12(19.8)	3(2.7)	16(9)	
	Discoloration	0	1(1.6)	3(2.7)	4(2.3)	
	Hydatid cyst	0	0	2(1.8)	2(1.12)	
	Fasciolosis	0	0	3(2.7)	3(1.7)	
Liver	Cirrhosis	0	0	7(6.3)	7(3.9)	0.00
	Abscess	0	1(1.6)	3(2.7)	4(2.3)	
	Hepatitis	0	0	7(6.3)	7(3.9)	
	Stelezia hepatica	0	0	3(2.7)	3(1.7)	
	Teniculosis and calcification	0	2(3.3)	0	2(1.12)	
	Teniculosis and abscess	0	1(1.6)	1(0.9)	2(1.12)	
	Teniculosis and hepatitis	0	0	1(0.9)	1(0.56)	
	Calcification and fasciolosis	0	0	2(1.8)	2(1.12)	
	Hepatitis and calcification	0	0	1(0.9)	1(0.56)	
	Abscess and hepatitis	0	0	1(0.9)	1(0.56)	
	Teniculosis and hydatid	0	0	1(0.9)	1(0.56)	
	More than three lesions	0	1(1.6)	2(1.8)	3(1.7)	
	Calcification	1(16.7)	3(4.9)	2(1.8)	6(3.4)	
	Teniculosis	0	3(4.9)	1(0.9)	4(2.3)	
	Marbling	0	3(4.9)	6(5.4)	9(5)	
	Pneumonia	0	0	5(4.5)	5(2.8)	
Lung	Hydatid cyst	0	0	5(4.5)	5(2.8)	
	Emphysema	0	1(1.6)	2(1.8)	3(1.7)	
	Abscess	0	`О ́	1(0.9)	1(0.56)	
	Emphysema and teniculosis	0	0	6(5.4)	6(3.4)	
	Marbling and teniculosis	0	0	1(0.9)	1(0.56)	
	Teniculosis and pneumonia	0	0	2(1.8)	2(1.12)	
	Hydatid and calcification	0	0	1(0.9)	1(0.56)	
	Unidentified Lesion	1(16.7)	0	2(1.8)	3(1.7)	

#### Survey on Pathological Lesion and its Financial Losses in Ovine Slaughtered at Jimma Municipal Abattoir, Jimma, Ethiopia

	Discoloration	0	0	1(0.9)	1(0.56)	0.00
Heart	Pericarditis	0	0	4(3.6)	4(2,3)	0.00
rioart	hydatid cyst	0	Õ	3(2.7)	3(1.7)	
	Abscess	0	1(1.6)	3(2.7)	4(2,3)	
Kidnev	Calcification	1(16.7)	1(1.6)	0	2(1 12)	
radinoy	Nenhritis	0	0(0)	2(1.8)	2(1.12)	
	Discoloration	0 0	0	1(0.9)	1(0.56)	
	Unidentified	0	Õ	2(1.8)	2(1 12)	
	Teniculosis	0	2(3,3)	1(0.9)	3(1.7)	
GIT	Foreign body	2(33,3)	2(3.3)	0	4(2.3)	
GIT	Abscess	2(00.0)	1(1.6)	2(1.8)	3(1.7)	
	Bruise	0	2(3.3)	0	2(1.12)	
	Abscess	0	2(3.3)	0	2(1.12)	
Carcass	Hydatid and Bruise	0	1(1.6)	0	1(0.56)	
04.0400	Discoloration	0	0	2(1.8)	2(1 12)	
liver and	Calcification	0	1(1.6)	0	1(0.56)	
kidnev	Abscess	0	0	2(1.8)	2(1 12)	
heart and	Hydatid cyst and abscess	0	0	1(0.9)	1(0.56)	
luna	i ijualia ojet ana abeeeee	Ū	Ū	.(0.0)	. (0.00)	
lung and	Hydatid cyst	0	0	1(0.9)	1(0.56)	
kidnev	Tydalla byot	Ŭ	Ũ	1(0.0)	1(0.00)	
raditoy	Calcification	0	1(1.6)	0	1(0.56)	
	Teniculosis	0	3(4.9)	2(1.8)	5(2.8)	
Liver and	Hydatid cyst	0	0	1(0.9)	1(0.56)	
Luna	Hydatid and calcification	0	0	1(0.9)	1(0.56)	
	Teniculosis and hydatid	0	0	2(1.8)	2(1.12)	
	Teniculosis	0	2(3.3)	0	2(1.12)	
Liver and GIT	Teniculosis and abscess	0	0	1(0.9)	1(0.56)	
Liver and GIT		0	0	1(0,0)	1(0,50)	
Liver and	Abscess	0	0	1(0.9)	1(0.56)	
heart	ADSCESS	U	0	1(0.9)	1(0.56)	
	Abagaga	0	0	O(1, 0)	0(1.10)	
Total	ADSCESS	U		∠(I.ŏ)	2(1.12)	179(100)
TOLAI			0(3.4) 01(	(99.0) I	11(99.9)	170(100)

\*AP=organs conditionally passed for human consumption

Table 4: Summary of direct financial losses and organs and carcass condemned at abattoir

Condemnation	Organ condemned						
status	Liver	GIT	Lung	Heart	Kidney	Carca	SS
PC	39	7	14	0	3	5(3kg)	68
TC	55	6	42	11	8	2*	124
Total	94	13	56	11	11	7	192
Price in US \$	99.33	38	43.556	8.8	6.33	370.8	

There was a condition of examining, a single to multiple lesions per organ e.g., we examined teniculosis and Calcification from a single liver; and all are recorded as different lesions.

\*=Whole carcass plus organs totally rejected at PM; except in carcass, PC indicates 50% loss.

#### b) Assessment of Direct Financial Losses

The direct financial loss was computed based on average cost/ price of individual condemned organs and carcasses during the study period, applying the formula given by Ogurinade and Ogunrinade (1980). The study indicated that there had been a total loss of 12,729,960ETB which is 56,576 Undue to a partial and total condemnation of organs and carcass at slaughterhouse annually. The study result also indicated there was a total condemnation of 2 (1.3%) whole carcasses (carcass plus organs) (Table 4). For the calculation, alive price of one sheep is considered as 4000ETB for total carcass condemnation.

## IV. DISCUSSIONS

The AM and PM inspections were conducted in the abattoir for the purpose of identifying an abnormality and removing animals' products with pathological lesions which were unsafe for human consumption and having poor aesthetic values (Van Llogtestijn, 1993; Grace yet al., 1999).

In this research, out of 384 study animals 11 (2.9%) were rejected and detained as unfit for human consumption suspecting different zoo notic diseases such as rabies in the case of salivation with high fever and tuberculosis in animals with a sign of emaciation with high coughing, depression and nasal discharge. Similarly, suspecting rabies in the case of salivation and bovine TB in animals with sign of emaciation with high depression and coughing was reported by Belina and Melees (2017). Radostits et al., (2000) stated that in domestic ruminants, cobalt deficiency results in appetence and loss of body weight, emaciation, weakness, decreased growth, unthrifty appearance, diarrhea, and anemia. Again during the AMI, 23(6%) of sheep were found to be showing signs of abnormalities such as lameness, tick infestation along with coughing and nasal discharge frequently encountered and passed with judgments of conditional approval where due attention was given for whole body part and specific organ at postmortem examinations.

One of the causes of lameness was trauma caused by hitting with a thick stick during driving to abattoir on foot and inappropriate vehicles and loading and off-loading negligence during transportation to marketplaces and to the abattoir. During the AM examinations, it was found that respiratory disorders were higher than other abnormalities encountered during the AMI 14(3.7%) nasal discharge and 5(1.3%) coughing. The respiratory signs such as the presence of nasal discharge and coughing were most probably related to stress due to lack of feed and water that may lead to immune suppression enhancing opportunistic pathogens. On the other hand, overcrowding during transportation is also a source of stress (Get chew, 2008). In agreement to the current study, coughing, depression and lameness are frequently observed abnormalities encountered during AMI (Man defroet al., 2015) at Elfora Export Abattoir, Ethiopia.

The rejection rate was significantly higher (p<0.05) for those poor body conditions than good and medium body conditions (Table 2). Because of poor BC by itself may be due to unidentified abnormalities that increase rejection probability. Jibatet *al.* (2008) studied and determined the rate of organs and carcasses condemned and the associated annual financial loss at HELMEX abattoir in Ethiopia and they reported out of 2688 sheep and goats examined 188(7%) carcasses were condemned due to poor body condition cases.

On the other hand, there was a significant difference (p=0.051) within the age groups of animals in rejection at AM more young than the adult which were 3.2% (6/185) and 1% (2/199) respectively. It may due to difficulty in protection from stress, shortage of feed and water, not getting enough rest. Herenda *et al.* (2000) stated that leanness (Poorness) is often observed in case of poor quality pasture and young growing animals which have had protein-deficient diet.

In the present study, organ condemnation rate showed that, liver and lung were the most frequently affected organs with the highest condemnation rate followed by GIT, kidney and heart and carcass significantly (p=00), which is 94(48.9%), 56(29.2%), 13(6.8%), 11(5.7%), 11(5.7%) and 7(3.6%) respectively. This finding is in agreement with reports of Cadmus and Adesokan (2009) who recorded that lungs (45.7%) and the liver (32.9%) were the most affected organs with the kidney (0.02%) and the heart (0.01%) being the least. The current study introduced that parasites are the major causes of organ condemnations. Parasitic causes

like, Cyst cercus teniculosis, hydatidosis, fasciolosis and Stelezia hepatica were found to be the major parasitic conditions responsible for organ condemnation. There was no statistical difference in the rate of organ and carcass condemnation from parasitic infestation considering the age and origin of animals. This shows that parasitic diseases of sheep are widely spread in all age groups and everywhere in the country.

The presence of small ruminant hydatidosis at slaughterhouse has been documented in Ethiopia. (Bekeleet *al.*, 1988) reported a prevalence rate of 16.4% in sheep which is higher than the finding in this study (7.4%). Similarly (Jobreet *al.*, 1996) reported prevalence rate of 11% and 6% from South Omo and DebreZeit slaughterhouses, respectively in sheep and goats.

In present finding, hydatid cysts were more frequently observed in lungs than liver of sheep (6.3%) and (3.6%) respectively. Additionally, similar findings also by different were reported authors (Khanet al., 2001, Dalimaiet al., 2002, Daryaniet al., 2007). However, the most common site for hydatid cyst was the liver followed by the lungs in the Middle East (Kamhawiet al., 1996). Lungs are most commonly affected by hydatidosis because at old age the liver capillaries are dilated, and most cysts passed directly to the lung. Secondly, the cyst passes to the lung via the thoracic duct without involving the liver (Gracey, 1986). And also, many researchers reported that liver and lung are the most commonly affected organs by hydatid cyst (Abunna and Hordofa, 2013; Denbarga, 2011; Jobre, 1996). The reason being that lung, and liver contain highest capillary bed in the body and therefore, the majority of the oncospheres were filtered out and trapped in the fine blood capillaries and only small number of oncospheres reaches the remaining organs (Gracey, 1986). In present study also lungs and liver 12(6.3%) and 7(3.6%) respectively, were affected by hydatidosis.

Out of 94 condemned liver teniculosis is the most frequent cause of organ lesion (34%) followed by calcification (25.5%), abscess 13(13.8%), hepatitis 10(10.6%), cirrhosis 7(7.4%), discoloration 4(4.3%), more than 3 lesion on liver 3(3.2%) and parasites like teniculosis, Stelezia hepatica, Fasciola species and hydatid cyst were found to be the major causes that rendered liver rejection from the local market (Table 3). Fascioliasis constitutes both economic and public health constrains to ruminant production. It is caused by two trematode species, Fasciola hepatica and F. gigantica, which develop in different livestock species mainly sheep and cattle; but, also in many other domestic herbivores (Gracey, and Collins, 1992). The reported prevalence of Fasciolaspp. (5.3%) was lower than other studies in bovine, like Belina and Melese (2017) study result showed fasciolosis and hydatidosis alone contributed 690(35.1%) gross pathological lesions.

Previous studies have indicated a higher economic loss resulting from a condemnation of edible organs and carcasses due to parasitic causes (Negategizeet al., 1993; Jembere, 2002; Jibat, 2006). In the current study, these parasitic causes of liver lesion might be due to improper wasting of condemned organ and the stray dog feed it at abattoir and selling of infected offal for dog which is final host for teniculosis and hydatidosis and stay them. Sissayet al., (2008) studied the prevalence and seasonal incidence of cestodeparasite infections of sheep in Eastern Ethiopia for two years (2003- 2005). Duringthis period, viscera including liver, lungs, heart, kidneys and the gastrointestinal tract were collected from 655 sheep slaughtered at four abattoirs. One of the most prevalent metacestodes was C. teniculosis. In sheep, the overall prevalence was 79% for C. teniculosis.

The causes for calcification abscess, hepatitis, cirrhosis and discoloration were difficult to identify grosslyand it may be due to systemic infectious diseases. Calcification is also another lesion that we encountered; it can be caused by injury, infection, and autoimmune disorders. Large-scale tissue damage is associated with extensive loss of cells, a situation referred to as tissue necrosis. The death of tissue in a specific area of the body leads to the release of signaling factors that attracts cells to clean up and heals the dead tissue. This process, known as an inflammatory response, attracts calcium into the damaged area as it heals (Carne, 2010).This study indicates 24(12.5%), 9(4.7%) and 3(1.6%) of Liver, lung, and kidney, respectively were affected by calcification.

Abscess was also a pathological condition; which is a collection of pus circumscribed by fibrous tissues. It occurs with great frequency throughout many organs and the carcasses of the meat animals and may be associated with a general condition or be found as isolated lesions (Libby, 1975). In present study 13(6.8%), 6(3.1%), 6(3.1%), 2(1%) and 2(1%) of liver, GIT, kidney, heart and carcass were affected by abscess. In agreement with (FSIS, 2009) stated that caseous lymphadenitis is a disease of sheep and goats caused by the C.Pseudotuberculosis. Postmortem findings may include, enlarged abscessed lymph nodes with greenish white-yellow caseous exudate, which tends to become dry and granular, cross-sections of lesions contain remnants of connective tissue capsules (resembles the concentric rings seen on the cut surface of an onion). Lesions found in many lymph nodes, especially the subiliac, superficial cervical, deep popliteal, tracheobronchial, and mediastinal lymph nodes, as well as lungs, heart, liver, spleen, and kidnevs. (Asrat. 2004) stated that occasionally the worms penetrate the bile duct wall into the liver parenchyma causing liver abscesses.

The study conducted in Gondar abattoir (Meseleet al., 2012) and Nekemte (Moje et al., 2014)

also revealed that livers and lungs are the most rejected organs by PM inspection and fasciolosis and hydatidosis are the major causes of rejections. However, in the current study different calcifications, cirrhosis, hepatitis, abscessations, emphysema, pneumonic lesions, marbling (contagious caprine pleuropneumonia (CCPP)) lesion, nephritis, foreign body, traumatic lesions and others nonparasitic abnormalities and unidentified lesion contributed to a condemnation of organs and carcasses were investigated.

Lungs were condemned because of C. teniculosis, hydatid cysts, marbling, emphysema, calcification-, pneumonia, other unknown caused lesion and abscess which were (35.7%), (21.4%),(17.8%),(16%), (16%), (12.5%), (5%) and (3.6%) respectively. C. teniculosis accounts for 35.7% as a principal cause of lung condemnation in sheep this might because of increased number of a stray dog in the area, the principal cause of lung condemnation was parasitic. However, the report observed during a retrospective study (Reassert al., 2013) reported pneumonia as a principal cause of lung condemnation in central Ethiopia accounting for 42.1% (Get chew, 2008).In current study, from the total lungs inspected higher 56 (29.2%) lungs were condemned. It may because of the animals unable to resist stress within a short period of time during transportation along way on foot, shortage of feed and water, stress due to hitting of animal by personnel who driving animal to market from the farmer and to abattoir and does not getting sufficient amount of rest at lair age may causes this respiratory problem.

FSIS (2009) reported that pneumonia is an inflammatory condition of the lungs that maybe caused by infectious agents, parasites, physical trauma, or foreign material in halation. In similar reports pneumonia might also be as a result of endemic diseases of sheep and goats such as pasteurellos are which is triggered by stress, contagious caprinepleuropneumonia (Radiostitis al., 2007). The other cause was marbled appearance (CCPP) lesions: CCPP is a disease peculiar to shoat and takes the form of a chronic inflammation of the lungs and pleura. It is not communicable to man and the carcass and the lungs found to be positive for CCPP can be passed for human consumption after a partial condemnation of the diseased part Gracey and Collins (1992).On the other hand, Emphysema is an abnormal and permanent enlargement of air spaces distal to terminal bronchioles with destruction of their alveolar walls, whereas oedema is a nonspecific lesion in which interstitium and alveoli are accumulated with fluid (Carne, 2010).

Also, different lesions of infectious and noninfectious causes like abscess, pericarditis, nephritis, and discoloration were found to be important causes for the condemnation of edible organs like liver, heart, and kidney. Similarly the same causes were found at central Ethiopia (Get chew, 2008; Regassaet al., 2013) and in goats slaughtered at Nigeria (Ojo, 1992).

In the present study, out of the 11 (5.7%) kidneys condemned abscess 6 (54.5%) account, whereas calcification accounts 3(27.2%), nephritis 2(18.2%) and other unidentified causes 2(18.2%) (Table 3).In this study abscess was a principal cause of kidney condemnation, however, the result in (Dejeneet al., 2013) study revealed out of the 57 (6.71%) kidneys condemned Nephritis 20 (2.35%) accounting for 11 (2.59%) and 9 (2.12%) kidneys in Ovine and Caprine respectively, was the principal cause of condemnation. Radostitiset al., (2000) stated that embolic nephritis occurs after septicemia or bacteremia when bacteria lodge in renal tissue.

The major causes of heart condemnation were found to be pericarditis, hydatid cyst and abscess. Out of the total of 11 (5.7%) hearts condemned due to gross abnormalities, pericarditis contributes about 4 (36.4%) and hydatid cyst also contribute 4(36.4%) and abscess 3(27.3%) out of condemned organs Table 3. The main cause of lesion in GIT condemnation primarily parasitic C.teniculosis and abscess 6(46.2%) and foreign body 4(30.8%). As a septic lesion, whenever localized abscess is found, partial condemnation is recommended Gracey and Collins (1992).

The main management practices that rendered organs and carcasses unfit for human consumption were bruising of the carcass mainly brought about by not proper handling of animals during transportation to the slaughterhouses by hitting the animal with thick stick and mechanical damage to organs due to faulty evisceration especially liver. Apart from affecting carcass value, bruising has also animal welfare implications as excessive use of sticks while driving to the abattoir, mishandling of animals during loading and unloading, improper transport vehicle and at slaughter could be responsible causes (Mungubeet al., 2006). It is stated that bruising of animals during transport is the major source of economic loss in Africa and Asia (Mitchell and slough, 1980). In the present study out of 7 carcass condemnations, 2 (28.6%) whole carcass was also totally condemned due to the yellowish discoloration, suspecting liver disease which may toxicity, systemic disease causing prehepatic and hepatic jaundice. Herendaet al. (2000) stated that icterus is the result of an abnormal accumulation of bile pigment, bilirubin, or of hemoglobin in the blood. Jaundice is divided into three main categories. Prehepatic jaundice occurs following an excessive destruction of red blood cells. Tick-borne diseases such as Babesiaovis and Anaplasmosis cause this type of icterus. Hepatic jaundice occurs due to direct damage to liver cells as seen in liver cirrhosis, systemic infections, and in chemical and plant poisoning. In sheep, jaundice may have been caused by phytogenic

chronic copper poisoning. Obstructive jaundice occurs when the drainage of the bile pigment bilirubin is blocked from entry into the intestine.

However, parasitic C. *teniculosis* and Stelezia hepatic have no public health importance; they are considered as the important cause of economic loss in the meat industry since viscera harboring them are rejected for aesthetic reasons. The threat these parasites pose to small ruminants' meat industry in Ethiopia is evident due to the present situation of improper disposal of offal at abattoirs and backyard slaughter. The presence of freely roaming stray dogs on grazing land together with livestock and the deeplyrooted habit of feeding dogs with offal, including sheep heads, are important risk factors. This may lead to the perpetuation of the life cycle between intermediate hosts (sheep) and the final hosts (dogs) for C. *teniculosis* and hydatidosis.

The financial loss in the abattoir was high, in this study analyzed those losses through condemnation of organs and carcass from local market. A total loss of(56,576 USD USD)was incurred in the abattoir. Carcass condemnation accounts highest part of the losses of the total direct losses whereas liver, lung, GIT, heart and kidney takes, respectively.

The indirect losses from body weight gain, mortality at the farms, public health implications (cause of treatment for a human when diseased upon eating of the affected edible organ which is zoonotic) were not included in the analysis in this study. Thus, the total financial loss attributable to diseases of ovine and. hence, abattoir wastage could be much higher. The economic analysis of livestock diseases in Ethiopia is scarce and inadequate because of lack of information on the prevalence and partly by the complexity of the analysis. Negategizeet al. (1993) have reported a financial loss associated with a liver condemnation due to ovine fasciolosis alone in the central highlands of Ethiopia amounting to be 2.3 million Ethiopian Birr (460, 000 USD). Similarly Jobreet al.(1996) have estimated a total annual loss of 1.3 million Ethiopian Birr (260 000 USD) resulting from offal condemnation and carcass weight loss

## V. CONCLUSION AND RECOMMENDATIONS

The current study revealed that, during the period different signs of diseases, and study abnormalities leading to conditional approval, rejection and detain of animals were encountered at AM inspection. In lesion survey, a total of 192 gross pathological lesions resulting in partial and total condemnations of liver, lung, kidney, heart, GIT and carcass were investigated. Different calcifications. parasitic teniculosis, hydatidosis, fasciolosis, Stelezia hepatica, pneumonic lesions, abscess, cirrhosis, marbling (CCPP), emphysema, hepatitis, nephritis, pericarditis, bruising (mechanical damage),

discoloration and foreign body were the main abnormalities recorded as causes of(56,576 USD) losses. There is perpetuation of the life cycle between intermediate hosts (sheep) and the final hosts (dogs) for C. *teniculosis* and hydatidosis by wasting the condemned organ near abattoir; and sometimes selling of affected organ. The results of the study showed that teniculosis and calcification were the two most frequently examined conditions, contributing 58(30.2%) of gross pathological lesions.

Based on this conclusion, the following recommendations are recommended:

- Awareness should be created for the animal attendants, farmers, customers, abattoir workers and butchers regarding the public health significance of diseases of animal origin and the related losses
- The government must empower veterinarians and other meat inspector more in passing professional judgments and, avoid complains of investors working in meat industry sector on inspectors judgments
- Immediate, safe and controlled elimination of all condemned abattoir materials and the sale of contaminated.

## References Références Referencias

- 1. Abebe, G. 1995.ILRI, Addis Ababa, pp: 133-138 *In:* Proceeding of an international conference of Current status of veterinary education and health research in Ethiopia In Veterinary Medicine impact on health and nutrition in Africa.
- 2. Abunna F., Hordofa D. 2013. Major Causes of Organ Condemnation for Cattle and its Financial Impact at WolaitaSoddo Municipality Abattoir, Southern Ethiopia .*Global Veterinaria*11: 730-734.
- 3. Alemu Y., MerkelR.C. (2008). Sheep and Goat Production Handbook for Ethiopia: Ethiopia Sheep and Goat Productivity Improvement Program/ ESGPIP/, Ministry of Agriculture and Rural: 15-53.
- 4. Arbabi M., Hooshyr H. 2006. Survey of Regions might have accounted for variation of the Echinococcosis and Hydatidosis in Kashan Region, Prevalence in different areas of a country Central Iran. *Iran J Public Health* 35: 75-81.
- Aseffa, M. 2005. Parasitic Causes of Carcass/Organ Condemnation at Asella Municipality Abattoir. Unpublished DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debrezeit, Ethiopia.
- 6. Asrat, M. 2004. Infection prevalence of ovine fasciolo is in irrigation schemes along the Upper Awash River Basin and effects of strategic an thelmintic treatment in selected upstream areas, Addis Ababa University School of Graduate Studies, Department of Biology. *M. Biomed. Sci.*

- 7. Bekele T., Mukassa M.E., Kasali O.B. 1988. The prevalence of cysticercosis and Hydatidosis in Ethiopian sheep. *Vet Parasitol.* 28: 267-270.
- Belina, D. and Melese, M. 2017. Pathological lesion survey and financial losses associated with organs and carcass condemnation in cattle slaughtered at selected abattoirs in Ethiopia. *Indian J. Vet. Pathol*, 41(1): 1-11.
- Cadmus, S.I.B., Adesokan, H.K. 2009.Causes and implications of bovine organs/offal condemnations in some abattoirs in Western Nigeria Trop. *Anim. Health Prod.*, http://dx.doi.org/ 10.1007/s11250-009-9334-7.
- Carne, H.R. 2010. Veterinary pathology hand book. Integrated Cased Based Applied Pathology. Vets 3011.
- 11. Dalimai A., Motamedi G.H., Hosseini M., Mohammad a B., Malaki H. 2002. Echinococcosis /hydatidosis in western Iran. *Vet Parasitol*. 105: 161-171.
- 12. Daryani A., Alaei R., Arab R., Sharif M., Dehghan M.H. 2007. The Prevalence, Intensity and Viability of Hdatidcysts in Slaughtered Animals in the Ardavil Province Northwest Iran's *Helminthol.* 81: 13-17.
- Dejene S., Abebe B., Degefu H. 2013. Study on the Major Health Problems That Causes Carcass and Organs Condemnation at Hashim's Export Abattoir, Debrezeit, Ethiopia *Global Veterinaria*11 (4): 362-371.
- 14. Denbarga Y., Demewez G., Sheferaw D. 2011. Major Causes of Organ Condemnation and Financial Significance of Cattle Slaughtered at Gondar Elfora Abattoir, Northern Ethiopia. *Global Veterinaria*7: 487-490.
- 15. Development, pp Statistical Agency (CSA), (2009). Central Statistical Authority Federal Democratic Republic of Ethiopia Agricultural Samples Enumeration Statistical Abstract, Pp 446-539.Central the Federal Democratic Republic of Ethiopia.
- Ethiopian Ministry of Agriculture Meat Inspection Regulations (EMAMIR), 1972. Legal Notice No. 428 NegariteGazexa. Addis Ababa, Ethiopia.
- 17. Ezana, G., 2007. Major disease of export oriented livestock in export abattoirs in /around adalibenwereda, Debreseit, Ethiopia.
- 18. FAO, 1993.Agro State Data. Statistical Division, www.fao.org/economic/ess publications/ ess-year book /en/ Rome, Italy.
- 19. FSIS, 2009.Public Veterinarian Training USDA FSIS Center for Learning Multi-speci Disposition Basics with a Public Health Focus1/29/09.
- 20. Fufa A., Loma A., Bekele M., Alemayehu R. 2010. Bovine fasciolosis: coprological, abattoir survey and its economic impact due to liver condemnation at Sodo Municipal abattoir, Southern Ethiopia. *Trop Anim Health Prod*.42: 289-292.

- Get chew, E.W. 2008. Major Diseases of Export Oriented Livestock in Export Abattoirs in/Around Ada Liben Woreda, Debrezeit. Online DVM Thesis, Faculty of Veterinary Medicine, Haramaya University, Ethiopia. http:// mahider. ilri.org/ handle / 10568/695.
- 22. Gatenby, R. M. 1991. Sheep. He tropical agriculturalist. MACMILLAN, ACCT. pp. 6-10.London and Basingstoke, education Ltd
- 23. GOR (Government of Oromia), 2006.Oromia reginal government socio-economic profile. Government of Oromia Region. Addis Ababa, Ethiopia.
- 24. Gracey J.F., Collins D.S. 1992.Meat Hygiene. 9th ed.Pp: 339-421.*ln:* Baillier Tindall. London.
- 25. Gracey, J.F. 1986.Meat Hygiene.8th ed. pp. 119-350.In: BaillierTindall. London.
- 26. Gracey J.F., Collins D.S. and Huey R.J. 1999.Meat hygiene. 10th Edition. pp: 758.*In:* London. W. B. Sounders Company Ltd.
- 27. Herenda D., Chambers P.G., Ettriqui A., Seneviratnap P. and Dasilvat T.J.P. 1994. Manual on meat Inspection for developing countries, pp: 139-148.
- Herenda D.; Chambers P.; Ettriqui A.; Seneviratna P. and da Silva T. 2000. Manual on meat inspection for developing countries. FAO *Anim. Prod. Health*, 119.
- Ibrahim, H., 1998. Small Ruminant Production Techniques. pp: 207. ILRI Manual 3. ILRI (International Livestock Research Institute), Nairobi, Kenya.
- Jacob, L., 1979. Seminar for animal health officials. Ministry of agriculture and settlement, Animals and Fisheries authority. Addis Ababa, Ethiopia
- Jembere, S., 2002: A Survey of Causes of Organs/Carcass Condemnation in Slaughtered Cattle at Nazareth Abattoir. Unpublished DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, DebreZeit
- Jibat, T., 2006. Causes of Organ and Carcass Condemnation in small ruminants slaughtered at HELMEX Abattoir. pp. 9–11. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debrezeit, Ethiopia.
- Jibat, T., Ejeta, G., Asafw, Y. and Wudie, A. 2008.Causes of abattoir condemnation in apparently healthy slaughtered sheep and goats at HELMEX abattoir, DebreZeit, Ethiopia, *Rev. Méd. Vét.*,159 (5): 305-31
- Jobre Y., Lebago F., Tiruneh R., Abebe G., Dorchies Ph. 1996. Hydatidosis in their selected regions in Ethiopia: an assessment trail on its prevalence, economic and public health importance. *Rev Med Vet* 147: 797-804.
- 35. Kamhawi S., Hijjawi N., Abu-Ghazaleh A., Abbas M. 1996. Prevalence of Hydatid Cyst in Livestock from

Five Regions in Jorda. *Ann Trop Med Hyg*.147: 797-804

- 36. Khan A.H., El–Buni A.A., Ali M.Y. 2001. Fertility of the Cyst of Echinococcus Granulosus in Domestic Herbiivorse from Benghazi.
- Libya, and the Ractivity of Antigens Produced from them.*Ann Trop Med Parasitol*.95: 337-342]\Libby, J.A., 1975. Meat Hygiene.4th ed. Pp. 40-168.*In*:Lea and Febiger, Philadelphia
- Mandefro A., Aragaw K., Hailu B., Alemayehu G., Chala G. 2015. Major Cause of Organ and Carcass Condemnation and Its Financial Loss at Bishoftu Elfora Export Abattoir. *International Journal of Nutrition and Food Sciences*.4: 364-372. doi: 10.11648/j.ijnfs.20150403.24
- 39. Mesele G., Guadu T., Bogale B., Chanie M. 2012. Pathological Conditions Causing Organ and Carcass Condemnation and Their Financial Losses in Cattle slaughtered in Gondar, Northwest Ethiopia. *Afr J Basic ApplSci*, 4: 200-2012
- 40. Ministry of information (MOI), 2005. Export products of Ethiopia. Press release of Ministry of information, department of press and Audio visual. Addis Ababa
- 41. Mitchell, J. R. and Slough, C.A.B. 1980. Helminthes in meat inspection.56-62.*In*: Guide to meat inspection in tropics: Common Wealth Agricultural Burro.
- 42. Moje N., Abdeta D., Kebede S., Terfa T., Desissa F., Regassa A. 2014. Major Causes of Organs and Carcass Condemnation in Cattle Slaughtered at Nekemte Municipality Abattoir, East Wollega, and Ethiopia. *Global Veterinaria*13: 278-284.
- 43. Mungube E.O., Bauni S. M., Tenhagen B. A, Wamae L. W., Nginyi J. M., Mugambi J. M.2006. The prevalence and economic significance of Fasciolagigantica and Stilesia hepatica in slaughtered animals in the semi-arid coastal Kenya. *Trop AnimHlth Prod*.3388:475-483.
- 44. Negategize P.K., Tekleye B. and Get chew J. 1993. Financial losses caused by ovine fasiolosis in Ethiopian highlands. *Trop. Anim. Hlth. Prod.*, 23: 155-167.
- 45. Negategize, P.K.J., Bekele, T., Tilahun, G., 1993. Financial losses caused by ovine fasciolosis in the Ethiopian highlands. *Trop. Anim. Health Prod*.25: 155–161.
- 46. Ojo, S., 1992.A Survey of Pathological Conditions in the Slaughter Goats. pp. 1–4. *In:* Zaria Slaughterhouse. Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria.
- PACE-Ethiopia, 2003. Experiences and the way forward on community Based animal health service delivery in Ethiopia. p. 6. *In:* Proceedings of workshop held in Addis Ababa, Ethiopia.
- 48. Radiostitis O.M., Gay C.C., Hincheliff K.W. and Constable, P.D. 2007. Veterinary Medicine. A text book of the disease of cattle, horse, sheep, pigs,

and goats.10ed ed. Pp 391-517. *In:* Ballier Tindals. London.

- Radostits O., Gay C., Blood D. and Hinchcliff K. (2000).Veterinary Medicine. A textbook of the diseases of cattle, sheep, pigs, goats and horses. 9<sup>th</sup> Ed. 1877pp. London: Harcourt, Publishers Ltd.
- Regassa A., Moje N., Megersaa B., Beyene D., Sheferaw D., Debela D., Abunnaa F., Skjerved E. 2013. Major causes of organs and carcass condemnation in small ruminants slaughtered at Luna Export Abattoir, Oromia Regional State, Ethiopia. *Preventive Veterinary Medicine*, 110: 139–148.
- 51. Sissay M., Uggla A. and Waller P. 2008. Prevalence and seasonal incidence of larval and adult cestode infections of sheep and goats in eastern Ethiopia. *Trop. Anim. Hlth. Prod.* 40(6):387-94.
- 52. Teka, G., 1997. Meat hygiene .pp: 99-113. *In:* Food hygiene principles and methods of food born disease control with special reference to Ethiopia.
- 53. Thornton, P., 2010. Livestock production: recent trends, future prospects. *Royal Soc.*, 365(1554): 2853-2867.
- 54. Hrushfield, M. 2007. Veterinary Epidemiology. ThirdEdition, Oxford. Pp 244-245, 249-251 and 258-259.*In*: Blackwell Science Ltd.
- 55. Van Iontestijn, J.G. 1993. Integrated quality. Meat safety: A new approach. *Meats Focus International*, 2:123-128.
- Vatta A.F., GumedeS.A. Harrison L.J.S., Krecek R.C., Letty B.A., MapeyiN. Pearson R.A. 2005. Goat keepers Veterinary Manual. Onderstepoort Veterinary Institute, Private Bag Onderstepoort, *Soutfrica. Veterinary Research*, 6: 42.
- 57. Yimam, M. 2003. Major Causes of Organ Condemnation in Ruminants Slaughtered at Gondar Abattoir, North West Ethiopia. Unpublished DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debrezeit.