

Analysis of Various Physiological Values of Growing Miniature Pigs in Isolator

Jin Soo Han

Received: 8 December 2017 Accepted: 2 January 2018 Published: 15 January 2018

Abstract

The objective of this study was to provide physiological values on the growing miniature pigs in isolator. Measurements were taken of each miniature pigs' body weight, temperature, hematological and serum biochemistries, and heart rate. The miniature pigs were divided into 2 groups. The piglet group ($n = 8$), consisting of animals approximately 4 weeks old and weighing 5 to 7 kg, were used to obtain normal physiological values at 6, 10, 14, 18, and 22 weeks of age. The adult group ($n = 8$), which consisted of animals approximately 40 weeks old weighing 19 to 21.3 kg, was used to obtain normal physiological values for miniature pigs at 43, 47, 51, 55, and 59 weeks of age. None of the animals displayed abnormal behavior or symptoms while they were bred in the isolators.

Index terms— isolator, miniature pigs, physiological values.

1 Introduction

Pigs have become widely used for biomedical research in recent years since there are many similarities in metabolic and cardiovascular function between swine and humans (Swindle et al., 1988). Miniature pigs specifically make good research animals, since they are easy to deliver, maintain, and utilize in isolators (Mandel and Travnicsek, 1987). Their small size relative to that of other large animals facilitates housing and handling, while their ample blood volume allows for more frequent serial blood collections than are possible with rodents. Furthermore, miniature pigs have received attention as potential donors in xenotransplantation because of their ability to be bred in an aseptic environment, their high fertility, and low costs (Park et al., 2006). Isolator systems are used to maintain laboratory and farm animals in a sterile environment to prevent contact-transmitted and airborne infections (Trexler, 1973). While isolators have been largely used in general laboratory animal research, most of the literature contains information pertaining to rodents, and there is little available information regarding optimal isolator design and conditions for miniature pigs.

The establishment of standard basic data is important in the biosciences, because it can reduce and refine the use of laboratory animals (Michael et al., 2006). Although there have been studies to acquire basic data regarding the use of miniature pigs, no data is available regarding their use within an isolator system. Therefore, the aim of this study, in addition to identifying baseline physiological values for miniature pigs growing within an isolator, was to identify optimal isolator designs and conditions for miniature pigs.

2 II.

3 Materials And Methods

4 P a) Advanced isolator

The miniature pigs were housed in groups of 2 in isolators (1200W × 900D × 950H m/m, SK-ISO-1700HBP600, Three-shine INC, Daejeon, Korea, Fig. 1). The internal temperatures of the isolators were set to $23 \pm 2^\circ\text{C}$, and their relative humidity was kept at $50 \pm 10\%$. The ventilation frequency was 50 times per h, illuminance was 100-150 Lux, and noise levels were kept below 58 phon, when possible. Feeding units and water nozzles were installed in the isolators, which consisted of space for breeding and movable roomdividing sub-frames set

on wheels. The sub-frames were equipped with a window for easy animal observation, and gloves for gnotobiotic handling of the animals. Gloves were protected from damage within the isolator by drawing them back into the glove housing after use, and a double-capped transport chamber was used to sterilize the goods coming in and out of the isolator. In short, the inner part of the cap was airtight, so that the outer cap could be removed to either add goods for sterilization and insertion, or extraction of sterilized waste.

The feed unit was securely fixed within the cage, and could be easily washed through a hole in its bottom. Automated watering valves were used to allow *ad libitum* consumption, and a cleaning box containing a water gun was attached for easy cleaning of the isolator's interiors. An uninterruptible power supply was installed to retain operation of the exhaust port in case of a blackout, ventilating the isolator via a safety filter to protect against the culling of miniature pigs midexperiment. The drain under the isolator was installed to make the excrement of the pigs fall downward, and the sub-drain switch valve can be used to remove the body wastes selectively.

5 c) Clinical symptoms

The health status of each miniature pig was monitored daily throughout the duration of the study. In case any abnormality was detected, a description of the severity and frequency of the symptoms was recorded by date.

6 d) Measurements of body weight and body temperature

The miniature pigs were weighed using a balance (HBS-510L; CAS, Korea). We measured the body surface temperature of the miniature pigs using a non-contact infrared thermometer, which allowed for measurement of their body temperatures without causing them any undue stress. Temperatures were measured on the center of the forehead, neck, lateral abdominal area and hip (eye exposure to the beam was avoided).

7 e) Hematological and serum biochemistry analysis

All pigs were anesthetized prior to blood collection using medetomidine 0.2 mg/kg (Domitor® ; Pfizer Korea, Seoul, Korea) intramuscularly and tiletamine-zolazepam 4.4 mg/kg (Zoletil® ; Virbac, Carros, France) intramuscularly. All miniature pigs were bled (5 mL) through their jugular veins. All blood samples were analyzed for evaluated using an automated hematology analyzer (FORCYTE; Oxford Science, USA) for CBC and differential blood cell counts. In addition to the red blood cell (RBC), hemoglobin concentration (Hb), and platelet (PLT) counts we did, we also quantified the content of several different cell populations within the white blood cell count (WBC), including: neutrophils (NEU), lymphocytes (LYM), monocytes (MONO), eosinophils (EOS) and basophils (BASO).

All serum samples were analyzed for complete total protein (TP), glucose (GLU), uric acid (URIC), calcium (CA), cholesterol (CHOL), total bilirubin (TBIL), creatine (CREA), alanine aminotransferase (ALT), albumin (ALB), aspartate aminotransferase (AST), and cortisol levels. All serum analysis was measured using a radioimmunoassay (RIA; PerkinElme, Finland) in the clinical laboratory of the Neodin Medical Institute (Seoul, Korea). The mean and SE were calculated for each of the measured parameters. Reference values for the hematology of pigs are documented well in previous literature.

8 f) Heart rate measurements

Heart rates of the 6-and 43-week-old animals were monitored and recorded. All pigs were anesthetized using medetomidine 0.2 mg/kg (Domitor® ; Pfizer Korea, Seoul, Korea) intramuscularly and tiletamine-zolazepam 4.4 mg/kg (Zoletil® ; Virbac, Carros, France) intramuscularly prior to collection of their heart rate data. During the 24 h of monitoring, we cleaned the isolators and fed the pigs while they were attached to the holter monitor for examination of their heart rates during their typical daily activities within the isolators. After 24 h, we anesthetized the animals once again, removed the monitors and then compiled the data.

9 g) Statistical analysis

All statistical analyses were performed using Graph Pad Prism version 4.0 for Windows (Graph Pad Software, San Diego, CA, USA). Data were recorded as mean $\pm$ standard error of the mean (SEM).

Comparisons of the two different groups were made by Student's unpaired t-tests and P values < 0.05 were considered significant.

10 Results

11 a) Clinical symptoms

All animals showed no clinical signs of disease and appeared healthy throughout the study.

12 b) Measurements of body weight and body temperature

The recorded body weight measurements are shown in Table 1. The weight of every animal that was bred in an isolator gradually increased. The results of the body temperature measurements are shown in lateral abdominal

area, and hip showed the highest temperature variations in all tests. The lowest temperature values occurred in the forehead region for all miniature pigs, and were lower in the 55-week-old miniature pigs (33.0°C) compared to other adult groups. The temperature of the abdominal region was higher in the 6-week-old miniature pigs (35.93°C) than in the other groups, and abdominal skin temperature values were considered to be very close to the rectal temperatures obtained of miniature pigs by using an infrared thermometer.

13 c) Hematological and serum biochemistry analysis

Hematological analysis results are shown in Tables 2 and 3. Total WBCs, neutrophil, and lymphocyte counts of the piglet group tended to decrease with age ($P < 0.05$). The total WBCs and neutrophil counts of the adult group tended to decrease with age, although data was not statistically significant between the age groups. The results of the serum biochemistry analysis are shown in Tables 4 and 5. Cortisol levels of the piglet group decreased with age. The cortisol values for the 6-week-old miniature pigs (18.04 ± 3.18 ug/dL) were higher than those for all other age groups. Although cortisol values of the piglet group tended to be greater than those of the adult group, the differences were not statistically significant. Serum biochemistry values did not differ among the age groups and were not statistically significant between the age groups. All hematological and serum biochemistry values were within the normal reference range. Bars represent the mean values and standard errors of body temperature ($n = 8$). One-way analysis of variance (ANOVA) tests revealed significant differences in 5 regions ($P < 0.05$); 1 Significant difference between the forehead and abdomen; 2 Significant difference between the forehead and rectum; 3 Significant difference between the neck and abdomen; 4 Significant difference between the neck and rectum; 5 Significant difference between the abdomen and hip; 6 Significant difference between the abdomen and rectum; 7 Significant difference between the hip and rectum.

Bars represent the mean values and standard errors of body temperature ($n = 8$). One-way analysis of variance (ANOVA) tests revealed significant differences in 5 regions ($P < 0.05$); 1 Significant difference between the forehead and abdomen; 2 Significant difference between the forehead and rectum; 3 Significant difference between the neck and abdomen; 4 Significant difference between the neck and rectum; 5 Significant difference between the abdomen and hip; 6 Significant difference between the abdomen and rectum;

14 Discussion

Pigs are exposed to many stress factors that may affect their health and welfare, including housing and management (Ruis et al., 1997). Typical signs of stress in pigs include depressed activity levels and decreased feed intake, which typically result in weight loss (Seymour et al., 1964; McGlone et al., 1987; Christon, 1988). None of the animals used in this study displayed abnormal behaviors or symptoms while they were bred in the isolators, and body weight measurements of all of the animals showed gradual body weight gains.

Temperature measurement is basic for medical evaluation, and is a tool that is frequently used in both clinical and research settings (Quimby et al., 2009). Body temperature is most commonly collected rectally in clinical settings, because it is an accurate reflection of core body temperature (Greer et al., 2007). This method is very difficult to utilize for animals housed within isolator environments, however, therefore we used a non-contact infrared thermometer. Since non-contact infrared thermometers typically require 2-4°C of calibration, the results should be considered when selecting a particular temperature region for experimental use with a non-contact infrared thermometer in miniature pigs. It is important to note that noninvasive temperature-measurement techniques that are rapidly accomplished could contribute to laboratory animal stress reduction and improved welfare (Chen et al., 2006). Results indicated that the areas exhibiting the highest temperatures in all tests were lateral abdominal area.

In order to measure the stress of breeding within an isolator, we monitored the cortisol levels in the animals' blood serum as well as the variation in their heart rates via a holter monitor. There was no significant difference between the average heart rates of 6- and 43-week-old animals. Although heart rates abruptly increased when there were workers outside of the isolators to clean or provide feed, this is likely due in response to cleaning noises and expectations of being fed. Additionally, while gradual increases and decreases in heart rates were observed during the animals' darkcycle sleeping hours, they were not regarded as abnormal symptoms of stress because reports indicate that heart rate also varies in humans during sleep (Bonnet and Arand, 1997).

Increased cortisol level is an important indicator of stress (Carroll et al., 2006), and serum concentrations have been widely used to assess the effects of different stressors on immune function (Bilandzi et al., 2006). Cortisol is known to be the primary glucocorticoid released during times of stress in pigs (Kojima et al., 2007), and administration of morphine and fentanyl during surgical procedures have been shown to decrease postoperative cortisol concentrations relative to controls in swine weighing 20 kg (Malavasi et al., 2006). The results indicate that cortisol levels were elevated in both the piglet and adult groups. Since transportation, weaning, and maternal separation have been shown to increase cortisol concentrations in pigs (Cooper et al., 2009; Kojima et al., 2008; Nyberg et al., 1988; ??arrot and Mission., 1989), the author attribute transportation stress to the increased cortisol levels in the 6- and 43-week old animals. Since cortisol levels were the highest in the 6-week-old animals, we assume that their young age caused them to be more susceptible to stress. The data agrees with previous studies, which indicated that increased stress resulted in elevation of WBC numbers and cortisol concentration (Morrow-Tesch et al., 1994; Kojima et al., 2009). These elevated WBC and cortisol concentrations

decreased with increasing age in all of this study’s miniature pigs. The author confirmed that all of the results from the CBC and serum biochemical tests were within the normal range. The data presented in this study provide a baseline for interpreting physiologic results of data gathered during the growth of miniature pigs within isolator systems.




Figure 1: Figure 1 :




Figure 2:

¹© 2018 Global Journals
²Significant difference between the hip and rectum.

3




Figure 3: Figure 3 :

4




Figure 4: Figure 4 :




Figure 5:

1

Figures 2 and 3. The center of the forehead, neck,

					Year
					2018
					31
d) Heart rate measurements					
Group	Age (weeks)		Body weight (kg)		
Piglet	6		6.06 ± 0.67		
	10		7.94 ± 0.57		
	14		9.35 ± 1.31		
	18 22 43 47		10.89 ± 2.48 12.24 ± 3.18 20.24 ± 0.67 21.68 ± 1.70		
Adult	51		23.56 ± 1.95		
	55		25.78 ± 2.69		
	59		27.40 ± 1.68		
Values are mean ± SE (n = 8).					

Figure 6: Table 1 :

2

	Normal Range	6 weeks	10 weeks	14 weeks	18 weeks	22 weeks
White blood cells (K/uL) 1	4.4-26.4	20.3 ± 6.55 1	14.28 ± 14.28	12.48 ± 4.50	10.94 ± 2.94	10.18 ± 3.54
Neutrophils (K/uL) 1	3.1-11.2	9.87 ± 5.95	5.07 ± 5.07	4.96 ± 1.70	4.98 ± 1.56	4.27 ± 1.66
Lymphocytes (K/uL) 1	4.3-13.6	8.85 ± 2.57	7.52 ± 7.52	6.31 ± 2.99	5.22 ± 1.39	4.84 ± 1.39
Monocytes (K/uL)	0.2-2.2	0.69 ± 0.36	0.69 ± 0.69	0.52 ± 0.15	0.36 ± 0.21	0.52 ± 0.37
Eosinophils (K/uL)	0.1-2.4	0.83 ± 0.74	0.98 ± 0.98	0.69 ± 0.82	0.35 ± 0.19	0.51 ± 0.76
Basophils (K/uL)	0.0-0.4	0.06 ± 0.03	0.04 ± 0.04	0.02 ± 0.01	0.02 ± 0.01	0.03 ± 0.03
Red blood cells (M/uL)	5.3-9.25	6.24 ± 1.51	6.69 ± 6.69	7.21 ± 1.00	5.73 ± 0.66	5.65 ± 1.38
Hemoglobin (g/dL)	9.0-15.8	9.51 ± 2.18	10.25 ± 10.25	10.53 ± 1.42	8.87 ± 1.00	8.73 ± 2.28
Platelets (K/uL)	148-898	374.06 ± 203.9	343.75 ± 343.8	370.13 ± 151.7	336.13 ± 90.0	289.00 ± 148.6

Figure 7: Table 2 :

3

	Normal Range	43 weeks	47 weeks	51 weeks	55 weeks	59 weeks
White blood cells (K/uL) 1	4.4-26.4	11.78 ± 3.95	11.23 ± 3.57	10.21 ± 1.39	10.11 ± 2.74	9.51 ± 3.17
Neutrophils (K/uL) 1	3.1-11.2	5.22 ± 2.01	4.82 ± 1.67	4.27 ± 1.12	4.03 ± 1.46	3.69 ± 1.53
Lymphocytes (K/uL)	4.3-13.6	5.46 ± 2.18	5.03 ± 2.45	5.21 ± 1.41	5.45 ± 1.17	4.97 ± 1.59
Monocytes (K/uL)	0.2-2.2	0.43 ± 0.23	0.56 ± 0.2	0.41 ± 0.20	0.38 ± 0.18	0.45 ± 0.21
Eosinophils (K/uL)	0.1-2.4	0.62 ± 0.52	0.79 ± 1.19	0.29 ± 0.28	0.24 ± 0.22	0.36 ± 0.45
Basophils (K/uL)	0.0-0.4	0.04 ± 0.03	0.04 ± 0.03	0.03 ± 0.04	0.02 ± 0.02	0.04 ± 0.02
Red blood cells (M/uL)	5.3-9.25	6.43 ± 0.57	6.80 ± 0.83	6.70 ± 1.61	6.29 ± 0.82	6.77 ± 1.86
Hemoglobin (g/dL)	9.0-15.8	9.91 ± 1.87	11.16 ± 1.88	10.42 ± 2.88	10.28 ± 1.72	10.46 ± 2.64
Platelets (K/uL)	148-898	391.9 ± 99.5	311.9 ± 90.7	284.2 ± 110.2	339.6 ± 94.2	323.8 ± 92.3

Figure 8: Table 3 :

4

	Normal 6 weeks			10 weeks		14 weeks		18 weeks		22 weeks	
	Range										
Total Protein(g/dL)	2.25 -	5.08	±	4.93	±	4.88	±	4.74	±	5.73	±
	8.0	1.75		1.97		1.37		1.48		1.25	
ALB (g/dL)	1.8 -	3.23	±	3.06	±	3.00	±	2.89	±	3.19	±
	3.3	0.83		0.77		0.37		0.73		0.52	
CREA (mg/dL)	0.5 -	0.59	±	0.63	±	0.71	±	0.71	±	0.85	±
	2.1	0.26		0.3		0.27		0.35		0.22	
URIC (mg/dL)		0.47	±	0.38	±	0.45	±	0.38	±	0.37	±
		0.34		0.38		0.32		0.38		0.4	
Glucose (mg/dL)	43	112.13	±	91.13	±	82.38	±	72.13	±	74.75	±
	-133	36.05		37.4		19.1		21.24		20.19	
TBIL (mg/dL)	0.0 -	0.08	±	0.08	±	0.08	±	0.08	±	0.08	±
	0.3	0.05		0.05		0.05		0.05		0.04	
AST (IU/L)	16	55.88	±	46.50	±	48.63	±	41.00	±	43.13	±
	-65	22.42		17.77		27.37		19.15		17.82	
ALT (IU/L)	9 -43	37.88	±	35.00	±	38.50	±	32.13	±	40.00	±
		11.53		10.2		17.57		16.69		16.24	
CHOL (mg/dL)	18	47.25	±	44.25	±	48.50	±	41.25	±	54.13	±
	-79	14.82		21.61		18.59		18.82		15.49	
Calcium (mg/dL)	6.5 -	9.98	±	8.70	±	8.55	±	7.98	±	8.95	±
	11.4	1.92		2.48		1.41		2.01		1.54	
Cortisol (ug/dL)		18.04	±	4.32	±	12.01	±	10.43	±	9.85	±
		3.18		3.81		5.13		5.03		5.73	

Values are mean $\pm$ SE (n = 8). ALB (albumin), CREA (creatine), URIC (uric acid), TBIL (total bilirubin), aminotransferase), ALT (alanine aminotransferase), CHOL (cholesterol). Normal range (Bollen et al., 2000;

Figure 9: Table 4 :

5

	Normal	43 weeks		47 weeks		51 weeks		55 weeks		59 weeks	
	Range										
Total Protein (g/dL)	2.25 -	6.16	±	5.73	±	6.45	±	6.19	±	6.75	±
	8.0	2.23		2.21		1.62		2.04		1.75	
ALB (g/dL)	1.8 -	2.93	±	2.81	±	3.31	±	3.14	±	3.18	±
	3.3	1.02		1.00		0.62		0.8		0.47	
CREA (mg/dL)	0.5 -	0.69	±	0.74	±	0.9	±	0.97	±	0.95	±
	2.1	0.37		0.32		0.25		0.28		0.1	
URIC (mg/dL)		0.38	±	0.47	±	0.33	±	0.35	±	0.3	±
		0.38		0.36		0.38		0.37		0.37	
Glucose (mg/dL)	43	91.25	±	69.38	±	90.88	±	79.38	±	89.13	±
	-133	55.74		42.39		39.44		45.06		46.56	
TBIL (mg/dL)	0.0 -	0.09	±	0.18	±	0.08	±	0.1	±	0.13	±
	0.3	0.06		0.3		0.05		0.05		0.05	
AST (IU/L)	16	60.5	±	48.88	±	39.38	±	34.75	±	48.38	±
	-65	28.25		19.05		12.98		14.95		25.65	
ALT (IU/L)	9 -43	27.38	±	30.50	±	28.63	±	26.63	±	31.25	±
		7.93		13.85		6.46		10.51		14.06	
CHOL (mg/dL)	18	43.63	±	46.25	±	47.75	±	48.25	±	54.25	±
	-79	21.79		20.42		23.56		20.73		24.21	
Calcium (mg/dL)	6.5 -	8.98	±	8.16	±	9.08	±	8.86	±	9.41	±
	11.4	2.34		2.37		1.6		2.31		1.8	
Cortisol (ug/dL)		9.99	±	9.79	±	8.2	±	9.85	±	9.74	±
		2.33		3.93		3.03		2.81		6.41	

Values are mean $\pm$ SE ($n = 8$). ALB (albumin), CREA (creatinine), URIC (uric acid), TBIL (total bilirubin), aminotransferase), ALT (alanine aminotransferase), CHOL (cholesterol). Normal range (Bollen et al., 2000;

Figure 10: Table 5 :

6

Year 2018		
34		
Age	6 weeks	43 weeks
Minimum	84.2 $\pm$ 10.4	68.6 $\pm$ 9.9
Average	121.9 $\pm$ 16.8*	111.7 $\pm$ 12.9
Maximum	186.6 $\pm$ 24.0	191.0 $\pm$ 10.7

[Note: Values are mean $\pm$ SE ($n = 8$). *Significant ($P < 0.05$) differences between values for 6 and 43 weeks. (Student's unpaired t -test, two-tailed)]

Figure 11: Table 6 :

-
- [Trexler ()] *An isolator system for the maintenance of aseptic environments*, P Trexler . 1973. p. .
- [Parrott and Mission ()] 'Changes in pig salivary cortisol in response to transport stimulation, food and water deprivation and mixing'. R F Parrott , B Mission . *Br Vet J* 1989. 145 p. 501.
- [Park et al. ()] 'Comparative analysis of various blood chemical values between miniature pigs and conventional pigs'. S H Park , D Y Kim , B K Park , H S Yoo , H J Youn , H Han . *Lab Anim Res* 2006. 22 (1) p. .
- [Greer et al. ()] 'Comparison of 3 methods of temperature measurement in hypothermic, euthermic, and hyperthermic dogs'. R J Greer , L A Cohn , J R Dodam , C C Wagner-Mann , F Mann . *J Am Vet Med Assoc* 2007. 230 p. .
- [Quimby et al. ()] 'Comparison of digital rectal and microchip transponder thermometry in cats'. J M Quimby , F Olea-Popelka , M Lappin . *J Am Assoc Lab Anim Sci* 2009. 48 p. .
- [Chen and White ()] 'Comparison of rectal, microchip transponder, and infrared thermometry techniques for obtaining body temperature in the laboratory rabbit'. P H Chen , C White . *J Am Assoc Lab Anim Sci* 2006. 45 p. .
- [Bilandzi? et al. ()] 'Cortisol and immune measures in boars exposed to three-day administration of exogenous adrenocorticotrophic hormone'. N Bilandzi? , M Zuri? , M Lojki? , B Simi? , D Mili? , I Bara? . *Vet Res Commun* 2006. 30 (3) p. .
- [Seymour et al. ()] 'Effects of dietary protein level and environmental temperature on performance and carcass quality of growing-finishing swine'. E W Seymour , V C Speer , V W Hays , D W Mangold , T Hanzen . *J Anim Sci* 1964. 23 p. 375.
- [Malavasi et al. ()] 'Effects of epidural morphine and transdermal fentanyl analgesia on physiology and behaviour after abdominal surgery in pigs'. L M Malavasi , G Nyman , H Augustsson , M Jacobson , M Jensen-Waern . *Lab Anim* 2006. 40 p. .
- [Mcglone et al. ()] 'Effects of heat and social stressors and within pen weight variation on young pig performance and agonistic behavior'. J J Mcglone , W F Stansbury , L Tribble . *J Anim Sci* 1987. 65 p. 456.
- [Kojima et al. ()] 'Effects of syndyphalin-33 on feed intake and circulating measures of growth hormone, cortisol, and immune cell populations in the recently weaned pig'. C J Kojima , S J Jenkins , T A Cooper , M P Roberts , J A Carroll , H Kattesh . *J Anim Sci* 2009. 87 p. .
- [Nyberg et al. ()] 'Effects of transport stress on concentrations of cortisol, corticosteroid-binding globulin and glucocorticoid receptors in pigs with different halothane genotypes'. L K Nyberg , I Lundstrom , Edfors-Lilja , M Rundgren . *J Anim Sci* 1988. 66 p. 1201.
- [Cooper et al. ()] 'Effects of transport stress, gender, and weaning weight on post-weaning performance in pigs'. T A Cooper , M P Roberts , H G Kattesh , C Kojima . *Prof. Anim Sci* 2009. 25 p. .
- [Kojima et al. ()] 'Effects of weaning and weaning weight on neuroendocrine regulators of feed intake in pigs'. C J Kojima , J A Carroll , R L Matteri , K J Touchette , G Allee . *J Anim Sci* 2007. 85 (9) p. .
- [Bonnet ()] 'Heart rate variability: sleep stage, time of night, and arousal influences'. M H Bonnet , Arand D L . *Electroencephalogr. Clin. Neurophysiol* 1997. 102 (5) p. .
- [Morrow-Tesch et al. ()] 'Heat and social stress effects on pig leukocyte numbers, mitogen-induced lymphocyte proliferation, antibody synthesis and plasma immunoglobulin G concentration'. J L Morrow-Tesch , J J Mcglone , J L Salak , R Hurst . *J Anim Sci* 1994. 72 p. 2599.
- [Carroll et al. ()] 'Hormonal profiles, behavioral responses, and short-term growth performance after castration of pigs at three, six, nine, or twelve days of age'. J A Carroll , E L Berg , T A Strauch , M P Roberts , H G Kattesh . *J Anim Sci* 2006. 84 (5) p. .
- [Kojima et al. ()] 'Physiological and immunological responses to weaning and transport in the young pig: Modulation by administration of porcine somatotropin'. C J Kojima , H G Kattesh , M P Roberts , T Sun . *J Anim Sci* 2008. 86 p. .
- [Swindle et al. ()] 'Swine as model in experimental surgery'. M M Swindle , A C Smith , Bjs Hepburn . *J Invest Surg* 1988. 1 (1) p. .
- [Ruis et al. ()] 'The circadian rhythm of salivary cortisol in growing pigs: effects of age, gender, and stress'. Maw Ruis , Te Brake , Jha Engel , B Ekkel , E D Buist , W G Blokhuis , H J Koolhaas , J . *Physiol Behav* 1997. 62 (3) p. .
- [Christon ()] 'The effect of tropical ambient on growth and metabolism in pigs'. R Christon . *J Anim Sci* 1988. 66 p. 3112.
- [Mandel and Travnicek ()] 'The minipig as a model in gnotobiology'. L Mandel , J Travnicek . *Nahrung* 1987. 31 p. .
- [Michael et al. ()] 'The principles of weight of evidence validation of test methods and testing strategies'. B Michael , Patric A Susanne , B Silvia , C Sandra , C Richard , C Robert , C Raffaella , C Rodger , C Chantra , E , Julia F Laura , G Marlies , H Thomas , H Sebastian , H Leonard , S , Laurie S Horst , S William , S Raymond , T Drew , W Valérie , Z . *Altern Lab Anim* 2006. 34 (6) p. .