

Location of the Axillary Nerve in Relation to Arm Length: A Cadaveric Study in a Kenyan Adult Population

R. Oluoch¹, H. R. Mwangi², A. Njoroge³ and G. Y. Elbadawi⁴

¹ Moi University

Received: 12 December 2018 Accepted: 31 December 2018 Published: 15 January 2019

Abstract

Background: The location of the axillary nerve from various bony landmarks has been described in the literature. Such descriptions can assist the surgeon to avoid iatrogenic injury to this nerve during deltoid splitting approaches to the shoulder and proximal humerus. **Objective:** To investigate the relationship between the axillary nerve and arm length in a cadaveric adult Kenyan population. **Methods:** The distance of the axillary nerve from important bony landmarks such as the anterior and posterior edges of the acromion, and the lateral epicondyle of the humerus were measured and recorded.

Index terms— axillary nerve, arm length.

Abstract-Background: The location of the axillary nerve from various bony landmarks has been described in the literature. Such descriptions can assist the surgeon to avoid iatrogenic injury to this nerve during deltoid splitting approaches to the shoulder and proximal humerus.

Objective: To investigate the relationship between the axillary nerve and arm length in a cadaveric adult Kenyan population.

reported to be around 10% of all brachial plexus injuries (4). Trans-deltoid incisions or approaches to the proximal humerus and the shoulder joint such as open reduction internal fixation (ORIF) of proximal humerus fractures, and open rotator cuff repair places this nerve at increased risk of injury either through transection or traction (3).

Traditionally, the axillary nerve is described to lie about 5cm from the edge of the acromial process of the scapula. This measurement, however, has individual variability and differs from study to study (3,5,6).

1 II.

2 Materials and Methods

This study conducted at the Human Anatomy laboratory of Moi University after ethical clearance from the Institutional Research and Ethics Committee (I.R.E.C) of Moi University. Fifty-one formalin prefixed left adult upper extremities disarticulated at the scapulohoracic junction were used. Only the left limbs were used since these were the majority.

Dissections were done using the deltopectoral approach to demonstrate the origin, course, and distribution of the axillary nerve beneath the deltoid muscle. A digital calibrated caliper [(Neiko® Tools Digital Caliper serial number 0.3.04.0487ECC) accurate to 0.01mm] was used to measure individual arm lengths and the lengths of the axillary nerve.

Hypodermic needles were introduced through the axillary nerve. This was to help in the representation of the course of the nerve on the outer surface of deltoid muscle and to minimize errors during subsequent measurements.

The distance from the acromion lateral edge to the lateral epicondyle of the humerus was measured and recorded as the arm length (AL). The course of the axillary nerve was represented on the outer surface of the deltoid muscle using hypodermic needles. A digital caliper was used to measure the distance from the anterior

43 edge of the acromion (AEA) to the axillary nerve and recorded as the anterior distance (AD) as illustrated in
44 figure 1 below.

45 3 Introduction

46 The axillary nerve originates from the posterior cord of the brachial plexus. It then descends inferolaterally on the
47 surface of subscapularis muscle to end in the quadrangular space by dividing into its two terminal divisions -the
48 anterior and posterior (1,2). The nerve supplies the shoulder joint, deltoid and the teres minor muscles.

49 Injury to this nerve can cause significant functional impairment such as limitation in shoulder abduction (3,4).
50 Axillary nerve injuries have been The distance from the posterior edge of the acromion (PEA) to the axillary nerve
51 was measured and recorded as the posterior distance (PD). All measurements taken recorded in centimeters (cm).
52 Correlation analysis was then performed between arm length (AL), and the anterior and posterior distances.

53 The ratio between arm length (AL) and the anterior distance (AD) was calculated for each cadaver and
54 recorded as the anterior index (AI) which is the distance of the nerve from the anterior edge of the acromion
55 divided by arm length, i.e. $AI = \frac{AD}{AL}$. Also, the ratio between arm length (AL) and the posterior
56 distance (PD) was calculated and recorded as the posterior index (PI) which is the distance of the nerve from
57 the posterior edge of the acromion divided by arm length, i.

58 4 Results

59 The average anterior distance (AD) was 6.46cm (SD 0.7cm) with a range of 5.15cm -8.68cm. The average
60 posterior distance (PD) was 5.88cm (SD 0.95cm) with a range of 4.42cm -9.99cm. There was a significant
61 correlation between AD and AL ($r = .335$; $p = 0.016$), where AD is expected to significantly ($p = 0.016$) increase by
62 0.104cm for every 1cm increase in AL.

63 The correlation between PD and AL (posterior index, PI) was not statistically significant ($r = 0.147$; $p = 0.304$).
64 A regression model fitted on the same shows that PD is predicted to increase by 0.062cm for every 1cm increase
65 in AL. However, this change is not statistically significant ($p = 0.304$). The average AI and PI were 0.2 ± 0.022 cm
66 (range 0.157-0.253cm) and 0.185 ± 0.033 (range 0.14-0.35cm) respectively.

67 5 IV.

68 6 Discussion

69 From the present study, the average anterior distance (AD) was 6.46cm (range 5.15-8.68cm) while the average
70 posterior distance (PD) was 5.88cm (range 4.42-9.99cm). The arm length had a mean of 31.96 cm (range 27.29-
71 38.74cm). Statistical analysis of these measurements revealed that for the AI, for every 1cm increment in the
72 arm length, the AD was predicted to increase by 0.104cm ($p = 0.016$). This increment represented a significant
73 change. While for the PI, for every 1cm increase in the arm length, the PD was predicted to increase by 0.062.
74 This data did not represent a significant change ($p = 0.304$).

75 The findings above are similar to those described by Cetik et al., (2006) on 24 embalmed adult cadaveric
76 shoulders (3). The authors reported that the axillary nerve curved inferior to the anterior and posterior edges
77 of the acromion process at an average distance of 6.08cm and 4.87cm respectively. They found a significant
78 correlation between arm length and both the anterior ($r = 0.79$, $p < 0.001$) and posterior distances ($r = 0.61$, p
79 $= 0.001$) respectively.

80 In a cadaveric study of 30 shoulders by Abhinav et al., (2008), the distance of the axillary nerve from the
81 lateral edge of the acromion was measured in varying degrees of adduction and abduction of the shoulder joint.
82 The average arm length (AL) was 31.0cm (range 27-34.5cm), with the axillary nerve located at an average length
83 of 6.0cm (range 4.5-6.5cm) from the lateral edge of the acromion. This distance reduced significantly when the
84 arm in an abducted position. Abduction moved the nerve closer to the acromion thereby putting the nerve at
85 risk during surgery (6). The present study did not consider the nerve distances with abduction or adduction
86 due to the stiff nature of the cadaveric specimens. In the current study, the average anterior (AI) and posterior
87 indices (PI) were 0.20 (range 0.157-0.253) and 0.185 (range 0.14-0.35) respectively. These findings are similar to
88 those reported by Cetik et al., (2006) who found figures of 0.20 and 0.16 for AI and PI respectively (3).

89 From the present study, a quadrangular safe area or zone which is located above the axillary nerve can be
90 useful and safe during deltoid splitting incisions in proximal humerus fractures, intramuscular injections, and
91 shoulder arthroscopy. This quadrangular shaped area is similar to that described by Cetik et al., (2006) as shown
92 in figure 3 Finally, the use of deltoid ratio, i.e. length and width of the deltoid muscle versus the distance from
93 the acromion to the axillary nerve can be used to predict the location of the axillary nerve preoperatively (8).
94 However, from the present study, measuring the deltoid length and width intraoperatively is a difficult task for
95 the surgeon. The surgeon is advised to use easily palpable bony landmarks, i.e. the acromion process of the
96 scapula (both the anterior and posterior edges), and the lateral epicondyle of the humerus to easily predict the
97 course of the axillary nerve.

98 V.


99 7 Conclusion

100 A 1cm rise in arm length had a predictable increase in nerve distances from anterior and posterior edges of
101 acromion by 0.104 cm and 0.062 cm respectively.

102 A preoperative template of a quadrangular "safe zone/ area" as landmarks on the proximal deltoid muscle
103 using minimum distances of 5.15cm and 4.42cm from anterior and posterior edges of acromion process of the
104 scapula respectively should protect the axillary nerve and its branches during surgery. ^{1 2}

¹© 2019 Global Journals

²© 2019 Global Journals Location of the Axillary Nerve in Relation to Arm Length: A Cadaveric Study in a Kenyan Adult Population



1

Figure 1: Figure 1 :





Figure 2:



2

Figure 3: Figure 2 :

Figure 4:

1

Number	Parameter	Range (cm)	Mean \pm SD, n=51
1	AD	5.15 -8.68	6.46 (0.70)
2	PD	4.42 -9.99	5.88 (0.95)
3	AL	27.29 -38.74	31.96 (2.27)

The arm length (AL) had a mean of 31.96cm (SD 2.27cm) with a range of 27.29cm -38.74cm as shown below.

Figure 5: Table 1 :

105 .1 Conflict of Interest

106 The author(s) declare (s) that there is no conflict of interests regarding the publication of this paper.

107 [Abhinav et al. ()] *A contribution to the calculation of a safe deltoid split. International journal of shoulder*
108 *surgery*, G Abhinav , B Sivaraman , N Matthew , Jst Grahame . 2008. 2 p. .

109 [Liu et al. ()] *Anatomic study of the axillary nerve in a Chinese cadaveric population: correlation of the course*
110 *of the nerve with proximal humeral fixation with intramedullary nail or external skeletal fixation. Archives of*
111 *orthopaedic and trauma surgery*, K Y Liu , T H Chen , J F Shyu , S T Wang , J Y Liu , P Chou . 2011. 131
112 p. .

113 [Duparc et al. ()] ‘Anatomical basis of the variable aspects of injuries of the axillary nerve (excluding the terminal
114 branches in the deltoid muscle)’. F Duparc , G Bocquet , J Simonet , P Freger . *Surgical and radiologic*
115 *anatomy: SRA* 1997. 19 (3) p. .

116 [Gray et al. ()] *Gray’s anatomy: the anatomical basis of clinical practice*, H Gray , S Standring , H Ellis , Bkb
117 Berkovitz . 2005. New York: Elsevier Churchill Livingstone. p. 1627. (39th ed. Edinburgh)

118 [Cetik et al. ()] ‘Is there a safe area for the axillary nerve in the deltoid muscle? A cadaveric study’. O Cetik , M
119 Uslu , H I Acar , A Comert , I Tekdemir , H Cift . *The Journal of bone and joint surgery American volume*
120 2006. 88 (11) p. .

121 [Sinnatamby ()] *Lasts Anatomy -Regional and Applied*, C Sinnatamby . 2011. Livingstone: Elsevier. 12.

122 [Visser et al. ()] ‘The incidence of nerve injury in anterior dislocation of the shoulder and its influence on
123 functional recovery. A prospective clinical and EMG study’. C P Visser , L N Coene , R Brand , D L
124 Tavy . *The Journal of bone and joint surgery British volume* 1999. 81 (4) p. .

125 [Kontakis et al. ()] ‘The position of the axillary nerve in the deltoid muscle. A cadaveric study’. G M Kontakis ,
126 K Steriopoulos , J Damilakis , E Michalodimitrakis . *Acta Orthop Scand* 1999. 70 (1) p. .