

Genital Mycoplasmas among Women in the Province of Basrah with an Evaluation of their Role in Some Cases

Reham M. Al-Mosawi

Received: 7 December 2018 Accepted: 5 January 2019 Published: 15 January 2019

Abstract

A total of 120 women attending the outpatient clinic of the obstetric and gynecology department of Basrah General Hospital, were evaluated for the role of genital mycoplasmas in certain infectious female genital conditions during the period from December, 1st 2003 to October, 2nd 2004. High vagina and endocervix swabs were cultured. Five species of genital mycoplasmas were isolated and identified on modified PPLO medium, the characteristics and biochemical properties of them were fit with those of *Mycoplasma hominis*, *Ureaplasma urealyticum*, *M.fermentans*, *M.genitalium* and *M.penetans* in which the last three species were identified for the first time in Iraq. *M.hominis*, *U.urealyticum*, *M.genitalium* and *M.penetans* were more frequently distributed in age group 20-39 years, and in women with lower socioeconomic status. A significant difference was found in the isolation of both *M.hominis* and *M.genitalium* from non-pregnant. Also, a significant difference was noted in isolation rates of *M.hominis*, *M.fermentans* and *M.penetans* from pregnant women at different gestational periods. A statistically significant difference at the level of ($P<0.01$) was found in isolation of both *U.urealyticum* and *M.genitalium* from endocervix region.

Index terms—

1 Introduction

he mycoplasmas are the smallest cell free-life microorganisms, characterized by their small cell size (0.3-0.8 μ m) and thus can pass through some filters used to remove bacteria. They have the smallest genome size and, as a result, lack many metabolic pathways and lack of a rigid bacterial cell wall (Mayer and Murray et al., 2002). Mycoplasmas are fastidious organisms that require specialized media for their isolation and identification (Razin and Freundt, 1984; Azinet al., 1998). Ultrastructural studies showed that they are constructed of only three organelles: plasma membrane, ribosomes and prokaryotic chromosomes. There is no evidence of any intracellular membrane structure, and because of the lack of rigid cell wall, mycoplasmas are pleomorphic (Lin, 1985). The family Mycoplasmataceae contains two genera that infect humans: *Mycoplasma* and *Ureaplasma*, which are usually referred to collectively as mycoplasmas. Seven species of mycoplasmas can be isolated from genitourinary tract, but in female genitourinary tract the most common mycoplasmas isolated are *U. urealyticum* and *M.hominis* have been implicated in human diseases (Cassell and Cole, 1981; Krause and Taylor-Robinson, 1992). *M.hominis* and *U.urealyticum* normally inhabit the urogenital tract in sexually mature men and women. Higher isolation rates are obtained from women than from men, possibly reflecting more favorable growth conditions in the vaginal tract. The degree of colonization with these species is related to sexual activity and to the number of sexual partners (Taylor-Robinson and McCormack, 1979). In 1980 new mycoplasma species were isolated from urethral specimens from 2 of 13 men with nongonococcal urethritis. It was later named *Mycoplasma genitalium* because of the host tissue location, however *M.genitalium* strains were originally isolated from the urogenital tract. This mycoplasma shared several properties with *M.pneumoniae* (Tully et al., 1981(Tully et al., 1983;Lind et al., 1984;Taylor-Robinson, 1995).

M.genitalium and *M.penetans* are recent additions to the pathogenic human mycoplasmas, reviewed by Marmion and Harris, (1996). *M.penetans*. This newly-identified mycoplasma was isolated by (Lo U et al

7 D) CULTIVATION AND ISOLATION OF MYCOPLASMAS (GENITAL MYCOPLASMAS)

., 1991) from the urogenital tract of Human Immuno Virus of positive homosexual men. In a study in west Africa, *M.genitalium* was associated with nongonococcal urethritis (NGU), particularly in *Trichomonas vaginalis* negative patients (Pepin et al., 2001).

Preliminary evidence for the involvement of the mycoplasma in the cervicitis and pelvic inflammatory disease was presented by Moller et al. (1984) and Uno et al. (1997). The etiological roles of genital mycoplasmas are: Acute pyelonephritis, bacterial vaginitis, pelvic inflammatory disease, chorioamnionitis, post-abortion and postpartum fever, pneumonia in new borns, non gonococcal urethritis, prostatitis and epididymitis (Mardh and Westrom, 1970;Shepard, 1970;Shepard, , 1980; Taylor-Robinson and McCormack, 1980; Krause and Taylor-Robinson, 1992;Taylor-Robinson, 1996). Furthermore, that mycoplasmas play a role in human infertility (Gnarpe and Fribreg, 1972;Stray-Pedersen et al., 1978; Asselletal., 1983; Othetal., 1983), abortion and stillbirth (Embreeetal., 1980; Kundsins et al., 1981), premature birth and low birth weight of infant (Braun et al., 1971; mbreeetal., 1980;Kundsins et al., 1981).

Despite numerous studies of these organisms, their role in most genital infection remains controversial and illdefined. The ubiquity and low virulence of genital mycoplasmas make it difficult to evaluate their role in producing genital infections, a situation that is further complicated by their frequent isolation together with other sexually transmitted diseases (Watts and Eschenbach, 1988). Furthermore the presence of both genital mycoplasmas (*U.urealyticum* and *M.hominis*) in a large proportion of healthy women complicates the assessment of the pathogenic role of these organisms (Taylor-Robinson and Furr, 1998). However, circumstantial evidence including high frequency of isolation of mycoplasmas from females with urogenital conditions or infertility warns against minimizing the importance of these microorganisms in human disease (Kundsins and Driscoll, 1970).

Mycoplasmas were the causative agents of disease in human (Krause and Taylor-Robinson, 1992) over many years passed. Despite, numerous studies conducted in Iraq on different genera of bacteria, there have been few studies on the role of genital mycoplasmas in the infectious disease (Simhairi, 1990;Al-Bahli, 1993).

2 II.

3 Materials and Methods

4 a) Study population

The population under study was women attending the outpatient clinic of obstetric and gynecology department of Basrah General Hospital. The samples obtained consisted of pregnant women at different gestational periods and non pregnant women including women using contraceptive devices besides women not using contraceptive devices and those with infertility and those with various complaints. Also women presented with different types of abortions are included. The samples were selected randomly from the women above and collected from different parts of Basrah. A total number of two hundred and twenty two women (222) were investigated. Their age ranged from 17 to 50 years. The investigation period extended from 1 st of December, 2003 to 2 nd of October, 2004. A control group consisted of one hundred (100) healthy women.

5 b) Collection and Inoculation of samples

Two swabs one from endocervix and the other from high vagina were obtained from each woman and each was inoculated. Onto a suitable medium. A sterile speculum was used. All swabs were transported to the laboratory within (one) hour for culture. For the isolation of mycoplasmas, each specimen was directly inoculated into the liquid phase, mixed up well and tilted for a while, once or twice, to cover the upper slanted portion in a simple monophasic -diphasic culture setup (MDCS) prior to incubation (Al-Sulamietal., 2002). For the isolation of bacteria other than mycoplasmas another two swabs from endocervix and high vagina were obtained from the same women included in the study. Then, each specimen after being transported to the laboratory was directly cultured onto MacConkey & Blood agar by the streaking method then incubation follows.

6 c) Culture Media

Media for isolation and identification species of genital mycoplasmas were applied depending on Marmion and Harris [5].

7 d) Cultivation and isolation of mycoplasmas (genital mycoplasmas)

Endocervix and high vaginal swabs were taken from the women under study, then, each specimen was directly inoculated into the liquid phase of the MDCS, mixed up well and then tilted once or twice to cover the upper portion of the slant for a while prior to incubation. All inoculated media were incubated aerobically at 37C o and observed daily for colour change from red to yellow in the liquid phase after 24 hrs. Then isolated colonies appeared after that on the slanted solid phase. We added sheep erythrocytes (7%) to the solid phase of the MDCS in several trials to detect the blood haemolysis by *M.fermentans* and *U.urealyticum*. as well, the egg yolk suspension (15 ml) was added to the standard PPLO agar for detect the lipolytic ability of *M.fermentans*.

8 III.

9 Results

In our knowledge this study performs the first isolation of *M.fermentans*, *M.genitalium* and *M.penetans* in Iraq with MDCS system and modified PPLO media according to the previous studies. Besides, it is the first isolation to both species: *M.hominis* and *U.urealyticum* by this method. The colonial growth was observed on the upper portion of the slant together with changing in colour of the liquid phase from red to yellow after 24 hrs. However, approximately 96 hrs. Period was necessary for full development of colonies on modified media of PPLO with MDCS for show the fried-egg appearance Figure 1 educational level and being frequently isolated from women with a low level ($p<0.01$; $p<0.05$; $p<0.05$) respectively. A difference was found to be significant in the isolation of both *M.hominis* and *M.genitalium* from non-pregnant when compared with pregnant women. A significant difference was also noted in the isolation rates of: *M.hominis*, *M.fermentans* and *M.penetans* from pregnant women at different gestational periods (trimesters) as shown in Table 3, 4. A statistically significant difference at the level of ($P<0.01$) was noted in the isolation of both *U.urealyticum* and *M.genitalium* from endocervix region in comparison with high vaginal region. This study revealed a significant association between the five species of genital mycoplasmas and some cases or symptoms in the female genital tract such as: vaginal discharge, itching and dysuria (Table 5, 6). Also, the genital mycoplasmas were found as a single infection in 20 cases (16.9%) and mixed infection with other causative agents in 21 cases (17.5%) Table 7. IV.

10 Discussion

Prevalence of mycoplasmas in the female genital tract depends on numerous factors such as age, level of socioeconomic status, sexually active, disorders in the menstruation, pregnancy, infertility, urogenital complaints and use of contraceptive device [19-21, Reham Al-Mosawi, 2009]. According to our knowledge, this study was the third in Iraq to determine the prevalence of genital mycoplasmas in Iraqi women and the second in Basrah. Further, the current study shows for the first time the isolation and identification of: *M.fermentans*, *M.genitalium* and *M.penetans* in Iraq besides the first isolation of both *M.hominis* and *U.urealyticum* by using modified PPLO medium and MDCS system for the primary isolation of genital mycoplasmas from clinical samples then, detection with biochemical test [18, Holt et al., 1994]. The modified PPLO medium consist of antimicrobial agents: such as thallium acetate and penicillin, the other supplements consist of glucose solution; sodium deoxyribonucleate (DNA calf thymus) solution; hydrogen phosphate solution and cresol red solution as an indicator of the mycoplasmal growth in addition to PPLO broth/agar media; yeast extract; sodium chloride and horse serum, this method offered multi merit symbolized by the less contamination as a result of elimination of transport media, fast of results appearance, supply a pliability in the of media represented by liquid and solid phases, it is inexpensive because its consumed a small quantities of liquid and solid media furthermore, it has a short incubation time period within one test tube. It is the first technique used to determine the prevalence of genital mycoplasmas in women in Basrah City in comparison with studies of Simhairi (1990 and Al-Bahli (1993).

Table 1 shows, the four species of genital mycoplasmas: *M.hominis*, *U.urealyticum*, *M.genitalium* and *M.penetans* were more frequently isolated in age groups 20-29 and 30-39 years whereas, *M.fermentans* tend to be frequently distributed in age group less than 20 years ($p<0.01$), this association could probably due to increase the sexual activity during this age of Iraqi women's life [21,26]. Concerning the educational levels as apparent in Table 2, the species: *M.hominis*, *M.fermentans* and *M.genitalium* were significantly related to the educational level and being frequently isolated from women with a low level. This is because the colonization of genital mycoplasmas in human is linked to younger age, lower socioeconomic status, and sexual activity with multiple partners and other factors [19,21]. A difference was found to be significant in the isolation of both *M.hominis* and *M.genitalium* from non-pregnant when compared with pregnant women. A significant difference was also noted in the isolation rates of: *M.hominis*, *M.fermentans* and *M.penetans* from pregnant women at different gestational periods (trimesters) as shown in Table 3, 4. Simhairi (1990), Al-Bahli (1993) found no significant difference in the prevalence of *M.hominis* and *U.urealyticum* among pregnant women as compared to non pregnant, as well as, to gestational stage of pregnancy. Their results were somewhat consistent with the results of the study. Similar as well as contrary observations regarding the prevalence of *M.hominis* of other strains in pregnant women have been reported by many investigators (Csonka et al., 1966; Jones, 1967; Harwick et al., 1970; Delouvois et al., 1975; McCormack, 1979 and Waska et al., 1986a). This controversy may be attributed to several factors like strain variations of the microorganisms, difficulty in determining a precise matching control with respect to sexual experience, especially in relation to the number of partners and population peculiarities, like differences in the socioeconomic status and hygienic standard. In present study we found, a statistically significant difference at the level of ($P<0.01$) in the isolation of both *U.urealyticum* and *M.genitalium* from endocervix region in comparison with high vaginal region (Table 5), the results of present study in agreements with R. M. Al-Mosawi, 2009. And in study of Upadhyaya et al. (1983) the frequency of the isolation of *U.urealyticum* was significantly higher in the infertile group than in a group of pregnant women. Thereupon, the results of the current study which show that both *U.urealyticum* and *M.genitalium* were recovered in a higher isolation rates from endocervix than high vagina of infected women may be associated with the inflammatory disease leading to infertility. Several investigators have demonstrated some kind of association between female infertility and *U.urealyticum* infection (Gnarpe and Friberg, 1973a; Fowlkers et al., 1975; Upadhyaya et al., 1983; Taylor-Robinson, 1986).

10 DISCUSSION

This study revealed a significant association between the five species of genital mycoplasmas and some cases or symptoms in the female genital tract such as: vaginal discharge, itching and dysuria (Table6)my study in accordance, with R. M. Al-Mosawi, 2009 except of the species *M.genitalium* which tend to be frequently distributed in women complaining of urethral discharge followed by vaginal discharge and itching also, Simhairi (1990) found that, the conditions like vaginal discharge, itching and monilia-associated infection were more frequent among patients with *M.hominis* infection than among those having no such infection. This reflects a significant degree of association between *M.hominis* infection and those conditions.

Similar results were reported by several investigators (Bercovici et al., 1962;Mendel et al., 1970;Taylor-Robinson and McCormack, 1979) who showed a high isolation rate of *M. hominis* in females with monilial and other vaginal conditions. Also, in the current study, the genital mycoplasmas were found as a single infection in 20 cases (16.9%) and mixed infection with other causative agents in 21 cases (17.5%) as showed in Table 7, these results were in agreement with R. M. Al-Mosawi, 2009.




Figure 1: Figure 1 :




Figure 2: Figure 2 :

Volume XIX Issue I Version I
 D D D D)
 (
 Medical Research
 Global Journal of

Figure 3:

1

Age year	No. of tested women	No. and (%) of women + ve in				
		M.hominis	M.fermentans	U.urealyticum	M.genitalium	M.penetrans
<20	10	1 (10.0)	1 (10.0)	0 (0)	0 (0)	0 (0)
20 -29	50	7 (14.0)	3 (6.0)	6 (12.0)	2 (4.0)	2 (4.0)
30 -39	40	6 (15.0)	1 (2.5)	4 (10.0)	3 (7.5)	1 (2.5)
? 40	20	2 (10.0)	0 (0)	2 (10.0)	0 (0)	0 (0)
Total	120	16	5	12	5	3
	X 2	1.694	7.361	11	13.08	7.192
	P	NS	0.01	0.01	0.01	0.01

Figure 4: Table 1 :

2

Level of education	No. of tested women	No. and (%) of women + ve in				
		M.hominis	M.fermentans	U.urealyticum	M.genitalium	M.penetrans
LOW	65	11 (16.9)	4 (6.1)	7 (10.7)	4 (6.1)	3 (6.4)
Moderate	35	4 (11.4)	1 (2.8)	3 (8.5)	1 (2.8)	0 (0)
High	20	1 (5.0)	0 (0)	2 (10.0)	0 (0)	0 (0)
Total	120	16	5	12	5	3
X 2		7.533	6.231	0.675	6.231	3.112
P		0.01	0.05	NS	0.05	NS

Figure 5: Table 2 :

3

Women sub-jects	No. tested	No. and (%) of women + ve in				
		M.hominis	M.fermentans	U.urealyticum	M.genitalium	M.penetrans
Pregnant	40	3 (7.5)	1 (2.5)	4 (10.0)	0 (0)	2 (5.0)
Non-pregnant	50	8 (16.0)	2 (4.0)	5 (10.0)	2 (4.0)	1 (2.0)
X 2		12.040	0.346	0	4.000	2.25
P		0.01	NS	NS	0.05	NS

Figure 6: Table 3 :

4

Pregnant women	No. tested	No. and (%) of women + ve in					Total of G.M (%)
		M.hominis	M.fermentans	U.urealyticum	M.genitalium	M.penetrans	
First trimester	10	1 (10.0)	1 (10.0)	1 (10.0)	0 (0)	1 (10.0)	4 (40.0)
Second trimester	10	0 (0)	0 (0)	1 (10.0)	0 (0)	0 (0)	1 (10.0)
Third trimester	20	2 (10.0)	0 (0)	2 (10.0)	0 (0)	1 (5.0)	5 (25.0)
X 2		10.110	23.333	0	0	10.000	34.000
P		0.01	0.01	NS	NS	0.01	0.01

Figure 7: Table 4 :

5

Sources of isolation	No. and (%) of women + ve in				
	M.hominis	M.fermentans	U.urealyticum	M.genitalium	M.penetans
Endocervix	7 (43.7)	2 (40.0)	7 (58.3)	3 (60.0)	1 (33.3)
High vagina	9 (56.2)	3 (60.0)	5 (41.6)	2 (40.0)	2 (66.6)
X 2	6.75	20.0	17.33	20.0	36.11
P	0.01	0.01	0.01	0.01	0.01

Figure 8: Table 5 :

6

Uorgenital complaints	No. of tested women	No. and (%) of women + ve in				
		M.hominis	M.fermentans	U.urealyticum	M.genitalium	M.penetans
Vaginal discharge	60	9 (15.0)	4 (6.6)	7 (11.6)	4 (6.6)	2 (3.3)
Dysuria	20	2 (10.0)	0 (0)	1 (5.0)	0 (0)	0 (0)
Itching	40	5 (12.5)	1 (2.5)	4 (10.0)	1 (2.5)	1 (2.5)
Total	120	16	5	12	5	3
X 2		4.23	8.36	5.66	4.88	4.25
P		0.05	0.01	0.01	0.05	0.05

Figure 9: Table 6 :

7

Total of genital mycoplasmas	Alone	S.aureus	In conjunction with K.pneumoniae	P.aeruginosa	E.coli	Proteus. spp
41	20	7	5	4	3	2

Figure 10: Table 7 :

173 [Williams Wilkins, C. O. Baltimore London] , *Bergey's Manual of Systematic Bacteriology* Williams & Wilkins,
174 C. O. Baltimore London (ed.) 1.

175 [Japan. Int. J. STD-AIDS. University Gifu, Japan] , *Japan. Int. J. STD-AIDS. University Gifu, Japan* 8 (4) p.
176 .

177 [Sc] , M Sc . Al-Mustansiriya University. (Arabic

178 [Mardh and Westrom (]) , P A Mardh , L T Westrom . 1970.

179 [Tully et al. (]) 'A newly discovered mycoplasma in the human urogenital tract'. J G Tully , D Taylor-Robinson
180 , R M Cole , D L Rose . *Lancet* 1981. p. .

181 [Al-Sulami et al. (]) 'A single method for rapid isolation and identification of *Mycoplasma pneumoniae* from
182 clinical specimens'. A A Al-Sulami , S S Hammadi , G J Al-Gizawi . *Eastern Mediterranean Health Journal*
183 2002. 8 (1) p. . (Arabic))

184 [Lilia et al. (]) 'Association of *Mycoplasma hominis* and *Ureaplasma urealyticum* with some indicators of
185 nonspecific vaginitis'. C R Lilia , G Constantino , Z Ivonne , Y Antonio , G Silvia . *Revista Latinoamericana*
186 *de Microbiologia* 2000. 42 p. .

187 [Braun et al. (]) 'Birth weight and genital mycoplasmas in pregnancy'. P Braun , Y H Lee , J O Klein , S M
188 Marcy , T A Klein , D Charles , P Levy , E H Kass . *N. Engl. J. Med* 1971. 284 p. .

189 [McCormack et al. (]) 'Colonization with genital *Mycoplasma* in women'. W M McCormack , B Rosner , Y H Lee
190 . *Am. J. Epidemiol* 1973b. 97 p. .

191 [McCormack et al. (]) 'Colonization with genital *Mycoplasma* in women'. W M McCormack , B Rosner , Y H Lee
192 . *Am. J. Epidemiol* 1973b. 97 p. .

193 [Shepard (]) 'Current status of *Ureaplasma urealyticum* in nongonococcal urethritis'. M C Shepard . *Asean. J.*
194 *Clin. Sci* 1980. 1 p. .

195 [Lo et al. (]) 'Enhancement of HIV-1 cytotoxic effects in CD 4 + lymphocytes by the AIDS-associated my-
196 coplasma'. S C Lo , S Tsai , J R Benish , J W Shih , D J Wear , D M Wong . *Science* 1991. 251 p.
197 .

198 [McCormack (]) 'Epidemiology of *Mycoplasma hominis*'. W M McCormack . *Sexually Transmitted Diseases. J.*
199 *Amer. Ven. Dis. Assoc* 1983. 10 p. .

200 [Pepin et al. (]) 'Etiology of urethral discharge in West Africa: the role of *Mycoplasma genitalium* and
201 *Trichomonas vaginalis*'. J Pepin , F Sobela , S Deslandes . *Bull Wld. Hlth Organ* 2001. 79 p. .

202 [Taylor-Robinson (]) 'Evaluation of the role of *Ureaplasma urealyticum* in infertility'. D Taylor-Robinson .
203 *Pediatric infect. Dis* 1986. 5 p. .

204 [Iwasaka et al. (]) 'Hormonal status and mycoplasma colonization in the female genital tract'. T Iwasaka , T
205 Wada , Y Kidera , H Sugimori . *Obstet. Gynecol* 1986a. 86 p. .

206 [Lin (]) 'Human macoplasma infection: Serologic observations'. J L Lin . *Rev. Infect. Dis* 1985. 2 p. .

207 [Taylor-Robinson (]) 'Infections due to species of *Mycoplasma* and *Ureaplasma*: anupdate'. D Taylor-Robinson .
208 *Clin. Infect. Dis* 1996. 23 p. .

209 [De-Louvois et al. (]) 'Microbial flora of the lower genital tract during pregnancy: relationship to morbidity'. J
210 De-Louvois , R Hurley , V C Stanley . *J. Clin. Pathol* 1975. 28 p. .

211 [Cassell et al. (]) 'Microbiologic study of infertile women at the time of diagnostic laparoscopy: association of
212 *Ureaplasma urealyticum* with defin sub population'. G M Cassell , J B Younger , M B Brown , R E Blackwell
213 , J K Davis , P A Marriott , S Stagno . *N. Engl. J. Med* 1983. 308 p. .

214 [Simhairi (]) *Microbiological study of some mycoplasma species of female genitalia and evaluation of their role in*
215 *some disease conditions*, R Z Simhairi . 1990.

216 [Razin et al. (]) 'Molecular biology and pathogenicity of mycoplasmas. Microbiol'. S Razin , D Yogev , Y Naot .
217 *Mol. Biol Rev* 1998. 62 p. .

218 [Bercovici et al. (]) 'Mycoplasma (pleuro pneumonia-like organisms) invaginitis'. B Bercovici , S Persky , R
219 Rozansky , S Razin . *Am. J. Obstet. Gynecol* 1962. 84 p. .

220 [Gnarpe and Friberg (]) 'Mycoplasma and human reproductive failure: The occurrence of different mycoplasma
221 in couples with reproductive failure'. H Gnarpe , J Friberg . *Am. J. Obstet. Gynecol* 1973a. 114 p. .

222 [Mayer and Murray (]) 'Mycoplasma and Ureaplasma'. G Mayer , G M Murray . MBIM 650/720. *Medical*
223 *Microbiology* 2002. 3. South Carolina University

224 [Tully et al. (]) 'Mycoplasma genitalium, a new species from the human urogenital tract'. J G Tully , D Taylor-
225 Robinson , D L Rose , R M Cole , J M Bove . *Int. J. Syst. Bacteriol* 1983. 33 p. .

226 [Harwick et al. (]) 'Mycoplasma hominis and abortion'. H J Harwick , R H Purcell , J B Lippa , F R Felkety .
227 *J. Infect. Dis* 1970. 121 p. .

- [Jones ()] 'Mycoplasma hominis in pregnancy'. D M Jones . *J. Clin. Pathol* 1967a. 20 p. .
- [Marmion and Harris ()] 'Mycoplasma pneumoniae and other medically important members of the family Mycoplasmataceae'. B Marmion , R Harris . *Mackie & Macrtiney Practical medical microbiology* Collee, J., Marimion, B., Fraser, A. and Simomons, A., (ed.) 1996. 30 p. . Churchill Livingstone (th ed)
- [Mendel et al. ()] 'Mycoplasma species in the vagina and their relation to vaginitis'. E B Mendel , D F Rowan , J H M Graham , D Dellinger . *Obstet. Gynecol* 1970. 35 p. .
- [Kundsins and Driscoll ()] 'Mycoplasmas and human reproductive failure'. R B Kundsins , S G Driscoll . *Surg. Gynecol. Obstet* 1970. 131 p. .
- [Gnarpe and Friberg ()] 'Mycoplasmas and Human reproductive failure'. H Gnarpe , J Friberg . *Am. J. Obstet. Gynecol* 1972. 114 p. .
- [Cassell and Cole ()] 'Mycoplasmas as agents of human disease'. G H Cassell , B C Cole . *N. Engl. J. Med* 1981. 2 p. .
- [Taylor-Robinson and McCormack ()] 'Mycoplasmas in human genitourinary infections'. D Taylor-Robinson , W M McCormack . *The Mycoplasmas* 1979. Academic Press Inc. II p. 307.
- [Krause et al. ()] *Mycoplasmas which infect humans "Molecular biology and pathogenesis*, D Krause , D Taylor-Robinson , R Elhaney , Mc , L Finch , J Baseman , J Maniloff . 1992. Washington, D. C: American Society for Microbiology. p. .
- [Shepard ()] 'Nongonococcal urethritis associated with human strains of T-mycoplasma'. M C Shepard . *JAMA* 1970. 211 p. .
- [Moller et al. ()] 'Pelvic inflammatory disease after hysterosalpingography associated with Chlamydia trachomatis and Mycoplasma hominis'. B R Moller , J Allen , B Toft , K B Hansen , D Taylor-Robinson . *Br. J. Obstet. Gynecol* 1984. 91 p. .
- [Embree et al. ()] 'Placental infection with Mycoplasma hominis and Ureaplasma urealyticum : Clinical correlation'. J E Embree , V W Krause , J A Embil , S Macdonald . *Obstet. Gynecol* 1980. 56 p. .
- [Al-Bahli ()] *Prevalence of genital Mycoplasma in women with selected obstetric and gynecological conditions*, S A H Al-Bahli . 1993. College of Medicine, Basrah University. (Arabic (M. Sc. thesis))
- [Al-Sulami et al. ()] 'Prevalence of genital mycoplasmas among women in the province of Basrah and an evaluation of their role in some cases'. A Al-Sulami , G J Al-Gizawi , R M Al-Mosawi . *J. Basrah Researches (Sciences)* 2007. 33 (2) p. .
- [Lind et al. ()] 'Serological crossreactions between Mycoplasma genitalium and Mycoplasma pneumoniae'. K Lind , B Q Lindhardt , H J Schutten , J Blom , C Christiansen . *J. Clin. Microbiol* 1984. 20 p. .
- [Al-Mosawi ()] 'Signs and symptoms of urethritis and cervicitis among women with or without genital mycoplasma infection in governorate of Basrah'. R M Al-Mosawi . *Journal of Basrah Researches* 2009. 35 (3) p. . (Sciences))
- [Toth et al. ()] 'Subsequent pregnancies among 161 couples treated for T-mycoplasmal genital tract infection'. A Toth , M L Lesser , G Brooks , D Labriola . *N. Engl. J. Med* 1983. 308 p. .
- [Fowlers et al. ()] 'T-mycoplasma and human infertility: Correlation of infection with alteration in seminal parameters'. D M Fowlers , J Macleod , W M O'Leary . *Fertil. Steril* 1975. 26 p. .
- [T-mycoplasma in the genitourinary tract of the female Microbiol. Scand. Sect. Br] 'T-mycoplasma in the genitourinary tract of the female'. *Microbiol. Scand. Sect. Br* 78 p. .
- [Taylor-Robinson and McCormack ()] 'The genital mycoplasma. Part II'. D Taylor-Robinson , W M McCormack . *N. Engl. J. Med* 1980. 320 p. .
- [Taylor-Robinson ()] 'The history and role of Mycoplasma genitalium in sexually transmitted diseases'. D Taylor-Robinson . *Genito-urin. Med* 1995. 71 p. .
- [Razin and Freundt (ed.) ()] *The mycoplasmas*, S Razin , E A Freundt . Krieg, R. N. & Holt, G. J. (ed.) 1984.
- [Holt et al. (ed.) ()] *The Mycoplasmas or (Mollicutes): cell wall-less bacteria In :Bergey's Manual of determinative bacteriology*, J G Holt , N R Krieg , P H Sneath , J T Staley , S T Williams . Williams & Wilkins, U.S.A. (ed.) 1994. 30 p. .
- [Upadhyaya et al. ()] 'The role of Mycoplasmas in reproduction'. M Upadhyaya , B M Hibbard , S M Walker . *Fertil. Steril* 1983. 39 p. .
- [Watts and Eschenbach ()] 'Treatment of chlamydia, mycoplasma and group B-Streptococcal infections'. D H Watts , D A Eschenbach . *Clin. Obstet. Gynecol* 1988. 2 p. .
- [Csonka and Williams ()] 'Tstrain mycoplasma in non-gonococcal urethritis'. G W Csonka , R E O Williams . *Lancet* 1966. 1 p. .

282 [Uno et al. ()] M Uno , T Deguchi , A Saito , M Yasuda , H Komeda , Y Kawada . *Prevalence of Mycoplasma*
283 *genitalium in asymptomatic men in*, 1997.

284 [Taylor-Robinson and Furr ()] 'Update on sexually transmitted mycoplasmas'. D Taylor-Robinson , P M Furr .
285 *Lancet* 1998. 351 (3) p. . (Suppl)

286 [Kundsin et al. ()] 'Ureaplasma urealyticum incriminated in perinatal morbidity and mortality'. R B Kundsin ,
287 S G Driscoll , P A Pelletier . *Science* 1981. 213 p. .

288 [Stray-Pedersen et al. ()] 'Uterine T-mycoplasma colonization in reproductive failure'. B Stray-Pedersen , J Erg
289 , T M Reikram . *Am. J. Obstet. Gynecol* 1978. 3 p. .

290 [Krause et al.] *Washington, D. C. (1992). Mycoplasmas which infect humans "Molecular biology and pathogenesis,*
291 D Krause , D Taylor-Robinson , R Elhaney , Mc , L Finch , J Baseman , J Maniloff . American Society for
292 Microbiology. p. .