

Reproductive Health Education Package to Reduce Risky Sexual Behaviors among Undergraduates in Selected State Universities in Sri Lanka: A Controlled Trial

Upuli Amaranganie Pushpakumari Perera¹ and Chrishantha Abeysena²

¹ University of Colombo

Received: 7 December 2018 Accepted: 3 January 2019 Published: 15 January 2019

Abstract

To assess the effectiveness of a reproductive health educational package to reduce RSB among undergraduates in selected state universities in Sri Lanka. Methods: A quasi-randomized controlled trial was conducted among undergraduates in 2014. Two universities were selected randomly either to the intervention or as the control group. From each of the selected universities, 10 clusters of students were selected randomly. A reproductive educational package was developed with reference to a curriculum by Centre for Disease Control, adjusted locally. The intervention group (n=309) received two health educational sessions with a gap of one week while the control group (n=431) did not receive any. The primary outcome was the reduction of RSBs while the secondary outcomes were the improvement of overall knowledge and overall attitudes on reproductive health at three months after the intervention. Intention to treat analysis was performed. Baseline imbalances were controlled by applying logistic regression. Results were expressed as adjusted odds ratios (AOR) and 95

Index terms— health education, randomization, reproduction, sexual behaviors, undergraduates.

Abstract-Objective: To assess the effectiveness of a reproductive health educational package to reduce RSB among undergraduates in selected state universities in Sri Lanka.

Methods: A quasi-randomized controlled trial was conducted among undergraduates in 2014. Two universities were selected randomly either to the intervention or as the control group. From each of the selected universities, 10 clusters of students were selected randomly. A reproductive educational package was developed with reference to a curriculum by Centre for Disease Control, adjusted locally. The intervention group (n=309) received two health educational sessions with a gap of one week while the control group (n=431) did not receive any. The primary outcome was the reduction of RSBs while the secondary outcomes were the improvement of overall knowledge and overall attitudes on reproductive health at three months after the intervention. Intention to treat analysis was performed. Baseline imbalances were controlled by applying logistic regression. Results were expressed as adjusted odds ratios (AOR) and 95% confidence interval (CI).

Results: The reproductive health package reduces the RSB at three months of follow up after the intervention although it was not statistically significant (AOR=0.80; 95% CI 0.28-2.31). Further, the package significantly improves the knowledge (AOR=11.75; 95% CI 7.04-19.63) and desirable attitudes towards reproductive health (AOR=6.13; 95% CI 3.64-10.32).

Conclusions: The educational intervention may reduce RSB after three months. It was effective for the improvement of good knowledge and desirable attitudes on reproductive health.

Keywords: health education, randomization, reproduction, sexual behaviors, undergraduates.

1 I.

Plain English Summary Risky sexual behaviors are behaviors leading to unwanted pregnancies and sexually transmitted infections. This research was done to assess the success of the developed package to lessen the risky sexual behaviors. A health education package was developed with the help of available packages from other countries to suit for young adults in Sri Lanka. Then the package was introduced to a group of selected undergraduates in two universities in the Western province of Sri Lanka. Before the introduction of the package, their level of risky sexual behavior has been assessed with a paper of questions. After three months of the introduction of the package, the risk level was assessed again. The results revealed that the educational intervention may reduce risky sexual behaviors. It also revealed the improvement of the knowledge and positive attitudes on risk behaviors and reproductive health.

2 II.

3 Introduction

Risky sexual behaviors (RSB) are behaviors leading to unintended pregnancies and sexually transmitted infections (STI) as defined as Centre for Disease Control (CDC) [1]. Three studies reported that the prevalence of risk behavior was higher than the level expected by parents and teachers [2,3,4]. Being in the age groups of adolescents and young adults, undergraduates may not be spared in practicing RSBs. Adolescents and young adults need different kinds of programs to prevent RSB. There is inconsistency among youth with respect to cognitive and social maturity and sexual experience. Therefore, interventions need to be tailor-made to meet the different requirements of different youth groups [5]. One review showed that determinants of safer sex behaviors among college students were alcohol usage, religiosity, barriers to condom use and perceived social norms. [6] Sexuality education has often concentrated on the risks associated with sexual behavior such as unintended pregnancies and STIs. Interventions for young people are usually implemented in schools, through peer educators, as community activities, and as mass media campaigns.

Several studies were performed to aim the reduction of RSB among adolescents. Two quasiexperimental studies have proven that educational interventions were effective in improving knowledge in R reproductive health (RH) in youth [7,8]. Another trial, which was conducted among adolescent girls between 15 to 19 years of age in India, revealed significant improvement of knowledge at one month after implementation of a RH package. [9]. Another quasiexperimental intervention targeted for adolescents of schools in China showed that an educational intervention is effective for increasing knowledge and improving attitude, not the behavior [10]. Similarly, a cluster randomized trial which was carried out to assess theoretically based sex education programme conducted by teachers revealed that the intervention did not reduce sexual risk taking in school going adolescents [11]. In contrast, another cluster randomized trial reported that in addition to sex education, skills building and motivational intervention was effective for reducing RSB among African-American male adolescents [12]. A review reported that sexual health education programs based on a comprehensive curriculum are able to improve knowledge and lessen the RSB [13]. It further reported that sex education programs do not increase sexual activity among adolescents and young people. Results from a Cochrane review showed that a combination of educational and contraceptive-promoting interventions showed the reduced the risk of unintended pregnancies among adolescents [14]. A study carried out to assess the relationship between condom availability and RSB among adolescents in high schools in Massachusetts revealed the success of programmes in human immunodeficiency virus (HIV) prevention [15].

Promoting Sexual Health is a comprehensive curriculum designed to provide information and build skills that reduce unintended pregnancies, reduce the transmission of STI including HIV and improve the quality of sexual experience [16]. The curriculum was designed for young adults in ages 18-24 years who are or to be sexually active to promote sexual health by addressing several aspects of sexuality such as the positive side of sexuality, pleasure, and satisfaction. Further, it is based on a logic model that specifies the important goals, particular behaviors that will lead to those health goals, and cognitive, sexual, psychosocial factors that affect those behaviors. Several authors have used similar contents within their interventions to reduce RSB [8,9,10,11,15,17].

There is a dearth of published literature on interventions for risky sexual behavior in Sri Lanka. Hettiarachchi has developed and successfully delivered a new educational intervention for teachers to increase the knowledge and attitude in sexual and RH among school children and teachers in Sri Lanka [18]. There are cultural taboos that have made it difficult for parents, teachers, and community leaders to openly discuss sexual issues among themselves or with adolescents [19]. The same cultural taboos that have hampered even marketing campaigns to promote greater use of contraceptives including condoms. This emphasizes the need for culturally sensitive reproductive education package for reducing RSB. Only those students who perform well in advance level examination are selected for state universities in Sri Lanka. Most of the university students stay away from their families. They are young adults and have more independent than school going adolescents. Therefore the university students are more vulnerable than the school children. Findings of this study are intended to inform the relevant authorities to design and deliver risk behavior reducing program for the youth in universities. Therefore, the objective was to assess the effectiveness of an educational intervention to reduce RSB among undergraduates in the selected state universities in Sri Lanka.

4 III.

5 Materials and Methods

A quasi-randomized controlled trial was conducted among the second and third year undergraduates in the University of Kelaniya and the University of Sri Jayewardenepura in 2014.

The exclusion criteria were undergraduates from foreign countries due to their different socio-cultural background and clergymen undergraduates due to the sensitive nature of the selected subject.

The intervention was a health education intervention conducted by four medical officers. It consisted of two-day interactive sessions of three and a half hours, conducted one week apart. No educational sessions were conducted to the control group. The intervention was designed using the curriculum on 'Promoting Sexual Health' [16]. Expert opinion was sought from the National STD/AIDS control programme and Health Education Bureau (HEB) for adapting the curriculum. The component which aim to improve the quality of sexual experience were removed. Case studies were changed according to local culture. The baseline information on knowledge, attitudes, and behaviors on reproductive health from previous national survey was utilized for the modification of the intervention. The available 'information, education and communication' (IEC) materials on family planning, STI and RSB were used with the expert opinion for the development of new materials. The developed intervention covered relevant topics in the prevention of STIs and unintended pregnancies. It included facts on contraceptive methods, i.e. effectiveness, side effects and ways of overcoming barriers. Furthermore, it described STIs with possible consequences and informs about places for testing. Dual protection given from condom was highlighted and condom demonstration was included emphasizing the problems in use. Multiple and overlapping partners' role in STIs was discussed. A mutual monogamous relationship was encouraged and the importance of the ability to insist on condom use was emphasized. In addition developed self-efficacy and skills to express or refuse sex. Almost all aspects in the prevention of STI and unintended pregnancies were covered by the developed intervention. The developed intervention was pretested among a group of 20 undergraduates outside the study population to test the feasibility, accessibility, co-operation and other logistical aspects of the designed intervention. Necessary changes were made to the intervention accordingly.

The primary outcome was the reduction of RSB among undergraduates during the three months of postintervention period. The secondary outcomes were overall knowledge and attitudes on RH among undergraduates at three months after the intervention. RSB is defined as a practice of one or more following behavior/s, having more than one sexual partner, alcohol use with sexual activities, inability to use condoms to prevent STI with commercial sex workers or non-commercial partners and inability to use in sexual activities with commercial sex workers or non-commercial partners. The overall knowledge on reproductive health was based on the knowledge on unsafe abortions (7 statements), contraceptives (10 statements), condoms (7 statements), STIs (28 statements) and HIV/AIDS (13 statements). One mark was given for each correct answer and zero marks was given for incorrect and do not know answers. A composite score on knowledge ranged from 0 to 65 and expressed as percentages. Equal or more than 75% was categorized as good knowledge and less than 75% as average knowledge for data analysis. The overall attitudes were based on attitude towards the use of contraceptives (5 statements), condoms (8 statements) and HIV/AIDS (9 statements). Each statement comprised of five responses as strongly agree, agree, neutral, disagree, strongly disagree. One mark was allocated for desirable attitudes while minus one mark for undesirable attitudes and zero marks for neutral responses. The range was from (-22) to +22. To make the values positive, +22 was added. The composite score ranged from 0 to 44. Cutoff marks were given for desirable and undesirable attitudes according to the inter quartile range. Those who scored 75th centile or above were taken as desirable attitudes and those who scored below 75% were taken as undesirable attitudes.

The University of Kelaniya and the University of Sri Jayewardenepura were selected as the intervention group and the control group respectively by tossing a coin. Ten clusters were selected randomly from all the clusters in each university. A cluster was defined as a tutorial group or a whole batch according to the structure of the selected undergraduate's group. We expected that the intervention will reduce the RSB from 13% to 6% among undergraduates [4]. Therefore, the calculated sample size was 301 for each group considering power as 80% and a false positive rate of 5%.

The developed intervention was introduced with IEC materials and all sections were discussed in detail. The opportunities were given to clarify any doubts. The intervention was conducted in the lecture halls for ten groups of undergraduates. Informed written consent was taken from the participants. A pre-tested self-administered questionnaire was used to assess pre and post-intervention data. The questionnaire comprised of five sections, the section I on socio-demographic and economic characteristics of the undergraduates, section II on other risk behaviors including alcohol, smoking and illicit narcotic drugs, section III on knowledge in selected aspects of RH, section IV on attitudes in selected aspects of RH, and section V on sexual behavior. The validity of the questionnaire was ensured by assessing the face, content and consensual validity. Content validity was assessed by checking whether or not all aspects of measures were covered using literature review and expert opinion. Consensual validity was determined by assessing the agreement of the experts on whether or not the conceptual definition has been used appropriately in the tool. A multi-disciplinary panel of experts in the fields of public health and reproductive health was used for assessment of validity. On the data collection day, the selected undergraduates (the tutorial group or batch which was identified) were taken to a separate lecture hall. The

participants were seated with appropriate gaps from each other to maintain confidentiality. An envelope was given to each participant to hand over the filled questionnaire.

Baseline data between the intervention and the control groups were analyzed using the chi-square test. The effectiveness of the intervention was assessed by comparing post-interventional data between intervention and control groups, the results were expressed as odds ratios (OR) and its 95% confidence intervals (CI). Baseline imbalances were controlled by applying multiple logistic regression. Sensitivity analysis was performed assuming that all participants who did not complete the follow-up data in the intervention group had high risk of RSB, average knowledge and undesirable attitudes, and those who were in the control group had a low risk of RSB, good knowledge and desirable attitudes.

6 IV.

7 Results

Of 309 undergraduates who were eligible and invited, only 300 post-interventional data could be collected from the intervention group. Of 427 undergraduates who were eligible and invited in the control group, 297 participated in the post-interventional assessment.

There were no statistically significant differences in the intervention and the control groups with respect to sex, ethnicity, religion, age and marital status (Table 1).

The proportion of undergraduates staying at boarding places were higher among the control group than the intervention group ($p < 0.001$). The control group had more undergraduates with family income $\geq 50,000$ rupees per month than the intervention group ($p = 0.014$). There were more undergraduates in Bio-Science stream in the intervention group (35.9%) than the control group (7.0%) ($p < 0.001$). With regard to school type, the control group consisted of 35.9% from mixed schools (both girls and boys) compared to 44.6% in the intervention group ($p = 0.04$). There were no differences with regard to the importance of religion and participating in religious activities between intervention and control groups. With regard to the availability of relatives and friends to discuss sexually related issues, there was no difference between the groups.

Significant differences were observed between the intervention and the control groups with regard to some risk behaviors before the intervention. Attended nightclub ever in life, usually going to cinema halls, taken alcohol ever and using internet > 2 hours per day showed significant differences between the two groups. (Table 2)

The undergraduates in the intervention group reported a higher percentage of good knowledge on contraceptives ($p < 0.001$) than the control group (Table 3).

Knowledge on condoms, knowledge on HIV/AIDS, knowledge on STIs and overall knowledge on RH did not show significant differences between the two groups. There were no differences with regard to attitudes in contraceptives, condoms, HIV/AIDS or overall attitudes in RH among undergraduates between the intervention and the control groups (Table 3 and 4). There was no significant difference of RSB within last three months between the intervention and the control group before the intervention (Table 4).

There was no significant difference observed between the intervention and the control group after three months of intervention with related to RSB (OR=1.39, 95% CI 0.81-2.39) (Table 4). However the adjusted OR was 0.8 (95% CI 0.28-2.31) which indicate a lower risk of RSB of the intervention group even though it was not statistically significant.

Overall good knowledge on reproductive health was higher (52.2%) among the intervention group compared to the control group (12.8%) at three months after the intervention (OR=7.48, 95% CI 4.97 -11.26) (Table 4). Even after adjustment for baseline imbalances, it remained higher (adjusted OR=11.75; 95% CI 7.04-19.63). The overall positive attitudes towards RH among undergraduates was higher in the intervention group (50.7%) compared to the control group (26.9%). The difference was significant statistically (OR=2.79, 95% CI 1.98 -3.92) (Table 4). Even after adjustment for baseline imbalances it was higher (adjusted OR = 6.13; 95% CI 3.64-10.32). The sensitivity analysis has not changed the direction of the association with any outcome (Supplementary Table 1). V.

8 Discussion

The intervention was not effective for reducing RSB at three months after the intervention. The overall knowledge and desirable attitudes on RH had been improved significantly in the intervention group when compared to the control group after three months of the intervention.

The two groups were allocated to the intervention and control group randomly to minimize selection bias. Contamination effect was minimized by selecting two separate universities for the intervention and control groups. There are only four universities in the western province, therefore considering a university as a cluster is not adequate for a cluster randomized trial. We randomly selected 10 clusters from each university. Even though we did not consider cluster effect for the calculation of sample size and statistical analysis. It was not feasible to conduct a cluster randomized trial within a university due to a risk of contamination. Some of the published results from RH interventions based on quasi-experimental studies [7,8,9,10]. It is natural to have a higher rate of loss to follow up among the control group as they were not intended for any benefits. The selection of three months of follow up period was done according to available literature and considering the feasibility of the collection of post-interventional data [12]. However, one study reported that the time period of three months may not be

adequate to detect a significant reduction of RSB [9]. Further, the undergraduates in our study were a low-risk cohort and therefore the usefulness of the intervention is limited. However, the intervention maybe effective for more vulnerable adolescents who have fallen out of the education system. The impact of the un-blindness of the investigators did not affect the outcome assessments as the use of a self-administered questionnaire. Further, the blindness of the participants was not feasible for health educational intervention. After controlling the baseline imbalance between the intervention and the control group, changed the effect of the intervention from harmful effect to a protective effect for RSB. However, it was not statistically or practically significant.

In contrast to our findings, a similar type of health education intervention carried out among male adolescents in America has improved RSB among them [12]. Even though the follow-up period was similar they have used different IEC materials such as videos, games, exercises and role-playing. The observed difference of results in the present study could be due to methodological variations, the difference in intervention curriculum and the baseline characteristics of the study population. A review of sexual education interventions to assess the impact on sexual behaviors revealed that sex education programmes were effective at reducing RSB among adolescents and young people [13]. The review revealed the effectiveness of reducing the number of sexual partners, increasing condom usage and increasing contraceptive usage.

Reproductive health-related information can be delivered successfully by health education interventions in various academic institutions through various channels. Knowledge and attitudes were improved among school-going adolescents by delivering a health educational intervention directed at school teachers in a district of Sri Lanka [18]. It represented a different study population; school children in Grade nine were educated through their teachers.

Knowledge and attitude improvement were observed among them and their evidence supports the findings of the present study. Another health educational intervention among female students between 15 to 19 years of age carried out in India, reported improvement of knowledge in reproductive health after one month of postintervention [9].

VI.

9 Conclusions

The educational intervention may reduce RSB after three months. It was effective for the improvement of good knowledge and desirable attitudes on reproductive health. Our study shows that educated youth improved their knowledge and attitudes about reproductive health, which is not surprising with their educational backgrounds. The intervention can be applied even for youths in working places without many alterations. However further research is needed to assess whether RSB for this group is improved in the long term follow them up.

10 VII.

11 Limitations

Follow up period after the intervention was three months period which may be inadequate to see significant changes in behavior. Even though it was unfeasible, unblinding of the participants might underreport of their self-assessed RSB. Consent for publication: Not applicable.

12 Availability of data and material:

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Clinical Trial registry: SLCTR/2014/003, registered on 1 st of Feb 2014

Funding: The Medical Research Institute of the Ministry of Health, Sri Lanka funded for the data collection of the study.

Authors' contributions: Both authors have contributed equally to the design the study. UP and CA analyzed and interpreted the data. UP was responsible for the conduct of the literature review and implementation of study and a major contributor in writing the manuscript. All authors read and approved the final manuscript.

Figure 1:

1

Variable		Intervention	N=309	Control	N=427	n	?2 p value
		n (%)		(%)			
Sex	Male Female	87 (28.2)	222 (71.8)	127 (29.7)	300 (70.3)		0.22 0.64
Ethnicity	Sinhalese Mus- lim	306 (99.0)	3 (1.0)	399 (93.4)	28 (6.6)		13.9 <0.001
Religion	Buddhist Non Buddhist	290 (93.9)	19 (6.1)	388 (90.9)	39 (9.1)		2.2 0.14
Resident 1	Boarding place Home/Relative	197 (63.8)	112 (36.2)	332 (77.9)	94 (22.1)		17.8 <0.001
Age 2	? 22 years ? 23 years	99 (32.1)	209 (67.9)	125 (29.5)	299 (70.5)		0.59 0.44
Marital sta- tus 1	Unmarried Ever married	305 (98.7)	4 (1.3)	418 (97.9)	9 (2.1)		0.68 0.41
Family income (Rs.) 3	<50000 >50000	242 (84.3)	45 (15.7)	363 (90.5)	38 (9.5)		6.1 0.014
Financial assistance (Rs.) 4	<3000 >3000	200 (92.2)	17 (7.8)	343 (93.0)	26 (7.0)		0.12 0.72
Academic year	Second Third	192 (62.1)	117 (37.9)	241 (56.4)	186 (43.6)		2.4 0.12
Study stream	Non-Biology Bi- ology	199 (64.4)	110 (35.6)	397 (93.0)	30 (7.0)		95.0 <0.001

Figure 2: Table 1 :

School type 5	Non-mixed	104 (44.6)	129	112 (35.9)	200	4.26
	Mixed	(55.4)		(64.1)		0.04
Importance of religion	Very im- por- tant	257 (83.2)	52	350 (82.0)	77	0.18
	Not very im- por- tant	(16.8)		(18.0)		0.67
Participating religious activities 6	Weekly or more	141 (45.9)		205 (48.6)		0.50
	Less than weekly	166 (54.1)		217 (51.4)		0.48
Access to a relative to talk in sexual issues	Yes	238 (77.0)		315 (73.8)		1.02
	No	71 (23.0)		112 (26.2)		0.31
Access to a friend to talk in sexual issues	Yes	266 (86.1)		384 (89.9)		2.57
	No	43 (13.9)		43 (10.1)		0.11
Variable	Intervention N=309 n (%)			Control N=427 n (%)		?2 p value
Attend night clubs*						
Yes		12 (3.9)		4 (0.9)		7.25
No		297 (96.1)		420 (99.1)		0.007
Attend nightclubs in previous one month						
Yes		6 (1.9)		3 (0.7)		2.28
No		303 (98.1)		424 (99.3)		0.13
Using Internet facilities						
Yes		290 (93.9)		378 (88.5)		6.07
No		19 (6.1)		49 (11.5)		0.014
Using Internet facilities >2 hours per day						
Yes		105 (34.0)		95 (22.2)		12.46
No		204 (66.0)		332 (77.8)		<0.001
Going to cinema halls*						
Yes		187 (60.5)		219 (51.5)		5.85
No		122 (39.5)		206 (48.5)		0.02
Going to cinema halls in last month						
Yes		30 (9.7)		27 (6.3)		2.88
No		279 (90.3)		400 (93.7)		0.09
Had taken alcohol ever						
Yes		85 (27.5)		76 (17.8)		9.89
No		224 (72.5)		351 (82.2)		0.002
Had taken alcohol in last three months						
Yes		48 (15.5)		42 (9.8)		5.42
No		261 (84.5)		385 (90.2)		0.02
Had smoked ever						
Yes		33 (10.7)		30 (7.0)		3.06
No		276 (89.3)		397 (93.0)		0.08
Had smoked in last three months						
Yes		18 (5.8)		19 (4.4)		0.71
No		291 (94.2)		408 (95.6)		0.39
Had taken Cannabis ever						
Yes		15 (4.8)		18 (4.2)		1.61
No		294 (95.2)		409 (95.8)		0.89

3

Knowledge/Attitude	Intervention N=309 n (%)	Control N=427 n (%)	?2 value	p
Contraceptives				
Good knowledge	112 (36.2)	88 (20.6)	22.15	
Average Knowledge	197 (63.8)	339 (79.4)	<0.001	
Condoms				
Good knowledge	126 (40.8)	156 (37.2)	0.95	
Average Knowledge	183 (59.2)	268 (62.8)	0.33	
STI				
Good knowledge	42 (13.6)	62 (14.5)	0.13	
Average Knowledge	267 (86.4)	365 (85.5)	0.72	
HIV/AIDS				
Good knowledge	172 (56.2)	244 (57.4)	0.10	
Average Knowledge	134 (43.8)	181 (42.6)	0.75	
Contraceptives				
Desirable attitude	85 (27.5)	113 (26.5)	0.10	
Undesirable attitude	224 (72.5)	314 (73.5)	0.75	
Condoms				
Desirable attitude	92 (29.9)	140 (32.8)	0.71	
Undesirable attitude	216 (70.1)	287 (67.2)	0.40	
HIV/AIDS				
Desirable attitude	127 (41.1)	174 (40.9)	0.002	
Undesirable attitude	182 (58.9)	251 (59.1)	0.96	
1 missing data=5, 2 missing data=1, 3 missing data=2				

Figure 4: Table 3 :

Outcomes	Pre-intervention			Post-intervention			
	Intervention	Control	OR	Intervention	Control	OR	AOR
	n	group	95 %	n	group	95 %	95 %
	group		CI	group		CI	CI
	N=309	N=427	p value	N=300	N=297	p value	p value
Risky sexual behavior			1.36			1.39	0.8 *
Yes	48	51	(0.89-2.1)	34	25	0.81-2.39	0.28-2.31
No	261	376	0.16	266 (88.7)	272 (91.6)	0.23	0.68
	(84.5)	(88.1)					
Overall knowledge on reproductive health			2.02			7.48	11.75 **
Good knowledge	63	48	1.34-3.0	157	38	4.97-11.26	7.04-19.63
	(20.4)	(11.2)		(52.3)	(12.8)		
Average knowledge	246	379	<0.001	143	259	<0.001	<0.001
	(79.6)	(88.8)		(47.7)	(87.2)		
Overall attitudes in reproductive health			1.04			2.79	6.13 ***
Desirable attitudes	79	106	0.74-1.46	152	80	1.98-3.92	3.64-10.32
	(25.6)	(24.8)		(50.7)	(26.9)		
Undesirable attitudes	230	321	0.82	148	217(73.1)	<0.001	<0.001
	(74.4)	(75.2)		(49.3)			

OR odds ratio, AOR adjusted odds ratio

* Adjusted for pre-intervention RSB, attended night club in last one month, reside outside the home and bio

** Adjusted for pre-intervention knowledge on reproductive health, reside outside the home and bio-science

*** Adjusted for pre-intervention attitudes on reproductive health, pre-intervention knowledge on reproductive health, bio-science stream.

Figure 5: Table 4 :

.1 Acknowledgements

All the undergraduates who participated for the study and the staff of the Universities, data collectors and the members of the Board of Study in Community Medicine, Postgraduate Institution of Medicine.

.2 List of Abbreviations

- [Oringanje et al.] , C Oringanje , M M Meremikwu , H Eko , E Esu , A Meremikwu , J Ehiri . (Interventions for preventing unintended pregnancies among adolescents)
- [Cochrane Database of Systematic Reviews ()] , 10.1002/14651858.CD005215.pub3. *Cochrane Database of Systematic Reviews* 2016. 2. (Art. No.: CD005215)
- [Silva et al. ()] 'Adolescent and youth reproductive health in Sri Lanka. Institute of policy studies in Sri Lanka'. De Silva , W I Somanathan , A Eriyagama , V . *Colombo. Dental Education* 2003. 75 (4) p. .
- [Manjula et al. ()] *An educational intervention study on adolescent reproductive health among preuniversity girls in Davangere district, South India. Annals of tropical medicine and public health*, R Manjula , S V Kashinakunti , R G Geethalakshmi , D Sangam . 2012. 5 p. .
- [Blake et al. ()] 'Condom availability programs in Massachusetts high schools: relationships with condom use and sexual behavior'. S M Blake , R Ledsky , C Goodenow , R Sawyer , D Lohrmann , R Windsor . *Am J Public Health* 2003. 93 (6) p. .
- [Kanekar and Sharma ()] 'Determinants of safer sex behaviors among college students'. A Kanekar , M Sharma . *Acta Didactica Napocensia* 2010. 3 (1) p. .
- [Pedlow and Carey ()] 'Developmentally-Appropriate Sexual Risk Reduction Interventions for Adolescents: Rationale, Review of Interventions, and Recommendations for Research and Practice'. C T Pedlow , M P Carey . *Annals of Behavioral Medicine* 2004. 27 (3) p. .
- [Coyle et al. ()] 'Draw the Line/Respect the Line: A Randomized Trial of a Middle School Intervention to Reduce Sexual Risk Behaviors'. K K Coyle , D B Kirby , B V Marín , C A Gómez , S E Gregorich . *American Journal of Public Health* 2004. 94 (5) .
- [Li et al. ()] *Effect of social cognitive theory-based HIV education prevention program among high school students in Nanjing, China, Health Education Research*, X Li , L Zhang , R Mao , Q Zhao , B Stanton . 2011. 26 p. .
- [Hettiarachchi et al. ()] 'Effectiveness of an educational intervention on sexual and reproductive health education directed at school teachers in the Kalutara district'. R Hettiarachchi , S Sivayogan , Sap Gnanissara . *Sri Lanka Journal of Social Sciences* 2008. 31 (32) p. 17.
- [Jemmott et al. ()] 'Effectiveness of an HIV/STD Risk-Reduction Intervention for Adolescents When Implemented by Community-Based Organizations: A Cluster-Randomized Controlled Trial'. J B Jemmott , L S Jemmott , G T Fong , K H Morales . *American Journal of Public Health* 2010. 100 (4) p. .
- [Ajuwon and Brieger ()] 'Evaluation of a schoolbased Reproductive Health Education Program in rural South Western'. A J Ajuwon , W Brieger . *Nigeria African Journal of Reproductive Health* 2007. 11 (2) p. .
- [Wight et al. ()] 'Limits of teacher delivered sex education: interim behavioral outcomes from randomized trial'. D Wight , G M Raab , M Henderson , C Abraham , K Buston , G Har , S Scott . *British Medical Journal* 2002. 324 (15) .
- [Brener et al. ()] *Methodology of the Youth Risk Behavior Surveillance System Morbidity and Mortality Weekly Report*, N D Brener , L Kann , S A Kinchen , J A Grunbaum , L Whalen , D Eaton , J Hawkins , Ross J . 2004. 53. (RR-12)
- [Charles et al. ()] *Promoting Sexual Health: A Curriculum to Reduce Unintended Pregnancy, Prevent Sexually Transmitted Infection and Improve Sexual Experience*, V Charles , D Kirby , G Lepore , J Walker , K Coyle . 2011. ETR Associates, California.
- [Parwej et al. ()] 'Reproductive health education intervention trial'. S Parwej , R Kumar , I Waleela , A Aggarwal . *Indian journal of Paediatrics* 2005. 72 (4) p. .
- [Fernando ()] *Sexual behavior and substance abuse among youth in the coastal region in Galle district. Thesis (MD in Community medicine)*, N Fernando . 2009. Postgraduate Institute of Medicine, University of Colombo
- [Perera and Reece ()] 'Sexual behavior of young adults in Sri Lanka: Implications for HIV prevention'. B Perera , M Reece . *AIDS Care* 2006. 18 (5) p. .
- [Kirby ()] *The impact of sex education on the sexual behavior of young people, Population Division Expert Paper No.*, D Kirby . 2011. 2011. United Nations New York. 12.
- [Thalagala and Rajapaksha ()] *UNICEF National survey on emerging issues among adolescents in Sri Lanka*, N Thalagala , L Rajapaksha . 2004.