



GLOBAL JOURNAL OF MEDICAL RESEARCH: H
ORTHOPEDIC AND MUSCULOSKELETAL SYSTEM
Volume 19 Issue 3 Version 1.0 Year 2019
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

A Comprehensive Review of Surgical Supplies

By A. K. Mohiuddin

World University of Bangladesh

Abstract- Injury to the skin provides a unique challenge, as wound healing is a perplexing and multifaceted procedure. Intense wounds can possibly move from the intense wound to endless wounds, requiring the doctor to have a careful comprehension of outside intercessions to bring these wounds again into the healing course. Careful improvement/ dressings are applications for wounds, consumes, and ulcers. They ought to be viewed as steady of healing; are attractive however not fundamental in a crisis. There are right now a lot of dressings accessible in the market to help in wound healing. Before picking a dressing for particular damage, a doctor must evaluate cautiously the necessities of the wound to comprehend which dressing would guarantee most extreme intrigue. Fundamentally, there is nothing called best decision, and it is vital that the benefits/negative marks of each dressing framework be comprehended. This article has given a structure to help with dressing evaluation. This article uncovers estimation of wound healing and the elements of wound dressings. An assortment of dressings and their individual subtleties are point by point.

Keywords: *wound dressing; gauzes; absorbent; cotton fiber; mesh; sponges; napkins.*

GJMR-H Classification: *NLMC Code: WE 168*



Strictly as per the compliance and regulations of:



© 2019. A. K. Mohiuddin. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

A Comprehensive Review of Surgical Supplies

A. K. Mohiuddin

Abstract- Injury to the skin provides a unique challenge, as wound healing is a perplexing and multifaceted procedure. Intense wounds can possibly move from the intense wound to endless wounds, requiring the doctor to have a careful comprehension of outside intercessions to bring these wounds again into the healing course. Careful improvement/dressings are applications for wounds, consumes, and ulcers. They ought to be viewed as steady of healing; are attractive however not fundamental in a crisis. There are right now a lot of dressings accessible in the market to help in wound healing. Before picking a dressing for particular damage, a doctor must evaluate cautiously the necessities of the wound to comprehend which dressing would guarantee most extreme intrigue. Fundamentally, there is nothing called best decision, and it is vital that the benefits/negative marks of each dressing framework be comprehended. This article has given a structure to help with dressing evaluation. This article uncovers estimation of wound healing and the elements of wound dressings. An assortment of dressings and their individual subtleties are point by point.

Keywords: wound dressing; gauzes; absorbent; cotton fiber; mesh; sponges; napkins.

I. BACKGROUND

Wound healing is a dynamic and complex procedure which requires appropriate condition to advance healing procedure. Generally, wet-to-dry dressings have been utilized broadly for wounds requiring debridement. In 1600 BC, Linen strips absorbed oil or oil secured with mortars was utilized to block wounds. Dirt tablets were utilized for the treatment of wounds by Mesopotamian starting point from around 2500 BCE. They cleaned wounds with water or milk before dressing with nectar or tar. Wine or vinegar use for cleaning the wounds with nectar, oil and wine as further treatment was trailed by Hippocrates of old Greece in 460-370 BCE. They utilized fleece bubbled in water or wine as a gauze. There was a noteworthy leap forward in the clean procedure during the nineteenth century, anti-infection agents were acquainted with control contaminations and reduction mortality. Present day wound dressing landing was in 21th century. Woven spongy cotton bandage was utilized in 1891. Until the mid-1900's, it was solidly accepted that wounds mended all the more rapidly whenever kept dry and revealed while 'shut wounds recuperate more rapidly than open wound' written in an Egyptian restorative content - Edwin smith careful papyrus in 1615 BC. Oscar Gilje in 1948 portrays clammy chamber impact for

healing ulcers. In the mid 1980's, the main present day wound dressing was presented which conveyed significant attributes giving dampness and retaining liquids (for example polyurethane froths, hydrocolloids, iodine-containing gels). During the mid-1990's, manufactured wound dressings ventured into different gathering of items which incorporates hydrogels, hydrocolloids, alginates, engineered froth dressing, silicone networks, tissue cements, vapor-porous cement movies and silver/collagen containing dressing. At the point when the wound is shut with dressing they are ceaselessly presented to proteinases, chemotactic, supplement and development factors, which is lost in the wound uncovered. In this way, during late twentieth century, generation of occlusive dressing started to secure and give clammy condition to wound. These dressings help in quicker re-epithelialization, collagen combination, advances angiogenesis by making hypoxia to the wound bed and diminishes wound bed pH which prompts decline in the wound disease. With the progression in innovation, in excess of 3000 items have been created to treat various kinds of wounds by focusing on different parts of healing procedure.

Author: Assistant Professor, Department of Pharmacy, World University of Bangladesh. e-mail: trymohi@gmail.com



Figure 1: Surgical dressing for wound healing (Source: 4g Inwi Configuration)

a) *Purpose of the study*

Discussion and projection of wound healing by market accessible careful supplies. The present audit follows the historical backdrop of dressings from its soonest origin to the present status and furthermore talks about the preferred position and restrictions of the dressing materials.

b) *Findings*

There is a mind-boggling measure of wound dressings accessible in the market. Present day world and innovation offered ascend to different method for wound healing with advancements. Practically a wide range of advancements are accessible in careful outlets, a couple of them are restricted to medical clinic settings. This suggests the absence of full comprehension of wound consideration and the board. The purpose of utilizing propelled dressings is to enhance explicit wound attributes to carry it as near "perfect" as could be allowed. It is simply after appropriately evaluating the wound qualities and getting learning about accessible items that the "perfect" dressing might be picked.

c) *Materials and Methods*

Research led an all year extensive writing search, which included specialized pamphlets, papers diaries, and numerous different sources. The present examination was begun from the earliest starting point of 2018. PubMed, ALTAVISTA, Embase, Scopus, Web of Science, and the Cochrane Central Register of was altogether looked. The catchphrases were utilized to

look for changed distributors' diaries like Elsevier, Springer, Willey Online Library, Wolters Kluwer were broadly pursued. Drug and specialized specialists, pharma organization delegates, medical clinic attendants and scientific experts were given their profitable recommendations. Projections depended on various sorts of careful supplies accessible in home and abroad.

d) *Research limitations*

Pictorial introduction of such a significant number of sorts of dressings are unrealistic to imitate in an article yet for a speedy survey, the article involves the majority of them. Likewise, sutures are not point by point which will be incorporated by the following article.

e) *Practical Implication*

The sole of this article was to detail a few kinds of careful supplies. Alongside understudies, specialists and experts of various foundation and controls, for example Advisors, medical procedure partners, specialists, attendants, emergency clinic experts and pharmacists need to procure much from this article.

II. INTRODUCTION


A professional service rendered by many pharmacists consists of supplying surgical instruments, sutures, surgical dressings, and other equipment employed by the surgical personnel during and after a surgical operation. Some pharmacists who have obtained the necessary background of information

carries a complete line of such supplies and even are able to provide operating tables and other heavy equipment. There are comparatively few such completely equipped pharmacies; the major outlet is through surgical supply houses. Every pharmacist, however, should be familiar with two of the products mentioned above, namely, Surgical Dressings and Sutures, which are discussed in detail below. The selection of the correct type of surgical dressing or suture is a crucial factor in protecting the welfare of the patient undergoing surgery. Many items in these categories are handled routinely by pharmacists, and all of these items come within the purview of their professional responsibility.

a) *Types of Wounds*

An intense wound is damage to the skin that happens abruptly because of mishap or careful damage. It recuperates at an anticipated and expected time span more often than not inside 8-12 weeks relying upon the size, profundity and the degree of harm in the epidermis and dermis layer of the skin. Ceaseless wounds for the most part result from decubitus ulcer, leg ulcer and consumes. Wound healing is a dynamic and complex procedure of tissue recovery and development advance through four unique stages (I) the coagulation and hemostasis stage (following damage); (ii) the fiery

stage, (soon after damage to tissue) during which swelling happens; (iii) the expansion time frame, where new tissues and veins are shaped and (iv) the development stage, in which renovating of new tissues happens.



(Source: Blog wound infection management tips Saturday, May 22, 2010)

Figure 2: Types of wounds




Figure 3: Distinct and overlapping phases of wound healing [4]

There are a limited number of reasons a wound becomes chronic; however, once these reasons are rectified, the wound resumes its natural course of healing.

- **Arterial:** Generally, an ABI of under 50 mm Hg, or a flat-out toe pressure under 30 mm Hg (or under 50 mm Hg for people with diabetes) demonstrates basic appendage ischemia and predicts disappointment of wounds to mend.
- **Venous:** Pressure-incited changes in vein divider penetrability at that point lead to spillage of fibrin and other plasma parts into the perivascular space. Collection of fibrin has immediate and negative impacts on wound healing as it down-directs collagen combination.
- **Infection:** Underlying irresistible procedures including cellulitic and osteomyelitis procedures will repress wound healing. Refined for vigorous, anaerobic, and contagious pathogens is suggested.
- **Pressure:** Increased pressure to the territory of concern will obliterate new tissue development and counteract appropriate perfusion of blood to the wound site. These regions should be offloaded to maintain a strategic distance from pressure in the territory.

- **Oncologic:** Always biopsy territories of worry in nonhealing wounds, as this can be an atypical introduction of certain kinds of malignancies.
- **Systemic:** There are different foundational ailments which hinder wound healing, with diabetes being the most well-known offender. It has been resolved that uncontrolled blood glucose levels smother the body's ordinary provocative reaction, just as causing microvascular malady which cutoff points healing.
- **Nutrition:** While serum egg whites has not been observed to be a decent indicator wound healing, there is some proof that protein hunger, just as inadequate degrees of specific nutrients and minerals, will confine the body's capacity to mend constant wounds.
- **Pharmacological:** Hydroxyurea has been accounted for in various cases to cause nonhealing ulcerations.
- **Self-incurred/psychosocial:** There are cases where a patient is causing the ulceration, either deliberately or because of resistance. This is frequently the hardest factor to spot and survive, however should dependably be a thought [1-7].




Figure 4: The cellular and molecular differences between acute healing wounds and chronic non-healing wounds [7]. The healing of intense wounds (left) starts with a transient provocative reaction as granulation tissue is framed, which gives a situation appropriate to the re-epithelialization required to finish fix. Constant non-healing wounds (right) are regularly tainted and display persevering aggravation. By definition, re-epithelialization has slowed down yet is hyper-proliferative. Granulation tissue is problematic with raised lattice metalloproteinases (MMPs) present together with poor fibroblast and vein invasion. Fibrin sleeves can likewise be available that counteract the dissemination of oxygen through the wound, rendering it hypoxic.

b) *Uses of Dressings*

Surgical dressing is a term applied to a wide array of products used for dressing physical injury or diseased tissues. Dressings may serve to:

- Provide a domain for soggy wound healing. Drying up of a wound is a main consideration in improving damage healing and expanding seaming. Dressings that avert drying up give an ideal situation to autolysis cell movement, granulation, and re-epithelialization.
- Prevent maceration by allowing dissipation or assimilation. In very secretory wounds, unnecessary dampness and autolytic chemicals will debilitate fixing tissue and will expand odds of microbial infection.
- Promote hemostasis.
- Protect the wound from further harm (mechanical harm, microbial intrusion, lack of hydration, maceration, synthetic harm, change in pH).
- Reduce heat misfortune.
- Control microbial development (by joining of antimicrobial medications).
- Promote autolysis.
- Promote healing.
- Provide pressure, advancing hemostasis, and decreasing edema.
- Provide support.
- Reduce torment, increment patient solace, and improve useful utilization of wound site.
- Reduce smell.
- Improve the presence of the wound site.
- Reduce in general expenses related with wound treatment [8-12], [25].

c) *Selection of Wound Dressing*

In view of the wound kind, appropriate dressing material must be utilized. Dressing choice ought to be founded on its capacity to a) give or keep up soggy condition b) upgrade epidermal movement c) advance angiogenesis and connective tissue amalgamation d) permit gas trade between wounded tissue and condition e) keep up fitting tissue temperature to improve the blood stream to the wound bed and improves epidermal relocation f) give insurance against bacterial infection and g) ought to be non-disciple to the wound and simple to evacuate in the wake of healing h) must give debridement activity to upgrade leucocytes movement and bolster the amassing of compound and l) must be sterile, non-dangerous and non-unfavorably susceptible. Likewise, determination ought to be made based on the level of exudation, nearness or probability of infection, nearness of necrotic tissue, and anatomical site. The right choice of a wound dressing depends on the sort of wound as well as on the phase of fix. The utilization of a wound dressing can't be considered in seclusion, but instead with regards to an incorporated wound-care program [13-18].

d) *Types of Wound Dressings*

Within this classification, dressings are considered on the basis of use.

- Primary/secondary wound dressings
- Secondary dressings
- Absorbents
- Bandages
- Adhesive tapes
- Protectives [19, 20]

e) *Specifications*

Surgical dressings and sutures are required to meet specific requirements of the USP for many characteristics. For these specific requirements and the performance of several of the official tests, eg, Absorbency test and Fiber length of cotton, Diameter of sutures, and Tensile strength of sutures, textile fabrics, and films refer to the detailed instructions provided in the USP [21-24].

III. CLASSIFICATION

For all intents and purposes, the natural technique for characterization utilizes the terms essential and auxiliary dressing. An essential dressing legitimately contacts the wound. It might give absorbability and may forestall drying up, infection, and attachment of the auxiliary dressing to the wound. An auxiliary dressing is put over an essential dressing, giving further protect, absorbability, compaction, or impediment [24-29]. Albeit a few dressings are exclusively essential or optional in nature, others have the qualities of both. The accompanying arrangement is utilized here:

a) *Primary Wound Dressings*

i. *Plain Gauze*

Plain Gauze has been used as a primary dressing but will stick to all but clean, engraved wounds. In spite of the fact that this property has been utilized to unbridle exudative, tainted, and necrotic wounds, this training might be excruciating and is frequently broken, causing the expulsion of injury tissue and new epithelium [15], [25], [30]. Bandage and non-woven wound dressings are dry woven or non-woven wipes and wraps with fluctuating degrees of sponginess, in view of plan. Texture arrangement may incorporate cotton, polyester or rayon. Accessible sterile or non-sterile in mass and with or without a glue fringe. They are utilized for purging, pressing and covering an assortment of wounds [31].



(Source: Web Indiamart)

Figure 5: White Plain Gauze Bandage

ii. *Impregnated Gauze*

Impregnated Gauze is used to reduce its adherence to wounds. Cotton, rayon, or cellulose acetic acid derivation cloth has been impregnated with an assortment of substances, for example, oil or paraffin (Aquaphor, Beiersdorf, Vaseline (Sherwood), KY jam (Johnson and Johnson), petrolatum emulsion (Adaptic,

Johnson and Johnson), zinc saline (NutraDress, Derma Sciences), or sodium chloride (mesalt, SCA Molnlycke). Coatings may wear off, permitting epithelial ingrowth and requiring a dressing change [32-35]. Operators most normally utilized incorporate saline, oil, zinc salts, petrolatum, xeroform and red. Signs change dependent on the compound. They are non-disciple and require an auxiliary dressing. An optional dressing ought to be utilized with these dressings to counteract drying up, give sponginess, and forestall the passageway of pathogens. At the point when utilized with a proper auxiliary dressing, these dressings might be utilized in vigorously oozing wounds [20], [31]. Silver, in ionic or nanocrystalline structure, has for a long time been utilized as an antimicrobial operator especially in the treatment of consumes (as silver sulfadiazine cream). Iodine likewise can bring down the microbiological load in incessant wounds. Alert is required in patients with a thyroid infection inferable from conceivable foundational take-up of iodine. Metronidazole gel is regularly utilized for the control of smell brought about by anaerobic microbes [15].



(Source: eBay)

Figure 6: Dermagran SPD-21 Hydrophilic Impregnated Gauze Wound Dressings 4" x 4" - Box/15

iii. *Film Dressings*

Film Dressings (transparent film, occlusive or semi-occlusive) are films of polyurethane with acrylic or polyether adhesives that provide a semipermeable membrane to water vapor and oxygen yet are waterproof. Reasonable for level, shallow wounds with

low to medium exudates. Advance soggy condition. Hold fast to sound skin yet not to wound. Permit visual checks. May be left set up a few days. Give no padding. Not for contaminated or vigorously oozing wounds [15]. In softly radiating wounds they license enough disintegration to advance soggy wound healing and

avert maceration. Film dressings dispose of microbes from wounds and license washing and consideration of the wound. Film dressings will cling great to flawless skin and have a low adherence for wound tissue. They ought not be utilized in contaminated or vigorously radiating wounds. Film dressings may wrinkle, framing channels for microbial passageway. Trouble in taking care of film dressings has been overwhelmed by extraordinary plan of different application frameworks [102]. Notwithstanding their utilization as wound dressings, glue movies have been utilized to ensure regions helpless against pressure, grating, or shear ulceration or for imbue ment or cannulation locales.

these dressings are profoundly versatile and adaptable, and can fit in with any shape and don't require extra taping. Examination of wound conclusion is additionally conceivable without expulsion of wound dressing due to straightforward movies. Subsequently these dressings are suggested for epithelializing wound, shallow wound and shallow wound with low exudates [3]. Instances of straightforward film dressings are Opsite™, Tegaderm™, Biooclusive™, Biooclusive (R) Transparent Dressing (Johnson and Johnson), Opsite (Smith and Nephew), Tegaderm (3M), and Dermasite (Derma Sciences) [36-40].



(Source: Wound Care Advisor)

Figure 7: Transparent Film Dressing

b) Primary/Secondary Wound Dressings

i. Composite Dressings

Composite Dressings have primary and secondary components that prevent adherence to the wound, with some degree of absorbency. The level of impediment given by these dressings shifts. Discharge (Johnson and Johnson), Telfa (Kendall), and Melolin (Smith and Nephew) comprise of gently permeable rayon or cotton cushions sandwiched between permeable polyethylene films. Nu-Derm (Johnson and Johnson) and Lyofoam A (Seton Healthcare Group) comprise of polyurethane froths with a film backing [32], [41-43]. A composite or mix dressings has different layers and each layer is physiologically particular. A large portion of the composite dressings have three layers. Composite dressings are wound covers that

consolidate physically unmistakable segments into a solitary item to give different capacities, for example, a bacterial obstruction, assimilation, and attachment. Ordinarily, they are made out of numerous layers and fuse a semi-or non-follower cushion that covers the wound. May likewise incorporate a cement outskirts of non-woven texture tape or straightforward film [26]. External most layer shield the wound from infection, center layer typically made out of absorptive material which keeps up dampness condition and help autolytic debridement, base layer made out of non-disciple material which keeps from adhering to youthful grinding tissues. Composite dressings have less adaptability and they are progressively costly [103].



(Source: Fashion Herald)

Figure 8: Composite Dressing

ii. *Hydrogels*

Hydrogels are insoluble hydrophilic materials made from synthetic polymers such as poly (methacrylates) and polyvinyl pyrrolidone. The high-water substance of hydrogels (70-90 %) helps granulation tissues and epithelium in a damp domain. They are mind boggling grids in which the scattering medium is caught rather like water in an atomic wipe. Hydrogels are nonadherent dressings that through semipermeable film permit a high rate of dissipation (and cooling) without trading off wound hydration. This causes them helpful in to consume treatment. Hydrogels are likewise helpful in shaggy regions where entanglement of hair into the dressing would not be awful [53], [110]. Delicate versatile property of hydrogels gives simple application and evacuation after wound is recuperated with no

harm. Temperature of cutaneous wounds is diminished by hydrogels giving mitigating and cooling impact. Undefined hydrogel dressings are definitions of water, polymers and different fixings with no shape, intended to give dampness to a dry wound and to keep up a soggy healing condition. The high dampness substance serves to rehydrate wound tissue. Demonstrated for fractional and full-thickness wounds, wounds with putrefaction, minor consumes, and radiation tissue harm. Impregnated hydrogel wound dressings are bandages and non-woven wipes, ropes and strips soaked with a shapeless hydrogel [31]. Instances of hydrogels are Geliperm (Geistlich), Vigilon (Bard), Flexderm (Dow Hickam), and Nu-Gel (Johnson and Johnson). The last is held together with a fusible fiber scrim [44-47].




(Source: Healthy Kin)

Figure 9: Hydrogel Sheet Dressing

iii. *Hydrocolloid Dressings*

Hydrocolloid Dressings combine the benefits of occlusion and absorbency. Hydrocolloids are scatterings of particles around which water atoms and solvated particles structure a shell-like structure. Liquid assimilation happens essentially by molecule swelling and growth of this structure. The hydrocolloid mass of these dressings comprises of gum-like materials, for example, guar or karaya, sodium carboxymethyl-cellulose, and gelatin, bound by a cement, for example, polyisobutylene. Hydrocolloid dressings show wet tack (attachment to a wet surface) in light of molecule swelling. This property encourages atraumatic evacuation. The dry tack of hydrocolloid dressings is

because of a glue, for example, polyisobutylene, which is inactivated by dampness [53]. The dry tack held by the dressing around the wound jam the edge seal. Exudate retention by most hydrocolloid dressings results in the arrangement of a yellow/darker coagulated mass that remaining parts on the wound in the wake of dressing evacuation. This might be wiped out from the wound and ought not be mistaken for discharge. Since hydrocolloids ingest water gradually, they are of little use on intensely radiating wounds. They are, be that as it may, helpful for respectfully to profoundly exudative incessant wounds. Instances of hydrocolloid dressings incorporate Duoderm (ConvaTec), Comfeel Plus (Coloplast), and RepliCare (Smith & Nephew) [48-51].




(Source: health.planetfem.com)

Figure 10: Hydrocolloid Dressings

iv. *Hydro-cellular foam dressing*

Hydro-cellular foam dressing--Present day wound dressing, initiates soggy wound condition and advances wound healing. Bandage empowers scab development, which weakens epithelialization and increments vascular endothelial development factor (VEGF) articulation. Interestingly, hydro-cell froth dressing quickens epithelialization and new vessel development in granulation tissue [104]. Hydrocellular Foam Dressings with Silicone Adhesive are demonstrated for use on a wide scope of intense and constant wounds like venous leg ulcers, pressure ulcers, diabetic ulcers and horrendous wounds [31]. A hydrocellular froth dressing has turned into a first decision for the treatment of tolerably to vigorously

radiating wounds which require exorbitant wound debridement for giving ideal conditions at the wound site while keeping up a wet wound condition [105,108]. may advance wound healing alongside reduction in aggravation by diminishing quality articulation levels of IL-1 β , IL-6, and IL-10 [106]. Hydrocellular froth dressing (HCF) assimilates over the top wound liquid, which contains different cytokines and development factors, and guarantees a clammy domain to advance wound healing. Nonetheless, the sub-atomic instruments hidden the wound liquid segment changes instigated by HCF are inadequately comprehended. dermal fibroblast expansion is upregulated by HCF because of expanded leptin level at the wound surface, and these impacts advance wound healing [107].



(Source: HighTideHealth.com)

Figure 11: Allevyn Adhesive Hydrocellular Foam Dressings

v. Calcium Alginate Dressings

Calcium Alginate Dressings are a solid, flexible, and common wound consideration dressing normally connected to diabetic wounds, venous wounds, full-thickness consumes, split-thickness unite giver destinations, pressure ulcers, pit wounds, and endless ulcers. Alginate dressings can likewise help wounds that are draining [110]. The calcium in these dressings settles blood stream, which moderates draining [25]. Alginic corrosive is a normally happening polysaccharide gotten from dark colored kelp. As the calcium

salt, these stringy nonwoven dressings are exceptionally retentive and are utilized on reasonably to exceedingly oozing wounds [13]. They might be held set up with cloth tape or a film dressing. They likewise might be utilized to pack wounds. Alginate dressings can ingest up to multiple times their weight in wound liquid [13, 25]. Instances of calcium alginate dressings are Sorbsan (Dow Hickam), Algosteril (Johnson and Johnson), and Kaltostat (Calgon Vestal) (Table 1) [52-56].

Table 1: Commercially available alginate-based dressings [109]

Commercially Names	Composition	Applications
Algicell™	Sodium alginate, 1.4% silver	Diabetic foot ulcer, leg ulcers, pressure ulcers, donor sites, and traumatic and surgical wounds.
AlgiSite M™	Calcium alginate	Leg ulcers, pressure ulcers, diabetic foot ulcers and surgical wounds.
Comfeel Plus™	Sodium carboxymethylcellulose and calcium alginate	Ulcers such as venous leg ulcers, pressure ulcers; burns, donor sites, postoperative wounds and necrotic wounds.
Kaltostat™	Sodium alginate	Pressure ulcers, venous ulcers, diabetic ulcers, donor sites, and traumatic wounds.
Sorbsan™	Calcium alginate	Arterial, venous, and diabetic leg ulcers Pressure ulcers, post-operative wounds, donor and graft sites and traumatic wounds.
Tegagen™	Sodium alginate	Diabetic and infected wounds.
Guardix-SG®	Sodium alginate and poloxamer	To avoid post-operative adhesions in thyroid and spine surgeries.
SeaSorb®	Calcium alginate	Good for high exuding wounds e.g., ulcers such as diabetic and leg pressure ulcers.
Algivon®	Calcium alginate and Manuka honey	It eliminates odour and ideal for necrotic wounds and wounds with odours.
Fibracol™Plus	Calcium alginate and collagen	Full and partial-thickness wounds, for ulcers such as pressure ulcers, venous ulcers, diabetic ulcers and second-degree burns.

Commercially Names	Composition	Applications
Hyalogran®	An ester of hyaluronic acid and sodium alginate	Used for ulcers, diabetic wounds, pressure sores, ischemic, necrotic wounds.
Tromboguard®	Sodium alginate, calcium alginate, chitosan, polyurethane and silver cations	Used to stop bleeding in postoperative wounds, traumatic wounds, gun shots, skin graft donor sites, bleeding from accidents.



(Source: Home Medical Supplies)

Figure 12: MAXORB AG+ CMC / Alginate Dressings

c) Secondary Wound Dressings

i. Absorbents

Surgical Cotton: Cotton is the basic surgical absorbent. It is official Purified Cotton USP.

Domestic cotton developed in the Southern US is appropriate for careful purposes. The tamed cotton plant achieves a tallness of 2 to 4 ft. Developing from the seeds is a case or boll that blasts open after maturing, uncovering a mass of white cotton strands. Every one of these filaments is a moment, hair-like cylinder, the external divider being unadulterated cellulose, the opening loaded up with plant liquids. At the point when the boll blasts open, the fiber falls into a level strip like structure, turned and multiplied upon itself in excess of multiple times from start to finish. Citrus extract can be utilized both for the expendable (non-sturdy) materials (outfits, veils, and sleeves for circulatory strain estimation) and the materials that expect solidness to washing [53,116].

The raw cotton fiber precisely cleaned of soil and checked into layers however not generally treated, has a constrained use for paddings and covers of solid surfaces. This structure is provided under the name nonabsorbent cotton. It additionally is utilized as often as possible as cotton connects the bacteriological research center on account of its non-retentiveness [117].

Absorbent Cottonis set up from the crude fiber by a progression of procedures that expel the regular waxes and all debasements and outside substances and render the filaments permeable. It is a for all intents

and purposes unadulterated, white cellulose fiber. Other than the recognizable move structure, Purified Cotton might be gotten in different arranged structures, for example, cotton balls or cotton-tipped tools [117]. Ongoing examinations have appeared perpetual wounds contain elevated amounts of tissue and cytokine devastating proteases including collagenase and neutrophil elastase. The decrease in catalyst action with dialdehyde cotton dressing was affirmed in arrangement by deciding elastase restraint with dialdehyde starch. The dialdehyde cotton dressing likewise diminished elastase action in human wound liquid in a portion reaction connection dependent on weight of bandage per volume of wound liquid [118]. Retentive cotton shows brilliant dampness engrossing and water-holding properties, and ingested particles can be effectively caught among AC strands without delivering gravity water. Notwithstanding, fine filaments are inclined to stay in the wounds even after evacuation [119].

Absorbent balls made of a uniform careful gooey rayon fiber likewise are accessible. These assimilate liquids quicker and hold their shape superior to anything cotton balls. Nonabsorbent Bleached Cotton, arranged by an altered fading process that holds the water-repellent regular oils and waxes, additionally is accessible. This cotton is distinguished effectively by its luxurious feel. Since it is repellent to water, it doesn't end up tangled or inelastic. Subsequently, it is well-adjusted to pressing, cushioning, and padding of dressings over damaged



zones and as nonabsorbent sponsorship on clean napkins, joins, and seepage dressings. All around dyed and checked cotton having receptiveness at the very least 12 hours is utilized in pharmaceutical, keeping up aseptic conditions for assembling process and furthermore for pressing purposes. Non-spongy blanched cotton has water repellent properties on the grounds that while purging off filaments the common wax is held [63], [120-121].

Rayon, or recovered cellulose, is produced using wood or cotton linters. Subsequent to dissolving it in a blend of salt and carbon disulfide, cellulose string is reprecipitated in a corrosive coagulating shower by section through fine openings in a metal plate. Since plant lignins have been expelled, just as the more round cross area, rayon strands are milder and more brilliant than cotton [57-62].

ii. *Surgical Gauzes*

The capacity of careful cloth is to give a permeable material of adequate elasticity for careful dressings. It is known as Absorbent Gauze USP. During the time spent making careful bandage, the crude cotton fiber is cleaned precisely and after that spun or wound into a string, and the string thus is woven into an open-work fabric that is dark and nonabsorbent. It is faded white and offered permeable by much indistinguishable procedures from those utilized in the availability of careful cotton [63]. The bandage in this way treated is dried by passing a constant length through a tentering machine. Tenterhooks rectify, stretch, and hold it tight as it is dried. When it leaves this hardware, the dried bandage is cut into lengths, collapsed, rolled, and stuffed. Bandage is grouped by its work, or number of strings per inch [57, 122]. A few sorts of careful dressing require a nearby coincided bandage for additional quality and more noteworthy security, while different uses, for example, essential wound dressings, retentive optional dressings, and bigger dressings to retain purulent issue or other seepage require gentler, progressively permeable clothes with an increasingly open structure. Different types of cushions, packs, and dressings are produced using careful bandage, alone or in association proportional with retentive cotton, tissue paper, and different materials [63, 57].

Filmated Gauze is a collapsed retentive bandage with a slender, even film of cotton or rayon appropriated over each layer [124]. This filmation cushions up and gives plentiful dressing volume, yet costs not as much as cloth alone of equal volume. It has snappy ingestion and unordinary non-abrasiveness [57]. Head recorded swabs are intended for use on surface wounds and cuts. These swabs are a multi-layer nonwoven Polyester/Viscose, which sandwiches caught cotton fiber focuses, giving improved retention [117, 126].

Nonwoven Surgical Sponges—Nonwoven textures have been built up that are reasonable options in contrast to woven cotton bandage for use in wound cleaning, wound dressing, and tissue-taking care of. These nonwoven textures rely upon thick snare of their manufactured strands (Dacron, rayon, and so forth) to give the texture an adequate elasticity moving toward that of woven cotton dressing. They ordinarily offer more prominent permeable limit than cotton dressing wipes of practically identical mass, while creating less build up. Claim to fame adaptations of the nonwoven wipes are accessible pre-fenestrated for IV tubing or channel dressing techniques [127]. One maker (Johnson and Johnson) gives both a nonwoven wipe to wound dressing (Sof-Wick: delicate surface, exceptionally permeable or Topper: exceedingly spongy, less dressing changes) and a nonwoven broadly useful purging/prep wipe (Nu Gauze: cloth like surface, more retentive than bandage) [128]. Furthermore, another general wipe which consolidates the best properties of woven and nonwoven dressing, has been made from another texture innovation. Mirasorb (Johnson and Johnson) is produced using a cotton mix, is more retentive and versatile than woven bandage, gives less adherence to solid tissue, and diminishes wound harm and tissue injury upon expulsion. Cotton balls and careful bandages (CSG) have been routinely utilized for cleaning liquids and blood in medical procedures. Polyurethane sheets (PUS) are progressively utilized rather than CSG [57], [63], [129].

Selva-Edge Gauze Strips in widths of 1/4 to 2 inches are planned uncommonly and woven for utilize both as pressing strips in medical procedure of the nose and sinuses, nasal hemostasis, and so on, and as seepage wicks in the treatment of bubbles, abscesses, fistulas, and other depleting wounds. The ravel-verification, selva edges on the two sides take out every single free string. These dressings are accessible unmedicated or sedated with 5% iodoform. These strips are realistic in sterile structure stuffed in fixed glass containers. Nu Gauze Packing Strips are bundled in polystyrene holders [127, 129].

Gauze Pads or Sponges are collapsed squares of careful cloth. These are folded to the point that no cut dressing edges or free strings are uncovered. This keeps free strands from entering the wound. The cushions are collapsed to such an extent that each size might be unfurled to bigger sizes without uncovering cut edges or free strings. Cleaned bundles of these much of the time utilized allgauze wipes are accessible in carefully designed bundles [57, 90]. Such sterile units especially are appropriate to the various plate sets arranged in medical clinics. Bandage cushions and cloth wipes are utilized in various applications and are incredible for general cleaning, dressings, preparing, pressing and debriding wounds. It can likewise be utilized as a transitory retentive dressing over wounds.

The contrast between these things are that cloth cushions accompanied one for each pack, while dressing wipes accompany at least two for every pack [130, 131].

X-ray Detectable Gauze Pads are same as all-cloth cushions yet contain additions treated with barium sulfate. Wipes utilized in medical procedure can be in charge of genuine inconvenience or even passing when they are not effectively distinguished and evacuated [133]. They are nontoxic, delicate, and nonabrasive. They remain for all time perceivable in light of the fact that they neither disintegrate in the body nor are influenced by either disinfection or time [63,136]. Instances of X-beam perceivable wipes incorporate Vistec and Kerlix (interesting, crinkleweave, delicate, and spongy), both made by Kendall [135]. X-Ray Detectable Sterile Gauze Sponges are prudent, multi-reason cloth wipes perfect for use in surgeries. They are sterile, very spongy, and their x-beam perceivability redresses inadvertent maintenance (gossypiboma) in patients. Exceptionally retentive (16-handle), fine work 100% cotton dressing is for all intents and purposes build up free. Latex free [134].

Ray-Tec X-Ray Detectable Sponges (Johnson & Johnson) contain a nonabrasive vinyl plastic monofilament that gives a trademark design in the X-beam. Composite permeable dressings have been progressed for explicit purposes. They normally comprise of layers of spongy dressing or nonwoven texture with fillers of cotton, rayon, nonwoven texture, or tissue paper in reasonable game plans. Composite wipes have dressing or nonwoven texture surfaces with fillers of cotton, rayon, nonwoven texture, or retentive tissue [63-70]. The RAY-TEC X-Rayable Sponge removed the mystery from careful accidents. On the off chance that a careful wipe was unintentionally left in a body during an activity, it could now be effectively identified by a x-beam. The spearheading wipes spared patients from conceivable exploratory medical procedure or extra pointless treatment following entanglements [136, 137].

iii. *Dressing Combines*

Dressing Combines are intended to give warmth and insurance and to assimilate enormous amounts of liquid that may deplete from an entry point or wound. Each consolidate comprises of a nonwoven texture spread encasing fiber with or without spongy tissue. They additionally may fuse a nonabsorbent layer of cotton, tissue, or plastic film to keep liquid from coming through to soil liners and bedding, however some consolidated dressings are altogether spongy [138,139]. Exorbitant draining is a confusion of wound debridement in patients accepting anticoagulation treatment. Chitosan is a straight, decidedly charged polysaccharide that has potential as a hemostatic topical dressing. Topical use of Opticell dressing with

chitosan has hemostatic impacts that could be a valuable device to control draining related with wound debridement [139, 140]. Drawn out essential anticoagulation could impede the healing procedure. A minor wound could be a day by day challenge for patients who use anticoagulants and antiplatelet drugs. Standard cloth dressings and direct pressure are regularly time-concentrated for controlling discharge for these patients [140]. In particular, the composites are antibacterial, hemostatic, biocompatible, great permeable for anticoagulated entire blood, and can keep up dampness balance for wound healing. For instance, the composites were found to hinder the development of both Gram positive and negative smaller scale life forms (counting *Escherichia coli* (ATCC 8739), *Staphylococcus aureus* (ATCC 25923), methicillin safe *S. aureus* (ATCC 33591), and vancomycin safe *Enterococcus faecalis* (ATCC 51299). They are nontoxic to fibroblasts, in particular fibroblasts were found to develop and multiply within the sight of the composites [141].

iv. *Laparotomy Sponges*

Laparotomy Sponges, otherwise called Abdominal Packs, Tape Pads or Packs, Walling-Off Mops, Stitched Pads, Quilted Pads, Gauze Mops, and so forth, are utilized to shape a nonabrasive divider that will block stomach or different organs from going into the field of activity and to help bolster body temperature during presentation. The employable control of non-compressible discharge is the single biggest effect that could be tended to in decreasing the mortality on the combat zone. Laprotomy cushions, generally utilized for drain clearing, are made of woven cotton, and, while compelling, their utilization requires a significant measure of room and includes weight. This represents no worry in conventional working rooms however is a deterrent for portable suppliers and suppliers in somber situations [142]. Be that as it may, wipes are made of four layers of 28×24 work dressing. The edges are collapsed in and trimmed. The whole pack is cross-sewed, and a circled tape 1/2-inch wide and 20-inches in length is connected to one corner. An alluring element of one sort is a X-beam perceptible addition so immovably fused into the bandage that it can't end up disconnected. Treated with barium sulfate, the monofilament is nontoxic and, were it to be left coincidentally in situ, would cause not any more outside body response than a conventional dressing [54].

v. *Sanitary Napkins*

Sanitary Napkins proposed for unique medical clinic use, also called V-Pads, Obstetrical (OB) Pads, Perineal Pads, Maternity Pads, and so on, are utilized in obstetrical, gynecological, or maternity cases. Napkins that have repellent tissue as an afterthought and back surfaces of the napkin for the most part are favored in light of their more noteworthy liquid holding limit. Clean

napkins for the most part accompany two sizes of filler, 3×9-inch or 3×11-inch. The napkin spread by and large is produced using a nonwoven texture or a nonwoven texture bolstered with an open-work scrim. Bundled, sanitized napkins are accessible and utilized by and large to decrease cross-pollution conceivable outcomes. lessening mental worry during the feminine cycle time frame is a significant personal satisfaction issue for ladies. Wearing a sterile napkin (SN) is accepted to impact mental pressure reactions of ladies during everyday living exercises [54], [71,72].

vi. *Disposable Cleaners*

Disposable Cleaners made from various types of nonwoven fabrics are available. They generally offer advantages over paper in wet strength and abrasion resistance, plus having better cleaning ability. Their advantages over cloth are reduced laundry expense and cross-contamination possibilities.

vii. *Eye Pads*

Eye Pads are experimentally molded to fit easily and spread the eye totally, accordingly securing the eyebrow when taped. These cushions are made utilizing nonwoven texture. Eye patches are commonly utilized post eye medical procedure, during the evening so as to ensure the eye and lessen inconvenience. It might possibly be utilized during the day. The Eyelid must be shut before use of the fix. An appropriately set fix must apply delicate pressure on the eye cover to stay shut and the eye itself does not rub against the fix. Since the measure of light going into the eye diminishes it helps in facilitating the inconvenience. The eye can at present move under the fix. At times if the careful wound isn't appropriately fixed, an extra day of fixing may be required. In progressively extreme cases patient may need to get the wound re-worked for appropriate conclusion. Eye patches are favored as they help in restricting the swelling. It is prudent to evacuate eye patches at whatever point conceivable to permit healing and utmost the dampness inside the fix which can likewise now and then be unsafe. Different sides are encased to keep the cotton from getting away and the cushion from contorting. Whenever wanted, the cushion might be collapsed and utilized as a pressure dressing. Eye cushions particularly are helpful in the outpatient facility of the emergency clinic, the modern therapeutic office, and the doctor's office. They are fixed in individual sterile envelopes [143-146].

viii. *Nursing Pads*

Nursing Pads are designed in a contour shape to fit comfortably under the nursing brassiere or breast binder.

ix. *Disposable Under-pads*

Disposable Under-pads are utilized for incontinent, maternity, and different patients with serious seepage. Such cushions cost not exactly the normal clinic made item and give a slick, spotless, simple to-

deal with cushion that is changed rapidly and effectively arranged [150]. The development of an underpad ought to achieve three objectives. In the first place, its sponsorship ought to have a low coefficient of grinding to counteract frictional skin wounds. Second, an internal retentive center ought to quickly contain dampness and scatter it all through the whole cushion. Third, the center and cover stock ought to effectively cooperate to hold dampness and forestall wet-back or liquid return [152]. Expendable briefs are accessible (Johnson and Johnson, Kendall). Incontinence items can be tremendously useful. They can counteract spilling onto your garments, control smell, and avoid skin disturbance [150]. Utilization of dispensable incontinence cushions diminishes medical clinic obtained pressure wounds (HAPIs) however not incontinence-related dermatitis (IAD) events [151]. Peacefulness Peach Sheet underpad can contain well more than one quart of liquid. This item accommodates excellent skin dryness, smell disposal, bacterial control and pH balance. Four sticky tape tabs secure the situating of about 5 square feet of insurance to wheelchairs, bedding and furniture. Perfect for the individuals who are in danger of skin breakdown, because of delicate skin or abnormal amounts of pee misfortune [152]. It was observed to be extremely cost effective, on the grounds that the Tuckable could stay set up over seven days without evolving [151].

x. *Cotton-Tipped Applicators*

Cotton-Tipped Applicators are used to apply medications or cleanse an area. Makes it easy to clean/culture hard to reach areas. Machine-made cotton-tipped applicators are uniform in size, resulting in no waste of cotton or medications [153]. The cotton is attached firmly to the stick and may be sterilized readily without affecting the anchorage of the cotton. They are available in 3- or 6-inch lengths [73].

d) *Bandages*

The function of bandages is to hold dressings in place by providing pressure or support. They may be inelastic, be elastic, or become rigid after shaping for immobilization. Common Gauze

i. *Roller Bandage*

Roller Bandage is listed in the USP as a form in which Absorbent Gauze may be provided. It is prepared from Type I Absorbent Gauze in various widths and lengths. Each bandage is in one continuous piece, tightly rolled and substantially free from loose threads and raveling's.

ii. *Muslin Bandage Rolls*

Muslin Bandage Rolls are made of heavier unbleached material (56 ×60 mesh). They are provided in the same widths as the typical gauze bandage. Muslin bandages are very strong and are used wherever gauze bandages do not provide sufficient strength or

support. They frequently are used to hold splints or bulky compression dressings in place [54, 57, 63].

iii. *Elastic Bandages are made in several types*

- A. *Woven Elastic Bandage* is made of heavy elastic webbing containing rubber threads. Good support and pressure are provided by this type of rubber elastic bandage [147].
- B. *Crepe Bandage* is elastic but contains no rubber. Its elasticity is owing to a particular weave that allows it to stretch to practically twice its length, even after repeated cleansings. This elasticity makes it especially serviceable in bandaging varicose veins, sprains, etc, because it conforms closely to the skin or joint surfaces, lies flat and secure, yet allows limited motion and stretches in case of swelling so that circulation is not impaired [148].
- C. *Conforming Bandage* is produced using two handles of exceptionally prepared, high caliber, 14×8-inch cotton cloth collapsed to the middle. This sort is a lot simpler to utilize and apply than standard roller swathe, since it will in general stick to itself during application, subsequently averting slipping. It promptly adjusts to all body designs without the need of turning around or contorting. A further bit of leeway is the way that there can be no unpleasant or frayed edge. Kling Conforming Gauze Bandage and Sof-King Conforming Bandage (Johnson and Johnson) are accessible in an assortment of sizes up to 6 inches wide. This bandage is utilized generally to hold dressings or supports solidly set up and once in a while as an essential dressing when adhering to the wound isn't an issue. A mercerized cotton Conforming Gauze Bandage sticks to itself and accordingly stays set up superior to anything dressing made of different materials. Sof-King is a one-utilize rayon and polyester mix wrap that gives more prominent mass to padding and more noteworthy retentiveness [20],[149], [63], [154].
- D. *High-Bulk Bandage* is made of multiple layers (typically six) of crimped cotton gauze. The high bulk of this bandage type is designed to provide padding protection in wound dressing applications. It also provides the absorbent capacity of a cotton dressing component [155]. One version (Sof-Band High Bulk, Johnson & Johnson) is made of mercerized cotton to help the bandage cling to itself, which facilitates application and improves dressing stability.
- E. *Compression Bandage* is made out of cotton weaved or woven with either thick, polyurethane, nylon, or elastane strings. Otherwise called flexible gauzes, are regularly utilized for the pressure some portion of RICE—Rest, Ice, Compression and Elevation—the best quality level of medical aid treatment for wounds and sprains. Likely the most

well-known brand name for a flexible swathe is an Ace wrap [156]. The wrap is agreeable and simple to apply. Its utilization is basically to keep up controlled degrees of pressure when pressure treatment is required. Similarly as with all pressure gauzes, these items ought to be used with alert on patients with stamped fringe ischemia or weakened blood vessel blood supply. Instances of pressure gauze incorporate Tensopress (Smith and Nephew), Yeinopress (Moliner), and Setopress (Seton Healthcare) [157].

iv. *Triangular Bandages*

Triangular Bandages usually are made by cutting a square of bleached muslin diagonally from corner to corner, forming two right triangles of equal size and shape. The length of the base is approximately 54 inches. These bandages were brought into prominence by Esmarch and still bear his name. They are used in first-aid work for head dressings, binders, and arm slings and as temporary splints for broken bones. A triangular bandage is used as an arm sling or as a pad to control bleeding. It may also be used to support or immobilize an injury to a bone or joint or as improvised padding over a painful injury. A tubular gauze bandage is used to retain a dressing on a finger or toe [158,159].

v. *Orthopedic Bandages*

Orthopedic Bandages are utilized to give immobilization and backing in the treatment of broken bones and in specific states of bones and joints. Mortar of Paris—impregnated dressing has been the ordinary material for this reason. All the more as of late presented are manufactured cast materials made of polyester cotton or fiberglass. Different sorts of plastic sheets likewise are offered that can be molded effectively and solidified to an unbending structure by cooling or substance response. These are valuable essentially for supports and restorative props. Separately bundled mortar of Paris wraps and braces are accessible in a wide assortment of sizes [53]. Flexible gauzes are regularly utilized in games to treat and anticipate sport wounds [162]. The Specialist brand (Johnson and Johnson) is produced using exceptionally treated mortar, consistently spread and solidly clung to the texture. This outcomes in a high solidarity to-weight proportion in throws produced using such gauzes. Engineered throws are connected like mortar of Paris. The utilization of an inelastic, short-extend pressure gauze following all out knee arthroplasty is a sheltered strategy that is satisfactory to patients [163]. The Delta-Lite Synthetic Casting System (Johnson and Johnson) offers both polyester, cotton texture impregnated with a polyurethane tar, and fiberglass throwing materials. Scotch cast Soft cast (3M) comprises of a sewed fiberglass substrate impregnated with a polyurethane tar containing a surface changing specialist (lessen tack, encourage application) [160]. The throws are water-

safe, light weight, and tough. Delicate, smooth and comparable to the body's shape. It is tear capable and quill capable, has low dampness maintenance, and is permeable enough to permit air course and to diminish skin disturbance. It very well may be utilized under a wide range of throwing materials, giving compelling, agreeable and safe throwing [161].

vi. *Orthoflex Elastic Plaster Bandages (Johnson & Johnson)*

Orthoflex Elastic Plaster Bandages (Johnson & Johnson) are plaster of Paris bandages containing elastic threads in the fabric and are intended for specialized prosthetic uses. Stockinette Bandages are made of stockinette material knitted or woven in tubular form without seams. Surgical stockinette is unbleached. Because it is soft and will stretch readily to conform comfortably to the arm, leg, or body, it is used to cover the skin prior to the application of a plaster of Paris or synthetic cast [165,166]. The physical properties of plaster bandages are a very important factor in achieving the basic functions of immobilization (maintaining bone fragments in the best possible position), which directly affects the speed and quality of fracture healing [164].

vii. *Cast Paddings*

Cast Paddings are soft, absorbent, protective paddings, applied like a bandage to the areas affected, before application of a cast. They are composed of various fiber constructions that conform and cling, absorb moisture, and allow the skin to breathe [74-80].

e) *Adhesive Tapes*

Surgical adhesive tapes are made in many different forms, varying both in the type of backing and in the formulation of the adhesive mass according to specific needs and requirements. The tapes available today may be divided into two broad categories: those with a rubber-based adhesive and those with an acrylate adhesive. Both types have a variety of uses. When strength of backing, superior adhesion, and economy are required (eg, athletic strapping), rubber adhesives commonly are used. Acrylate adhesives on a variety of backing materials are used widely in surgical dressing applications, when reduced skin trauma is required, as in operative and postoperative procedures; they are supplied in various strength and adhesion levels [81-85].

i. *Acrylate Adhesives*

Acrylate glues on a nonwoven or texture support have been acknowledged generally for use as careful tapes, owing to a great extent to what might be named their hypoallergenic nature. Since acrylate glues are essentially a uni-polymeric framework, they wipe out the utilization of an enormous number of segments in elastic based cements. In poly (alkyl-acrylate) cements, the ideal harmony between attachment, union, and stream properties is dictated by the selection of

monomers and the control of the polymerization responses. When the polymer is made, no other detailing or intensifying is required. Moreover, the acrylics have a phenomenal timeframe of realistic usability since they are not influenced promptly by warmth, light, or air, factors that will in general debase elastic based cements. Acrylate glues consolidate the correct parity of tack and long haul attachment. Their atomic structure allows the entry of water vapor so they are nonocclusive and along these lines when covered on a permeable sponsorship material don't cause over hydration in the stratum corneum. Horrendous reaction to careful tapes is limited significantly when tapes are developed to enable typical skin dampness to go through glue and support material. With this development, the dampness substance and quality of the horny cell layers remain generally ordinary. At the point when a permeable tape is evacuated, the planes of detachment create close to the outside of the stratum corneum, in the area of the normally desquamating cells. This permits rehashed utilization of tape over a similar site with insignificant harm to the skin. Hypoallergenic Surgical Tapes with acrylate cement are accessible with an assortment of permeable support materials. Rayon fabric material sponsorship gives a high-quality tape appropriate for fastening substantial dressings. Lighter dressing applications can be practiced with lower-quality, efficient, paperbacked careful tapes. A weaved sponsorship tape (Dermiform, Johnson and Johnson) gives a portion of the economies of paper careful tape with the quality and likeness of a material support. Different tapes include versatile fabric or froth backing materials for unique taping needs [169].

ii. *Rubber-Based Adhesives*

A second group of surgical adhesive tapes is the cloth-backed and plastic-backed rubber adhesives. These are used primarily where heavy support and a high level of adhesion are required. Modern rubber-based adhesive tape masses consist of varying mixtures of several classes of substances and are composed of an elastomer (para or pale crepe rubber in the case of natural rubber tapes, and synthetic elastomers made from polymers of isobutylene, alkyl-acrylate, or similar materials), one of several types of rosin or modified rosin, antioxidants, plasticizers and fillers, and coloring agents to give the tape the desired tint or whiteness [170].

Adhesive Tape Reactions: While skin responses some time ago were acknowledged by the medicinal calling as practically unsurprising sequelae to the utilization of sticky tape, with better comprehension of the components of such responses and advancement in research and innovation, the long-looked for target of hypo-reactivity has, in huge degree, been accomplished. Since sticky tape masses generally have comprised of heterogeneous and complex blends of

natural mixes, it isn't astounding that numerous specialists have credited sticky tape response to sensitivity. Later work, be that as it may, has demonstrated that a genuine hypersensitive reaction to the advanced sticky tape mass or its segments is a factor in just a little extent of clinical responses and that most watched responses are credited appropriately to different components, primarily mechanical disturbance and, to a lesser degree, concoction aggravation. There clearly is no critical distinction in response between patients with or without a background marked by hypersensitivity, however obvious explicit dermatitis may happen all the more promptly in people who have showed some other type of contact dermatitis. Unfriendly indications delivered by sticky tape are described by erythema, edema, papules, vesicles, and in extreme cases, desquamation. Tingling might be extraordinary, or it might be missing. The response might be exhibited promptly by patchtesting, and more often than not shows itself ahead of schedule—inside 24 to 48 hr. Typically, the response turns out to be increasingly serious the more drawn out the tape is left set up and keeps on expanding in force for quite a while after the tape is evacuated. This kind of response is durable and requires days for its total subsidence. Two particular sorts of bothering can result from the mechanical elements of expelling tape from the skin. One reaction—actuated vasodilation—is a generally nontraumatic, fleeting impact in which no genuine harm to the skin happens. A subsequent sort—skin stripping—is an awful reaction in which skin is evacuated with the tape and real harm to the epidermal layers results. Such mechanical skin evacuation is potentially the prevailing reason for clinical responses seen with the utilization of sticky tape. Substance bothering from sticky tape results when disturbing parts in the mass or sponsorship of the tape pervade the fundamental tissues of the skin. The tape development can impact the reactivity of such fixings considerably. For instance, numerous exacerbates that typically don't enter unblemished stratum corneum can infiltrate overhydrated corneum. At the point when bits of the stratum corneum are evacuated, the hindrance limit of the skin is harmed significantly. In this circumstance, any bothering parts of the tape have prepared access to fundamental tissues. These substances at that point can cause a level of aggravation that is far more prominent than would be seen on unblemished skin [81-85], [171, 172].

f) *Protectives*

Up to this point, protectives included just the different impermeable materials proposed to be utilized adjunctively with other dressing segments to anticipate the loss of dampness or warmth from a wound site or to shield garments or bed liners from wound exudate. Film dressings are phenomenal gadgets to ensure against

infection and dislodgement of vascular cannulae and seepage destinations. Also, they might be utilized to ensure unprotected zones against pressure wounds. Protectives additionally are utilized to cover wet dressings and hot or cold packs. In like manner use as protectives are plastic sheeting and waxed or plastic-covered paper. These anticipate the break of dampness or warmth from the dressing or the pack and secure attire or bed materials. Elastic sheeting is an elastic covered material, waterproof and adaptable, in different lengths and widths for use as a covering for bedding. A purported nursery sheeting is provided, covered distinctly on one side [86-88].

g) *Products for Adhesion Prevention*

Grips are unusual associations between organs or tissues that structure after injury, including medical procedure. They comprise of deliberate fibrin and fibrovascular scar tissue and convolute all regions of medical procedure. In gynecological medical procedure, bonds may result in barrenness and pelvic agony; in intestinal medical procedure they may result in intestinal hindrance; in cardiovascular medical procedure they may render a second sternotomy perilous, and in ligament medical procedure they will avert versatility. Albeit cautious tissue taking care of and great hemostasis may decrease attachment arrangement, there are not many demonstrated substances intended for the counteractive action of grips. Gynecare Interceed Absorbable Adhesion Barrier (Ethicon) is a sewn texture of oxidized recovered cellulose that is set at a site where bonds are suspected to happen. It swells and gels to frame a boundary between two adjoining surfaces, permitting re-mesothelialization to occur. The texture at that point debases horribly by around 14 days and infinitesimally by around 28 days. Interceed Barrier is shown for diminishing the occurrence of bonds in pelvic gynecological medical procedure. Other mechanical hindrances utilized for the avoidance of bonds incorporate Seprafilm (Genzyme) and Gore-Tex Surgical Membrane (Gore). More up to date items accessible for the counteractive action of postoperative bonds that are not site-explicit for application incorporate Gynecare Intergel Adhesion Prevention Solution, a ferric hyaluronate gel (Lifecore Biomedical) and Sepracoat, a weaken hyaluronic corrosive arrangement (Genzyme) [89, 90], [173,174].

h) *Operating Room Supplies*

i. *Hemostatic Products*

Hemostatic Products quicken hemostasis by giving a thrombogenic surface that advances platelet accumulation and fibrin polymerization. These topical hemostatic operators incorporate collagen, gelatin, cellulose, and thrombin. These incorporate collagen wipes and powders (Instat, Johnson and Johnson; Helistat, Integra Life Sciences; Actiofoam, Bard; Avitene, Davol; Helitene, Integra Life Sciences), gelatin wipes

(Surgifoam, Johnson and Johnson; Gelfoam, Upjohn), and Oxidized Regenerated Cellulose USP (Surgicel, Johnson and Johnson). Both oxidized cellulose and oxidized recovered cellulose are operators whose activities rely upon the development of a coagulum comprising of salts of polyanhydroglucuronic corrosive and hemoglobin. At the point when connected to a draining surface, they swell to form a darker coagulated mass that is ingested bit by bit by the tissues, as a rule inside 7 to 14 days. They are utilized in medical procedure for the control of moderate draining when suturing or ligation is unrealistic or ineffectual.

ii. *Thrombin (USP) solutions*

Thrombin (USP) solutions of bovine origin (Thrombinar, Jones Medical) promote hemostasis by catalyzing the conversion of fibrinogen to fibrin. They may be used in conjunction with fibrinogen concentrates prepared from autologous cryoprecipitate or from pooled donor blood.

iii. *Tissue sealants*

Tissue sealants are absorbable and are utilized for an assortment of signs including fixing of blood vessel punctures, fixing of air spills during pneumonic medical procedure, and supporting wound healing. The zone of tissue sealants is extending quickly, with new items achieving the market for various signs. Angio-seal (Kendall), an absorbable material, is utilized as a sealant for blood vessel punctures. AdvaSeal (Focal), an engineered absorbable sealant, is utilized to seal air spills during pneumonic medical procedure. Tissell (Immuno AE), a two-segment fibrin sealant, is utilized to advance wound healing just as accomplish hemostasis and tissue attachment. BioGlue, (Cryolife) is a cow-like egg whites based paste used to seal aortic aneurysms and anastomotic destinations.

iv. *Tissue glues*

Tissue glues are used for topical skin adhesives and replace the need for sutures, staples, or adhesive strips for certain types of lacerations requiring closely approximated wound edges. Derma bond (Closure Medical), an octyl cyanoacrylate, is used as a topical skin adhesive that sloughs from the wound as re-epithelialization of the skin occurs, providing sufficient time for wound healing. Indermil (Tyco Healthcare), a butylcyanoacrylate, is another topical skin adhesive.

v. *Disposable Sterile OR and OB Packs*

Disposable Sterile OR and OB Packs are prepared, packaged, and sterilized assemblies of diapering and gown units, designed to fulfill the operating and delivery room needs. They eliminate the problems of laundering, storage, assembly, and sterilization of muslin drapes and gowns. They introduce many special materials with particular properties of porosity; repellency to water, alcohol, blood and other fluids; abrasion resistance; and other desirable attributes.

vi. *Double packages*

Double packages of contamination-resistant paper have been developed to permit opening and use without compromising sterility. Retention of sterile characteristics until used, eliminates the need for re-sterilization. Face masks for use in the operating room and where contamination must be controlled generally are made of plied, fine-mesh gauze, shaped to cover the nose, mouth, and chin. They are laundered and autoclaved. Disposable face masks with special filtration material giving high retention of particulate matter and designed for more effective fitting are available from several manufacturers. Surgine Face Mask (Johnson & Johnson) claims a 94% filtration efficiency with high user comfort [54], [91], [175-178]

i) *Surgical Dressings*

i. *Adhesive Bandage*

Adhesive Absorbent Compress; Adhesive Absorbent Gauze: A compress of four layers of Type I absorbent gauze, or other suitable material, affixed to a film or fabric coated with a pressure-sensitive adhesive substance. It is sterile. The compact may contain an appropriate antimicrobial agent and may contain one or more suitable colors. The adhesive surface is covered by a suitable removable covering.

Description- The compress is substantially free from loose threads or ravelings; the adhesive strip may be perforated, and the back may be coated with a water-repellent film.

ii. *Gauze Bandage*

Type I absorbent gauze; contains no dye or other additives.

Description- One continuous piece, tightly rolled, in various widths and lengths and substantially free from loose threads and ravelings.

iii. *Oxidized Cellulose*

Absorbable Cellulose; Absorbable Cotton; Cellulosic Acid; Hemo-Pak (Johnson & Johnson); *Oxycel* (Deseret Medical) Sterile gauze or cotton that has been oxidized chemically to make it both hemostatic and absorbable; contains 16% to 24% carboxyl (COOH) groups.

Description- In the form of gauze or lint. Is slightly off-white in color, is acid to the taste, and has a slight charred odor.

Solubility- Insoluble in water or acids; soluble in dilute alkalis.

Comments- The estimation of oxidized cellulose in different surgeries depends on its properties of absorbability when covered in tissues and its surprising hemostatic impact. Ingestion happens between the second and seventh day following implantation of the dry material, contingent upon the sufficiency of the blood provided to the territory and the level of

concoction debasement of the embedded material. Complete retention of a lot of blood-doused dressing may take a month and a half or more, and genuine careful difficulties and sore arrangement have been accounted for as the consequence of inability to assimilate. Hemostasis relies on the stamped fondness of cellulosic corrosive for hemoglobin. At the point when presented to blood, either in vitro or in careful conditions, the oxidized cloth or cotton turns exceptionally dim darker or dark and structures a delicate coagulated mass that promptly shape itself to the forms of unpredictable surfaces and controls careful discharge by giving a falsely actuated cluster. Pressure ought to be applied on the bandage or cotton for around 2 min to encourage the fixing off of little, draining vessels.

Two factors require emphasis:

1. Cellulosic acid does not enter the physiological clotting mechanism per se but forms what might be termed an artificial clot as described and, therefore, is effective in controlling the bleeding hemophilic and
2. The hemostatic action of cellulosic acid is not enhanced by the addition of other hemostatic agents, such as thrombin (which in any case would be destroyed by the pH of the gauze unless some means of neutralization were practicable). The hemostatic effect of either one alone is greater than the combination.

It is helpful as a transitory pressing for the control of fine, venous, or little blood vessel drain, however since it restrains epithelialization, it ought to be utilized uniquely for the prompt control of discharge and not as a surface dressing. A cleaner and increasingly uniform item arranged from oxidized recovered cellulose has been created and is accessible as Surgicel Absorbable Hemostat. This offers numerous points of interest over the more seasoned, less-uniform oxidized cellulose got from cotton and, due to its compound consistency, guarantees reliable execution and defeats a large number of the challenges experienced with the more established kind of cotton item. The sewed texture strips don't part, might be sutured set up effectively if essential, and furnish instant and complete assimilation with least tissue response.

iv. *Oxidized Regenerated Cellulose*

Surgicel; Surgicel Nu-Knit; Surgicel Fibrillar (Johnson & Johnson)

Contains 18–24% carboxyl groups (COOH), calculated on the dried basis. It is sterile.

Preparation- Cellulose is dissolved and regenerated by a process similar to the manufacture of rayon, which is then oxidized.

Description- Creamy white gauze, lint, or woven material.

Solubility- Insoluble in water; soluble in alkali hydroxides.

Comments- Absorbable hemostatic.

v. *Purified cotton*

Gossypium Purificatum; Absorbent Cotton

The hair of the seed of cultivated varieties of Gossypium hirsutum Linnéor other species of Gossypium (Fam Malvaceae), freed from adhering impurities, deprived of fatty matter, bleached, and sterilized in its final container.

Description- White, soft, fine, filament-like hairs appearing under the microscope as hollow, flattened and twisted bands, striate and slightly thickened at the edges; practically odorless and practically tasteless.

Solubility- Insoluble in ordinary solvents; soluble in ammoniated cupric oxide TS.

vi. *Dextranomer*

Debrisan (Johnson & Johnson)

Dextranomer is a three-dimensional cross-linked dextran polymer prepared by interaction of dextran with epichlorohydrin.

Description- White, spherical beads, 0.1 to 0.3 mm in diameter; hydrophilic. Also available dispersed in polyethylene glycol, as a paste.

Solubility- Insoluble in water or alcohol. Each gram absorbs about 4 mL of aqueous fluid, the beads swelling and forming a gel.

Comments- Typically to cleanse secreting lesions such as venous stasis ulcers, decubitus ulcers, infected traumatic and surgical wounds, and infected burns. It absorbs the exudates, including the components that tend to impede tissue repair, and thereby retards eschar formation and keeps lesions soft and pliable.

vii. *Absorbable Dusting Powder*

Starch-derivative Dusting Powder

An absorbable powder prepared by processing cornstarch and intended for use as a lubricant for surgical gloves; contains not more than 2% magnesium oxide.

Description- White, odorless powder; pH (1 in 10 suspension) between 10 and 10.8.

viii. *Absorbent Gauze*

Carbasus Absorbens; Gauze

Cotton, or a mixture of cotton and not more than 53.0%, by weight, of purified rayon, in the form of a plain-woven cloth. If rendered sterile, it is packaged to protect it from contamination. *Description*—White cotton cloth of various thread counts and weights; may be supplied in various lengths and widths and in the form of rolls or folds.

ix. *Purified Rayon*

A fibrous form of bleached, regenerated cellulose. It may contain no more than 1.25% titanium dioxide.

Preparation- By the viscose rayon process.

Description- White, lustrous or dull, fine, soft, filamentous fibers, appearing under the microscope as round, oval, or slightly flattened translucent rods, straight or crimped, striate and with serrate cross-sectional edges; practically odorless and practically tasteless.

Solubility- Very soluble in ammoniated cupric oxide TS or dilute H₂SO₄ (3 in 5); insoluble in ordinary solvents.

Comments- Hemostatic.

x. *Adhesive Tape*

Sterile Adhesive Tape

Fabric and/or film evenly coated on one side with a pressure-sensitive, adhesive mixture. If rendered sterile, it is protected from contamination by appropriate packaging [20], [25], [57], [63], [66], [74], [92-97].

different prescriptions, including over-the-counter drugs. For instance, fluoroquinolones, for example, ciprofloxacin, will tie to calcium, magnesium and iron in the gastrointestinal framework and diminish the assimilation and adequacy of the anti-microbial. Pharmacists need to encourage patients to stop their multi-nutrients and additionally acid neutralizers when taking fluoroquinolones, or to take the items at isolated occasions of the day. Once more, high portions of anti-toxins may cause gastric aggravation, could be avoided by PPIs. By perceiving venous skin changes (for instance), realizing when to allude patients to their essential consideration suppliers and perceiving the patient's status to change, pharmacists can assume an indispensable job in supporting the convenient survey and reassessment of wounds. The needs (and the board plans) of the patient can change quickly and definitely, especially in the perplexing instance of wounds where there are a few simultaneous co-sullen conditions. The avoidance and treatment of wounds is mind boggling, regularly multi-factorial in starting point and best polished through a composed, inter-professional group approach. This work together tie approach postures difficulties, for example, encouraging correspondence when colleagues are physically situated in discrete workplaces and work various hours [98,99], [179-182]. For quick reference, a quick wound care chart is detailed in Table 2 and barriers to effective wound care in Table 3.

IV. PHARMACIST'S ROLE

In customary practice, it is frequently during these far reaching prescription audits that a drug specialist should ready to evaluate the patient's hazard factors for wounds and note any early indications of venous trade off. Network pharmacists are regularly in contact with the patient or patient's delegate every now and again as the patient will in general come back to the drug store a few times each month for an assortment of human services needs. The most evident commitment of a network drug specialist would be in the administration of neighborhood wound consideration, explicitly the administration of aggravation and infection control. Numerous anti-infection agents will meddle with

Table 2: Quick Reference Wound Care Chart [100]

Dressing Type	Wound Type	When to change dressing	Comments/Expectations
Paraffin dressings <ul style="list-style-type: none"> Cuticerin Jelonet Bactigras 	<ul style="list-style-type: none"> Superficial, clean, minor abrasions Skin tears that require review within 24 to 48 hours 	<ul style="list-style-type: none"> 2 days 	<ul style="list-style-type: none"> Prevent maceration Granulation Protect epithelizing wound
Long wearing impregnated mesh <ul style="list-style-type: none"> Mepitel (silicone) 	<ul style="list-style-type: none"> Epidermal and clean superficial dermal Epithelised burns 	<ul style="list-style-type: none"> 7 days 	<ul style="list-style-type: none"> Healing Epithelization
Absorbent Pads <ul style="list-style-type: none"> Mesasorb Zetuvit 	<ul style="list-style-type: none"> Oozing wounds Secondary dressing to absorb exudate 	<ul style="list-style-type: none"> PRN 3 days 	<ul style="list-style-type: none"> Absorb fluid Prevent maceration
Films <ul style="list-style-type: none"> Tegaderm Opsite 	<ul style="list-style-type: none"> Secondary dressing for burns and sutures Stage I pressure injuries 	<ul style="list-style-type: none"> 2 to 7 days "A week or a leak" 	<ul style="list-style-type: none"> Not to be used on infected wounds Keep dressings intact Keep dressings dry Water resistant not waterproof Protect against sheer and friction
Foams <ul style="list-style-type: none"> Allevyn Lyofoam 	<ul style="list-style-type: none"> Pressure injuries Ulcers Toe wounds 	<ul style="list-style-type: none"> 2 to 5 days When exudate is 1 cm from the edge of dressing 	<ul style="list-style-type: none"> Protection Draws fluid from wound preventing maceration and promoting healing

Soft silicone <ul style="list-style-type: none"> Mepilex border 	<ul style="list-style-type: none"> Skin tears Epithelising wounds Superficial dermal burns Epithelized burns 	<ul style="list-style-type: none"> 2 to 7 days (depending on skin tear) 	<ul style="list-style-type: none"> Healing Protect epithelializing tissue Prevent maceration Ensure hemostasis prior
Hydrocolloids <ul style="list-style-type: none"> Duoderm Duoderm extra thin Comfeel 	<ul style="list-style-type: none"> Stage I pressure injuries Superficial dermal burns 	<ul style="list-style-type: none"> 2 to 5 days Change if leakage occurs or rolled up	<ul style="list-style-type: none"> For wounds with minimal exudate Prevent maceration Protection
Hydrocolloid paste <ul style="list-style-type: none"> Comfeel paste 	<ul style="list-style-type: none"> Pressure injuries with slough Ulcers with slough 	<ul style="list-style-type: none"> 2 days 	<ul style="list-style-type: none"> De-slough Re-hydrate Fills dead space
Hydrogels <ul style="list-style-type: none"> Solugel Intrasite gel Purilon 	<ul style="list-style-type: none"> Necrotic ulcers Pressure injuries with slough 	<ul style="list-style-type: none"> Daily to 2 days 	<ul style="list-style-type: none"> Rehydrate De-slough Prevent maceration
Calcium Alginate <ul style="list-style-type: none"> Kaltostat Kaltostat rope 	<ul style="list-style-type: none"> Promote haemostasis Bleeding ulcers Bleeding skin tears Around bleeding insertion sites 	<ul style="list-style-type: none"> 24 hours (until haemostasis is achieved) 	<ul style="list-style-type: none"> Prevent trauma (soak off if stuck to wound) Haemostasis Absorption (absorbs up to 20 times its weight in fluid) Granulation
Hydrofibre <ul style="list-style-type: none"> Aquacel Durofibre 	<ul style="list-style-type: none"> Moderately to heavily exudating wounds Not bleeding wounds 	<ul style="list-style-type: none"> 2nd to 3rd daily 	<ul style="list-style-type: none"> Prevent maceration Maintain a moist wound environment
Cadexomer Iodine <ul style="list-style-type: none"> Iodosorb 	<ul style="list-style-type: none"> Heavily colonised infected wounds Infected or slow to progress wounds 	<ul style="list-style-type: none"> 2nd to 3rd daily In severe cases may need to be dressed daily	<ul style="list-style-type: none"> Check for iodine/shellfish sensitivities Change to paste when powder begins to crust up Remove all product before reapplication Can be painful on application

V. FACTORS DELAYING WOUND HEALING

Holistic assessment of the patient is an important part of the wound care process. A number of local and systemic factors can delay or impair wound healing. These may include:

- **Malnutrition:** Inadequate supply of protein, carbohydrates, fatty acids, and trace elements essential for all phases of wound healing
- **Reduced Blood supply:** Cardiovascular disorders and Ischemia
- **Medication:** Non-steroidal anti-inflammatory drugs and Corticosteroids.
- **Chemotherapy:** Suppresses the immune system and inflammatory response
- **Radiotherapy:** Increases production of free radical which damage cells
- **Psychological stress and lack of sleep:** Increase risk of infection and delayed healing
- **Obesity :** Decreases tissue perfusion
- **Infection:** Prolong inflammatory phase, use vital nutrients, impair epithelialization and release toxins

- **Reduced wound temperature:** Prolonged dressing changes or use of cold cleansing products.
- **Underlying Disease:** Diabetes Mellitus and Autoimmune disorders
- **Maceration:** Excess wound exudates or contact with bodily fluids reduces wound tensile strength
- Inappropriate wound management
- Patient compliance
- Unrelieved pressure
- Immobility
- Substance abuse including alcohol and cigarette smoke [101], [111-115]

Table 3: Barriers to effective treatment of wounds [109]

Barriers	Examples
Educational factor	Poor quality of research, lack of appropriate training, ritualistic practice and lack of appropriate skills.
Organizational factor	Lack of standardisation of practice that is acceptable, lack of expert opinion, instability in the health services.
Clinical factor	Bacterial infection, hypersensitivity,

Barriers	Examples
	malnutrition, poor tissue perfusion, copious exudate, too much or too little information on wound management.
Psychosocial factor	Social isolation resulting in depression and reduced motivation with treatment, pain resulting in loss of sufficient sleep and lack of self-care.

VI. FUTURE DIRECTIONS

The improvement of new and compelling intercessions in wound consideration remains a zone of exceptional research. Negative pressure wound treatment has without a doubt changed wound consideration starting now and into the foreseeable future and has demonstrated gainful for an assortment of wounds. Hydro conductive dressings are another class that is rising with concentrates in progress. Different modalities, for example, hyperbaric oxygen, development factors, biologic dressings, skin substitutes, and regenerative materials have likewise demonstrated effective in propelling the wound-healing procedure through an assortment of components. The eventual fate of wound healing now stays obscure. Barely any high caliber, randomized controlled preliminaries assessing wound dressings exist and don't obviously exhibit predominance of numerous materials or classifications. Relative viability research can be utilized as an instrument to assess topical treatment for wound consideration moving into what's to come. Until further information develop, instruction on the accessible items and legitimate clinical idea must win.

VII. CONCLUSION

Wounds will effectively secure microscopic organisms, except if steady measures are taken. The bacterial security managed by average retentive cellulose dressings has been demonstrated to be confined, especially within the sight of serous exudate that may jeopardize dressing respectability. Moreover, dressings may shed particles that wait the wound. On the other hand, numerous cutting edge dressings are impenetrable to microscopic organisms, are dispensed with totally, have been found to improve re-epithelialization rates and decrease the event of wound sepsis. As of late, it has been discovered that they could likewise assume a job in forestalling cross-tainting. Expelling run of the mill cellulosic dressings from bacterially colonized wounds frees wound microscopic organisms into the air, and the numbers are moderate to decay. Notwithstanding, utilizing an in vitro wound model, utilization of the hydrocolloid dressing on tentatively colonized wounds brought about altogether less quantities of airborne microbes. Dispersal from wet run of the mill dressings was lower than from dry dressings; in any case, the quantities of microscopic

organisms per liter of air following removal of the hydrocolloid dressing were roughly 20% of those watched for cloth. These discoveries have additionally been settled in the facility. To diminish the rate of complexities, damage care when all is said in done, and infection control approaches specifically, requires deliberately restrained cooperation.

ACKNOWLEDGEMENT

It's an extraordinary appreciation and respect to be a piece of medicinal services research and training. I am appreciative to Dr. Sarwar Ahmed Sobhan, General and Laparoscopic Surgeon; Hepatobiliary, Pancreatic and Colorectal Surgeon, BSMMU, Shahbag Dhaka for his valuable time to audit my paper. Likewise, I'm thankful to workshop Library of Faculty of Pharmacy, University of Dhaka and BANSDOC, National Scientific and Technical Documentation Center for the books, diaries and pamphlets. The best assistance was from my understudies who paid enthusiasm for my point as class address and urged to compose such article on careful improvement. Notwithstanding an incredible shortage of subsidizing this reason from any expert, the experience was adequate to continue look into.

Compliance with the Ethical Issues

- *Ethics approval and consent to participate*

Animal and Human experiment: N/A

Human Data Submission Approval: N/A

- *Consent for publication*

Consent to publish Individual Person's data: N/A

- *Availability of data and materials*

Data sharing: Please contact author for data requests.

- *Competing interests*

The authors declare that they have no competing interests

- *Funding*

Funding from individual/Organization: N/A

- *Authors' contributions*

The individual contributions of authors: N/A

REFERENCES RÉFÉRENCES REFERENCIAS

1. Eming S A, Martin P, Tomic-Canic M. Wound repair and regeneration: mechanisms, signaling, and translation. *Sci Transl Med.* 2014 Dec 3; 6(265): 265sr6. doi: 10.1126/scitranslmed.3009337. Review. PubMed PMID: 25473038; PubMed Central PMCID: PMC4973620.
2. Nagle S M, Wilbraham S C. Wound Assessment. [Updated 2019 Jan 19]. In: Stat Pearls [Internet]. Treasure Island (FL): Stat Pearls Publishing; 2018 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK482198/>
3. Dhivya S, Padma V V, Santhini E. Wound dressings - a review. *Biomedicine (Taipei).* 2015 Dec; 5(4): 22.

- doi: 10.7603/s40681-015-0022-9. Epub 2015 Nov 28. PubMed PMID: 26615539; PubMed Central PMCID: PMC4662938.
4. Thiruvoth F M, Mohapatra D P, Sivakumar D K, Chittoria R K, Nandhagopal V. Current concepts in the physiology of adult wound healing. *Plast Aesthet Res* 2015; 2: 250-6. Available from: <http://parjournal.net/article/view/1211>.
 5. Dreifke M B, Jayasuriya A A, Jayasuriya A C. Current wound healing procedures and potential care. *Mater Sci Eng C Mater Biol Appl*. 2015 Mar; 48: 651-62. doi: 10.1016/j.msec.2014.12.068. Epub 2014 Dec 19. Review. PubMed PMID: 25579968; PubMed Central PMCID: PMC4443476.
 6. Takeo M, Lee W, Ito M. Wound healing and skin regeneration. *Cold Spring Harb Perspect Med*. 2015 Jan 5; 5(1): a023267. doi: 10.1101/cshperspect.a023267. Review. PubMed PMID: 25561722; PubMed Central PMCID: PMC4292081.
 7. Nunan R, Harding K G, Martin P. Clinical challenges of chronic wounds: searching for an optimal animal model to recapitulate their complexity. *Dis Model Mech*. 2014 Nov; 7(11): 1205-13. doi: 10.1242/dmm.016782. PubMed PMID: 25359790; PubMed Central PMCID: PMC4213725.
 8. Global guidelines on the prevention of surgical site infection. Web WHO. Available From: <https://www.who.int/gpsc/ssi-guidelines/en/>
 9. FDA Executive Summary. Classification of Wound Dressings Combined with Drugs. Prepared for the Meeting of the General and Plastic Surgery Devices Advisory Panel September 20 -21, 2016. Available From: <https://www.fda.gov/ucm/groups/fdagov-public/@fdagov-afda-adcom/documents/document/ucm518494.pdf>
 10. Dumville J C, Gray T A, Walter C J, Sharp C A, Page T. Dressings for the prevention of surgical site infection. *Cochrane Database Syst Rev*. 2014 Sep 1; (9): CD003091. doi: 10.1002/14651858.CD003091.pub3. Review. Update in: *Cochrane Database Syst Rev*. 2016 Dec 20;12 :CD003091. PubMed PMID: 25178020.
 11. Gupta A, Rattan V, Rai S. Efficacy of Chitosan in promoting wound healing in extraction socket: A prospective study. *J Oral Biol Craniofac Res*. 2019 Jan-Mar;9(1):91-95. doi: 10.1016/j.jobcr.2018.11.001. Epub 2018 Nov 9. PubMed PMID: 30456164; PubMed Central PMCID: PMC6234520.
 12. USL Medical. Wound Management Guide. Available From: <https://www.usl.co.nz/file/open/1531/>
 13. Dabiri G, Damstetter E, Phillips T. Choosing a Wound Dressing Based on Common Wound Characteristics. *Adv Wound Care (New Rochelle)*. 2016 Jan 1; 5(1): 32-41. Review. PubMed PMID: 26858913; PubMed Central PMCID: PMC4717498.
 14. Britto E J, Morrison C A. Wound Dressings. [Updated 2019 Jan 19]. In: Stat Pearls [Internet]. Treasure Island (FL): Stat Pearls Publishing; 2018 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470199/>.
 15. Jones V, Grey J E, Harding K G. Wound dressings. *BMJ*. 2006 Apr 1; 332(7544): 777-80. Review. PubMed PMID: 16575081; PubMed Central PMCID: PMC1420733.
 16. Rando T, Kang A C, Guerin M, Boylan J, Dyer A. Simplifying wound dressing selection for residential aged care. *J Wound Care*. 2018 Aug 2; 27(8): 504-511. doi: 10.12968/jowc.2018.27.8.504. Pub Med PMID: 30086249.
 17. Broussard K C, Powers J G. Wound dressings: selecting the most appropriate type. *Am J Clin Dermatol*. 2013 Dec; 14(6): 449-59. doi: 10.1007/s40257-013-0046-4. Review. PubMed PMID: 24062083.
 18. Negut I, Grumezescu V, Grumezescu AM. Treatment Strategies for Infected Wounds. *Molecules*. 2018 Sep 18; 23(9). pii: E2392. doi: 10.3390/molecules23092392. Review. PubMed PMID: 30231567; PubMed Central PMCID: PMC6225154.
 19. Healthcare Rehabilitation. What dressing for what wound. Available From: <https://www.pharmac.govt.nz/assets/ss-wounds-5-and-6-product-selection-what-dressing-for-what-wound-prue-lennox.pdf>
 20. Cooper and Gunn's Dispensing for Pharmaceutical Students by S. J. Carter. Publisher: CBS Publishers & Distributors; 12th edition (December 1, 2008) ISBN-10: 8123909020 ISBN-13: 978-8123909028
 21. Najibi S, Banglmeier R, Matta J, Tannast M. Material properties of common suture materials in orthopaedic surgery. *Iowa Orthop J*. 2010; 30: 84-8.
 22. Khiste S V, Ranganath V, Nichani A S. Evaluation of tensile strength of surgical synthetic absorbable suture materials: an in vitro study. *J Periodontal Implant Sci*. 2013 Jun; 43(3): 130-5. doi: 10.5051/jpis.2013.43.3.130. Epub 2013 Jun 30. PubMed PMID: 23837127; PubMed Central PMCID: PMC3701834.
 23. ETHICON Wound Closure Manual. Available From: http://www.ups.upenn.edu/surgery/Education/facilities/measey/Wound_Closure_Manual.pdf
 24. Basu P, Narendrakumar U, Arunachalam R, Devi S, Manjubala I. Characterization and Evaluation of Carboxymethyl Cellulose-Based Films for Healing of Full-Thickness Wounds in Normal and Diabetic Rats. *ACS Omega*. 2018 Oct 31; 3(10): 12622-12632. doi: 10.1021/acsomega.8b02015. Epub 2018 Oct 4. PubMed PMID: 30411013; PubMed Central PMCID: PMC6217532.
 25. Sood A, Granick M S, Tomaselli N L. Wound Dressings and Comparative Effectiveness Data. *Adv Wound Care (New Rochelle)*. 2014 Aug 1; 3(8): 511-529. Review. PubMed PMID: 25126472; PubMed Central PMCID: PMC4121107.

26. Wound Dressings. Web WoundEducators.com. Available From: <https://woundeducators.com/resources/wound-dressings/>
27. National Collaborating Centre for Women's and Children's Health (UK). Surgical Site Infection: Prevention and Treatment of Surgical Site Infection. London: RCOG Press; 2008 Oct. (NICE Clinical Guidelines, No. 74.) Appendix C, Wound dressings for surgical site infection prevention. Available From: <https://www.nice.org.uk/guidance/cg74/evidence/full-guideline-242005933>
28. Unfallchirurg H T. Wound dressings. Overview and classification. Unfallchirurg 2012 Sep; 115(9): 774-82. doi: 10.1007/s00113-012-2209-9.
29. Wound Home Skills Kit: Surgical Wounds. American College of Surgeons Division of Education. Available From: https://www.facs.org/~media/files/education/patient%20ed/wound_surgical.ashx
30. 4 Types of Impregnated Wound Dressings. Web Advanced Tissue Dec 22, 2015. Available From: <https://advancedtissue.com/2015/12/4-types-of-impregnated-wound-dressings/>
31. Impregnated Dressings. Web Wound Source. Available from: <https://www.woundsource.com/product-category/dressings/impregnated-dressings>
32. Web Amazon. <https://www.amazon.in/Johnsons-K-Y-Personal-Lubricant-Jelly/dp/B01B00220K>
33. Web Amazon. <https://www.amazon.com/Personal-Lubricant-Jelly-Johnson-3pack/dp/B00QP3YO94>
34. Wound care swabs. Web Mölnlycke Health Care AB. Available From: <https://www.molnlycke.us/products-solutions/mesalt/>
35. Dabi S, Haddock T. Adaptive Transparent Film Dressings. Johnson & Johnson Worldwide Absorbent Products and Materials Research. Available From: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.843.3479&rep=rep1&type=pdf>
36. Technology and product reviews. Understanding film dressings. Available From: <https://www.vitalitymedical.com/pdf/understanding-film-dressings.pdf>
37. Transparent Adhesive. Web Amazon. Available From: <https://www.amazon.com/Smith-Nephew-Flexifix-Transparent-Adhesive/dp/B0015TI2ZA>
38. Transparent Dressings. Web Health products for you. Available From: <https://www.healthproductsforyou.com/c-transparent-dressings.html>
39. Transparent Dressings. Web Vitality Medical. Available From: <https://www.vitalitymedical.com/transparent-dressings.html>
40. Edwards H, Gibb M, Finlayson K, Jensen R. Wound Dressing Guide. Project funded by the Australian Government Department of Health and EBPA program. Available From: https://cms.qut.edu.au/_data/assets/pdf_file/0003/451767/Book2-wound-dressing-guide.pdf
41. Ouchless Non-Adherent Pad TELFA Ouchless 1961 |2 x 3 Inch Non Adherent Pad by Covidien. Available From: <https://www.vitalitymedical.com/telfa-ouchless-1961-2-x-3-inch-non-adherent-pad-by-covidien.html>
42. MELOLIN Absorbent Dressings. Available From: <http://www.smith-nephew.com/anz/producthome/melolin/>
43. Understanding Composite Wound Dressings. Web Wound Educators. September 12, 2013. Available From: <https://woundeducators.com/wound-dressings-composites/>
44. SMTL Dressings Data card. Available From: <http://www.dressings.org/Dressings/geliperm.html>
45. Amtech. WC983-Johnson and Johnson Nu-Gel Hydrogel with Alginate15gram. Available From: <https://www.amtech.co.nz/shop/Medical+Consumables+MEDSU/Woundcare+Dressings+MEDSUWCARE/Ointments+and+Gels+MEDSUWCARE/OINTMENTS+AND+GELS/WC983Johnson+and+Johnson+NuGel+Hydrogel+with+Alginate15gram+WC983.html>
46. Elasto-Gel Hydrogel Occlusive Dressing, Elastogel Drs Gel 4X4 in. Web United Health Supply. Available From: http://www.unitedhealthsupply.com/Elasto-Gel-Hydrogel-Occlusive-Dressing-Elastogel-Drs-Gel-4X4-in-1-BOX-5-EACH_p_49330.html
47. Advanced Tissue. Your Guide to Hydrocolloid Dressings. Wound care products Dec 05, 2014.
48. Advanced Tissue. 4 Unique Wound Healing Benefits of Hydrocolloid Dressings. Wound dressings Mar 20, 2015.
49. Wound Care advisor. What you need to know about hydrocolloid dressings. Available From: <https://woundcareadvisor.com/hydrocolloid-dressings/>
50. Comfeel® Plus Ensured protection. Faster healing.*Hydrocolloid dressings for moist wound healing. Available From: <https://www.coloplast.com.au/wound/wound-care/comfeel-plus/>
51. REPLICARE hydrocolloid dressing. Available From: <http://www.smith-nephew.com/professional/products/advanced-wound-management/repicare/>
52. Baranoski S, Ayello E A. Chapter 9. Wound Treatment Options. In: Wound Care Essentials: Practice Principles by Sharon Baranoski, Elizabeth A. Ayello. LWW medical book collection. Publisher: Lippincott Williams & Wilkins, 2008.
53. Barbolt T A, Liu S H. Chapter 109. Surgical Supplies. In: Remington: The Science and Practice of Pharmacy - 22nd edition by Loyd V. Ed. Allen. Publisher: Pharmaceutical Press, Published: 2013.
54. Chapter 8. Dressing Materials. In: Wound Healing and Ulcers of the Skin: Diagnosis and Therapy - The Practical Approach by Avi Shai, Howard I. Maibach. Publisher: Springer Science & Business Media, 2004.

55. Wietlisbach C M. Wound Care. In: *Fundamentals of Hand Therapy*, 2nd Edition by Cynthia Cooper. Published by Elsevier B.V. 2014.
56. Kujath P, Michelsen A. Wounds - from physiology to wound dressing. *Dtsch Arztebl Int*. 2008; 105(13): 239-48.
57. *Manual of Medical-Surgical Nursing*, 7th Edition by Frances Donovan Monahan, PhD, RN, ANEF, Marianne Neighbors, EdD, RN and Carol Green, PhD, MN, RN, CNE.
58. Dziejwski P, James S, Taylor D, Bosanquet N, Cutting K, West P. Modern dressings: healing surgical wounds by secondary intention. *Hosp Med*. 2003 Sep; 64(9): 543-7. Review. PubMed PMID: 14521072.
59. ELLIOTT J R. The development of cotton wool as a wound dressing. *Med Hist*. 1957; 1(4): 362-6.
60. <https://www.indiamart.com/proddetail/absorbant-cotton-wool-7265770873.html>
61. Chapter 13. Surgical Cotton and bandages. In: *Handbook on Medical and Surgical Disposable Products* by NPCS Board of Consultants & Engineers. Publisher: Niir Project Consultancy Services, 2014.
62. Dinah F, Adhikari A. Gauze packing of open surgical wounds: empirical or evidence-based practice?. *Ann R Coll Surg Engl*. 2006; 88(1): 33-6.
63. Kaplan M, Iyiköşker Hİ. A new complication of retained surgical gauze: development of malignant fibrous histiocytoma--report of a case with a literature review. *World J Surg Oncol*. 2012; 10: 139. Published 2012 Jul 9. doi:10.1186/1477-7819-10-139
64. Ali Algadiem E, Aleisa A A, Alsubaie H I, Buhlaiqah N R, Algadeeb J B, Alsneini H A. Blood Loss Estimation Using Gauze Visual Analogue. *Trauma Mon*. 2016; 21(2): e34131. Published 2016 May 3. doi:10.5812/traumamon.34131.
65. *Practice of Pharmacy: A Treatise on the Preparing, Standardizing, and Dispensing of Official, Unofficial, and Extemporaneous Pharmaceutical Products, with Descriptions of Medicinal Substances, Their Properties, Uses and Doses*. Authors: Joseph Price Remington, Ernest Fullerton Cook, Eric Wentworth Martin, Publisher: Mack, 1951.
66. *Medical-Surgical Care Planning* Publication Year: 2003 Edition: 4th Ed. Authors/Editor: Holloway, Nancy M. Publisher: Lippincott Williams & Wilkins (LWW) ISBN: 978-1-582-55224-8.
67. *Nursing the Surgical Patient* 3rd Edition Editors: Rosie Pudner eBook ISBN: 9780702044120 Paperback ISBN: 9780702030628 Imprint: Baillière Tindall Published Date: 10th March 2010.
68. <http://sonitechnicaltextiles.com/portfolio-view/gauze-swabpadsponge/>
69. <http://www.medicarehygiene.net/surgical-gauze.html>
70. Mohamed N G, Abidin N Z, Law K S, et al. The effect of wearing sanitary napkins of different thicknesses on physiological and psychological responses in Muslim females. *J Physiol Anthropol*. 2014; 33(1): 28. Published 2014 Sep 4. doi:10.1186/1880-6805-33-28.
71. *Investment Opportunities in India: Sanitary Napkin Project (Reasons for Investment, Core Project Financials, Potential Buyers, Market Size & Analysis)* by: NIIR Project Consultancy Services (NPCS) Team, Publisher: Niir Project Consultancy Services, 2014.
72. Chapter 7. Factors that Predispose the Ear to Otitis Externa. In: *Small Animal Ear Diseases - E-Book: An Illustrated Guide* by Louis N. Gotthelf. Publisher: Elsevier Health Sciences, 2004.
73. *Pye's Elementary Bandaging and Surgical Dressing: With Directions Concerning the Immediate Treatment of Cases of Emergency, for the Use of Dressers and Nurses*, Revised from H.W. Carson's Tenth Edition of Pye's Surgical Handicraft Author A. J. Cokkinis Edition 15 Publisher John Wright, 1933.
74. *Surgical Dressings and Wound Management* by Stephen Thomas Medetech Publisher: Cardiff 2010.
75. *Surgical Wound Healing and Management*, Second Edition Hardcover – Import, 24 Sep 2012 by Mark S. Granick (Editor), Luc Teot (Editor).
76. *Surgical dressings, ligatures and sutures Volume 6 of Pharmaceutical monographs* by Francis Fish, John Owen Dawson, Publisher: Publisher Heinemann Medical, 1967.
77. *On bandaging and other operations of minor surgery* by Fitzwilliam Sargent, Publisher: Publisher: Blanchard and Lea, 1862.
78. Shim H S, Choi J S, Kim S W. A Role for Postoperative Negative Pressure Wound Therapy in Multitissue Hand Injuries. *Biomed Res Int*. 2018; 2018: 3629643. Published 2018 Mar 26. doi:10.1155/2018/3629643.
79. <https://www.verywellhealth.com/surgical-bandage-change-instructions-3156920>
80. *Surgical dressings and wound healing: proceedings of a symposium held on 7-8 July 1970 at the University of Bradford*; edited by K. J. Harkiss London, Crosby Lockwood [for] Bradford University Press, 1971 153 p. illus. 23 cm. ISBN 0258968397.
81. *Surgical Dressing Maker Work Log: Work Journal, Work Diary, Log - 126 Pages, 6 X 9 Inches Orange Logs/Work Log Series* Author Orange Logs Publisher Create Space Independent Publishing Platform, 2018 ISBN 1718726422, 9781718726420.
82. *Adhesives and Adhesive Tapes* by Gerhard Gierenz, Werner Karmann, Published by John Wiley & Sons, 2008.
83. *Pressure sensitive adhesive tapes: a guide to their function, design, manufacture, and use* by John

- Johnston, Pressure Sensitive Tape Council, Publisher: Pressure Sensitive Tape Council, 2000.
84. Adhesive Medical Tape. Available From: <https://www.buyemp.com/category/adhesive-medical-tape>
 85. The Mechanics of Surgery Issue 1 of Norman surgery series by Charles Truax, Publisher: Norman Publishing, 1988.
 86. First Aid. Available From: <http://files.hostgator.co.in/hostgator249427/file/531103455modified-copy.pdf>
 87. Powers J G, Morton L M, Phillips T J. Dressings for chronic wounds. *Dermatol Ther.* 2013 May-Jun; 26(3): 197-206. doi: 10.1111/dth.12055. Review. PubMed PMID: 23742280.
 88. Global Guidelines for the Prevention of Surgical Site Infection. Geneva: World Health Organization; 2016. Appendix 26, Summary of the systematic review on advanced dressings.
 89. Biofilm Eradication and Prevention: A Pharmaceutical Approach to Medical Device Infections by Tamilvanan Shunmugaperumal, Publisher: Publisher: John Wiley & Sons, 2010
 90. Chapter 2. General Surgical Supplies and post-operative splints. In: *Surgical Equipment and Supplies* by Colleen J Rutherford, published by F.A. Davis, May 12, 2016, pp 55.
 91. Hybrid Polymer Composite Materials: Properties and Characterisation by Vijay Kumar Thakur, Manju Kumari Thakur, Asokan Pappu, Published by Elsevier Science, 2017.
 92. Nanocellulose Polymer Nanocomposites: Fundamentals and Applications by Vijay Kumar Thakur, Published by Wiley, Oct 28, 2014.
 93. USP DI Volume 1 Drug Information for the Healthcare Professional (USP DI: v.1 Drug Information for the Health Care Professional) by Thomson PDR, Micromedex, Publisher: Thomson Healthcare; 25th edition (January 2005).
 94. Lewis S Medical Surgical Nursing by Chintamani, Publisher: Elsevier India, 2011.
 95. First Aid Manual by D K, Publisher: Dorling Kindersley Limited, 2016.
 96. Biocompatible Polymeric Materials and Tourniquets for Wounds. *Topics in Applied Chemistry* by Jan W. Gooch, Publisher: Springer, 2010.
 97. Bushra R, Aslam N, Khan A Y. Food-drug interactions. *Oman Med J.* 2011; 26(2): 77-83.
 98. Jin S. A Day in the Life: A Pharmacist's Role in Wound Care. *The Wound Management Team Wound Care Canada* Volume 13, Number 2· Fall 2015 Page 34-37.
 99. Polk R E. Drug-drug interactions with ciprofloxacin and other fluoroquinolones. *Am J Med.* 1989 Nov 30; 87(5A): 76S-81S. Review. PubMed PMID: 2686430.
 100. Vilella M. Quick Reference Wound Care Chart. The Benchmark Group, Wound Management in the Primary Health Care Program. 16 November 2014. Available from Wound care. Web The Royal Children Hospital Melbourne. Available From: https://www.rch.org.au/rchcpg/hospital_clinical_guideline_index/wound_care/
 101. Chapter 22. Surgical Wound Care. In: Kim Cooper, Kelly Gosnell. *Foundations of Nursing E-Book*. Publisher: Elsevier Health Sciences, Oct 23, 2018.
 102. Laurie S. Wound dressing selection: types and usage 2011; Available from <http://www.woundsource.com/blog/wound-dressing-selectiontypes-and-usage>
 103. Kunugiza Y, Tomita T, Moritomo H, Yoshikawa H. A hydrocellular foam dressing versus gauze: effects on the healing of rat excisional wounds. *J Wound Care.* 2010 Jan; 19(1): 10-4. PubMed PMID: 20081568.
 104. Yamane T, Nakagami G, Yoshino S, Muramatsu A, Matsui S, Oishi Y, Kanazawa T, Minematsu T, Sanada H. Hydrocellular foam dressing promotes wound healing along with increases in hyaluronan synthase 3 and PPAR α gene expression in epidermis. *PLoS One.* 2013 Aug 22; 8(8): e73988. doi: 10.1371/journal.pone.0073988. eCollection 2013. PubMed PMID: 23991211; PubMed Central PMCID: PMC3750004.
 105. Yamane T, Nakagami G, Yoshino S, Shimura M, Kitamura A, Kobayashi-Hattori K, Oishi Y, Nishijima Y, Minematsu T, Sanada H. Hydrocellular foam dressings promote wound healing associated with decrease in inflammation in rat periwound skin and granulation tissue, compared with hydrocolloid dressings. *Biosci Biotechnol Biochem.* 2015; 79(2): 185-9. doi: 10.1080/09168451.2014.968088. Epub 2014 Nov 24. PubMed PMID: 25420023.
 106. Yoshino S, Nakagami G, Ohira T, Kawasaki R, Shimura M, Iwatsuki K, Sanada H, Kobayashi-Hattori K, Oishi Y, Yamane T. Hydrocellular foam dressing increases the leptin level in wound fluid. *Wound Repair Regen.* 2015 Sep; 23(5): 703-10. doi: 10.1111/wrr.12349. Epub 2015 Sep 14. PubMed PMID: 26242831.
 107. Postigo Mota S, Muñoz Bermejo L, Pinto Montealegre E, López Herranz M, Píriz Campos RM, López Corral JC. [Hydrocellular dressings or polyurethane foams]. *Rev Enferm.* 2011 Feb; 34(2): 66-7; quiz 68. Spanish. PubMed PMID: 21495394.
 108. Aderibigbe B A, Buyana B. Alginate in Wound Dressings. *Pharmaceutics.* 2018 Apr 2; 10(2). pii: E42. doi: 10.3390/pharmaceutics10020042. Review. PubMed PMID: 29614804; PubMed Central PMCID: PMC6027439.
 109. Lee K Y, Mooney D J. Alginate: properties and biomedical applications. *Prog Polym Sci.* 2012 Jan; 37(1): 106-126. PubMed PMID: 22125349; PubMed Central PMCID: PMC3223967.

110. Guo S, Dipietro LA. Factors affecting wound healing. *J Dent Res*. 2010 Mar; 89(3): 219-29. doi: 10.1177/0022034509359125. Epub 2010 Feb 5. Review. PubMed PMID: 20139336; PubMed Central PMCID: PMC2903966.
111. Karahan A, AAbbasoğlu A, Işık S A, Çevik B, Saltan Ç, Elbaş N Ö, Yalılı A. Factors Affecting Wound Healing in Individuals With Pressure Ulcers: A Retrospective Study. *Ostomy Wound Manage*. 2018 Feb; 64(2): 32-39. PubMed PMID: 29481325.
112. Avishai E, Yeghiazaryan K, Golubnitschaja O. Impaired wound healing: facts and hypotheses for multi-professional considerations in predictive, preventive and personalised medicine. *EPMA J*. 2017 Mar 3; 8(1): 23-33. doi: 10.1007/s13167-017-0081-y. eCollection 2017 Mar. Review. PubMed PMID: 28620441; PubMed Central PMCID: PMC5471802.
113. Thomas Hess C. Checklist for factors affecting wound healing. *Adv Skin Wound Care*. 2011 Apr; 24(4): 192. doi: 10.1097/01.ASW.0000396300.04173.ec. PubMed PMID: 21422844.
114. Gonzalez A C, Costa T F, Andrade Z A, Medrado A R. Wound healing - A literature review. *An Bras Dermatol*. 2016 Sep-Oct; 91(5): 614-620. doi: 10.1590/abd1806-4841.20164741. Review. PubMed PMID: 27828635; PubMed Central PMCID: PMC5087220.
115. Bischof Vukušć S, Flinčec Grgac S, Budimir A, Kalenić S. Cotton textiles modified with citric acid as efficient anti-bacterial agent for prevention of nosocomial infections. *Croat Med J*. 2011 Feb; 52(1): 68-75. PubMed PMID: 21328723; PubMed Central PMCID: PMC3046493.
116. Detailed feasibility analysis of gauze, bandages and absorbent cotton. Available From: <https://www.moea.gov.bt/wp-content/uploads/2017/10/hospital-PART-3.pdf>
117. Edwards J V, Yager D R, Cohen I K, Diegelmann R F, Montante S, Bertoniere N, Bopp A F. Modified cotton gauze dressings that selectively absorb neutrophil elastase activity in solution. *Wound Repair Regen*. 2001 Jan-Feb; 9(1): 50-8. PubMed PMID: 11350640.
118. Li T T, Lou C W, Chen A P, Lee M C, Ho T F, Chen Y S, Lin J H. Highly Absorbent Antibacterial Hemostatic Dressing for Healing Severe Hemorrhagic Wounds. *Materials (Basel)*. 2016 Sep 21; 9(9). pii: E793. doi: 10.3390/ma9090793. PubMed PMID: 28773912; PubMed Central PMCID: PMC5457081.
119. *Pharmaceutical Science in Homoeopathy and Pharmacodynamics* by K. P. Mazumdar, published by B. Jain Publishers, Jun 6, 2015, pp 71.
120. 2012 Emergency Response Guidebook: A Guidebook for First Responders during the Initial Phase of a Dangerous Goods/hazardous Materials Transportation Incident, published by United States Pipeline and Hazardous Materials Safety Administration, 2012.
121. *Regenerated Cellulose Fibres* by C Woodings, published by Elsevier, Apr 30, 2001.
122. Zaman H U, Islam J M, Khan M A, Khan R A. Physico-mechanical properties of wound dressing material and its biomedical application. *J Mech Behav Biomed Mater*. 2011 Oct; 4(7): 1369-75. doi: 10.1016/j.jmbbm.2011.05.007. Epub 2011 May 11. PubMed PMID: 21783147.
123. Repository Home. Investigation of surgical dressing in Keyatta National Hospital. Available From: <http://erepository.uonbi.ac.ke/handle/11295/77101>
124. Thomas S, Loveless P, Hay N P, Toyick N. Comparing non-woven, filmateci and woven gauze swabs. *J Wound Care*. 1993 Jan 2; 2(1): 35-41. PubMed PMID: 27911576.
125. Filmed Swabs – Sterile. Available from: <https://www.365healthcare.com/Woundcare/Swabs/Filmed-Swabs-Sterile~c~BD10>
126. Patel M, Bhrambhatt D. Nonwoven Technology for Unconventional fabrics. Available From: <https://pdfs.semanticscholar.org/e2f8/d32c2b1c05a3be7ed511197c7ad6b87fd460.pdf>
127. TRADITIONAL WOUNDCARE. Unsurpassed quality in a complete line of products. Available From: <https://docplayer.net/44575738-Traditional-wound-care-unsurpassed-quality-in-a-complete-line-of-products.html>
128. Surgical Supplies. Available From: <https://manualzz.com/doc/11065832/surgical-supplies>
129. Whats the Difference? Gauze Pads vs. Gauze Sponges. *Web Dukal* October 12, 2018.
130. Morgan N. Medical gauze 101. Available From: <https://woundcareadvisor.com/medical-gauze-101-vol4-no1/>
131. Patient Recovery Wound Care Catalog - Cardinal Health. Available From: <https://www.cardinalhealth.com/content/.../patient-recovery-wound-care-catalog.pdf>
132. Kumar GVS, Ramani S, Mahajan A, Jain N, Sequeira R, Thakur M. Imaging of retained surgical items: A pictorial review including new innovations. *Indian J Radiol Imaging*. 2017 Jul-Sep; 27(3): 354-361. doi: 10.4103/ijri.IJRI_31_17. PubMed PMID: 29089689; PubMed Central PMCID: PMC5644334.
133. Dynarex Sterile X-Ray Detectable Gauze Sponge 4" x 4" x 16 ply. Web Danlee Medical Products, Inc. Available From: <https://www.danleemedical.com/Dynarex-Sterile-X-Ray-Detectable-Gauze-Sponge-4-x-4-x-16-ply-item-3351-item-GA3351-P3579.aspx>
134. Kendall Vistec™ X-Ray Detectable Specialty Sponge 4" x 4" Sterile, 16-ply, Banded 10's. Available From: [Year 2019](https://www.edgepark.com/wound-care/bandages-and-dressings/absorptive-dressings/kendall-vistec-x-ray-detectable-specialty-sponge-4andquot%3B-x-</p>
</div>
<div data-bbox=)

135. X-Ray Detectable Gauze Sponges. Web Darby Dental Supply LLC.
136. RAY-TEC X-Rayable Sponge, 1944. Available From: <https://ourstory.jnj.com/ray-tec-x-rayable-sponge>
137. Abdel-Rahman R M, Abdel-Mohsen A M, Hrdina R, Burgert L, Fohlerova Z, Pavliňák D, Sayed O N, Jancar J. Wound dressing based on chitosan/hyaluronan/nonwoven fabrics: Preparation, characterization and medical applications. *Int J Biol Macromol.* 2016 Aug; 89: 725-36. doi: 10.1016/j.ijbiomac.2016.04.087. Epub 2016 May 2. PubMed PMID: 27151671.
138. Abou-Okeil A, Sheta A M, Amr A, Ali M A. Wound dressing based on nonwoven viscose fabrics. *Carbohydr Polym.* 2012 Sep 1; 90(1): 658-66. doi: 10.1016/j.carbpol.2012.05.093. Epub 2012 Jun 1. PubMed PMID: 24751090.
139. Jin NZ, Gopinath SCB. Potential blood clotting factors and anticoagulants. *Biomed Pharmacother.* 2016 Dec; 84: 356-365. doi: 10.1016/j.biopha.2016.09.057. Epub 2016 Sep 23. Review. PubMed PMID: 27668535.
140. Harkins A L, Duri S, Kloth L C, Tran C D. Chitosan-cellulose composite for wound dressing material. Part 2. Antimicrobial activity, blood absorption ability, and biocompatibility. *J Biomed Mater Res B Appl Biomater.* 2014 Aug; 102(6): 1199-206. doi: 10.1002/jbm.b.33103. Epub 2014 Jan 10. PubMed PMID: 24407857; PubMed Central PMCID: PMC4082766.
141. Sirkin M R, Cook P, Davis K G. Evaluating Alternatives to Traditional Cotton Laparotomy Sponges for Blood Absorption in the Austere and Mobile Surgical Environment. *J Spec Oper Med.* 2015 Winter; 15(4): 54-8. PubMed PMID: 26630095.
142. Eye Pads. Available From: <https://www.healthproductsforyou.com/c-eye-pads.html>
143. Eye Pad. Available From: <https://www.eurofarm-spa.com/en/products/hospital-line/eye-dressing/316-eye-pad.html>
144. Stevens S. Cleaning and dressing the eye after surgery. *Community Eye Health.* 2016; 29(94): 36. PubMed PMID: 27833264; PubMed Central PMCID: PMC5100474.
145. AMD-Ritmed Eye Pad Sponge Dressings Sterile. Available From: <https://www.usamedicalsurgical.com/eye-pad-sponge-dressings-by-amd-ritmed/>
146. Conforming Bandage. Available From: <https://www.elastoplast.com.au/products/muscle-joint-and-back-pain/conforming-bandages>
147. Mane V, Karthik D. An overview of crepe bandages. *Web Textile Today.* Available From: <https://www.textiletoday.com.bd/an-overview-of-crepe-bandages/>
148. A Guide to Canine and Feline Orthopaedic Surgery by Hamish Denny, Steve Butterworth, published by John Wiley & Sons, Apr 30, 2008.
149. Urinary Incontinence Products for Men. Available From: <https://www.webmd.com/men/guide/urinary-incontinence-products-for-men#1>
150. Francis K, Pang S M, Cohen B, Salter H, Homel P. Disposable Versus Reusable Absorbent Underpads for Prevention of Hospital-Acquired Incontinence-Associated Dermatitis and Pressure Injuries. *J Wound Ostomy Continence Nurs.* 2017 Jul/Aug; 44(4): 374-379. doi: 10.1097/WON.00000000000000337. PubMed PMID: 28549048.
151. Tranquility Peach Sheet Underpads - Heavy Absorbency, 21.5" x 32.5". Product Description. Available From: <https://www.newleafhomemedical.com/tranquility-peach-sheet-underpads-heavy-absorbency-21-5-x-32-5/>
152. Medline - Cotton Tipped Applicators. Available From: <https://bio-medical.com/coviden-curity-cotton-tipped-applicator-qtip.html>
153. Conforming Bandages. What Are They? Web Vitality Medical. Available From: <https://www.vitalitymedical.com/conforming-bandages.html>
154. Chapter 7. Perioperative Nursing. Lippincott Manual of Nursing Practice Eleventh, North American Edition by Sandra M Nettina MSN APRN BC ANP, published by LWW; Eleventh, North American edition (November 2, 2018)
155. Brouhard R. When to Use a Compression Bandage. Medically reviewed by Richard N. Fogoros, MD. Web [verywellhealth.com](http://www.verywellhealth.com), February 08, 2019. Available From: <https://www.verywellhealth.com/elastic-bandages-1298333>
156. SMTL Dressings Datacard. Tensopress. Setopress Data search. Available From: <http://www.dressings.org/Dressings/tensopre.html>
157. Dressings and Bandages. Available From: <https://www.stjohn.org.nz/First-Aid/First-Aid-Library/Immediate-First-Aid1/Dressings-and-Bandages/>
158. Vuković D. 15 Uses for a Triangular Bandage. Available From: <https://www.primalsurvivor.net/triangular-bandage-uses/>
159. Orthopaedic Bandage. Available From: <http://endomed.co.za/product/orthopaedic-bandage/>
160. 3M™ Scotchcast™ Soft Cast Casting Tape. Available From: https://www.3m.com/3M/en_US/company-us/all-3m-products/~/3M-Scotchcast-Soft-Cast-Casting-Tape/?N=5002385+3293316389&rt=rud
161. Fousekis K, Billis E, Matzaroglou C, Mylonas K, Koutsojannis C, Tsepis E. Elastic Bandaging for Orthopedic- and Sports-Injury Prevention and Rehabilitation: A Systematic Review. *J Sport Rehabil.* 2017 May; 26(3): 269-278. doi:

- 10.1123/jsr.2015-0126. Epub 2016 Aug 24. Review. PubMed PMID: 27632889.
162. Brock T M, Sprowson A P, Muller S, Reed M R. STICKS study - Short-sTretch Inelastic Compression bandage in Knee Swelling following total knee arthroplasty - a feasibility study. *Trials*. 2017 Jan 9; 18(1): 6. doi: 10.1186/s13063-016-1767-5. PubMed PMID: 28069060; PubMed Central PMCID: PMC5223465.
163. Antabak A, Barišić B, Andabak M, Bradić L, Brajcinović M, Haramina T, Haluzan D, Fuchs N, Durković S, Curković S, Luetić T, Sisko J, Prlić I. [Physical Properties of Plaster Bandages]. *Lijec Vjesn*. 2015 Nov-Dec; 137 (11-12):372-6. Croatian. PubMed PMID: 26975068.
164. Plain stockinette / Tubular bandage. Available From: <https://www.lohmann-rauscher.com/en/products/plaster-room/skin-protection/plain-stockinette-tubular-bandage/>
165. Tubular Stockinette. Web Bandage Plus. Available From: <https://www.bandagesplus.com/bandaging-supplies/bandages/tubular-stockinette#>
166. The Complete Technical Guide for Adhesive Tape. Web ecotape.com. Available From: <https://www.ecotape.com/blog/complete-technical-guide-adhesive-tape/>
167. Tape An Excursion Through The World Of Adhesive Taps. <https://www.atera.com/uploads/Bestanden/Afera%2050th%20Anniversary%20Publication.pdf>
168. Zajackowski M j. Acrylate-Polyether Based Pressure Sensitive Adhesives. Available From: <https://www.pstc.org/files/public/Zajackowski07.pdf>
169. Rubber-Based Adhesives. Web Adhesives Research, Inc. Available From: <https://www.adhesivesresearch.com/technologies/chemistries/rubber-based-adhesives/>
170. Pressure-Sensitive Adhesives and Applications 2nd Edition by Istvan Benedek, published by CRC Press, Published February 3, 2004.
171. Adhesives Technology Handbook 2nd Edition by Sina Ebnesajjad Arthur H. Landrock Sina Ebnesajjad, published by Elsevier 7th August 2008.
172. Darmas B. Use of barrier products in the prevention of adhesion formation following surgery. *J Wound Care*. 2008 Sep; 17(9): 405-8, 411. Review. PubMed PMID: 18833900.
173. González-Quintero V H, Cruz-Pachano F E. Preventing adhesions in obstetric and gynecologic surgical procedures. *Rev Obstet Gynecol*. 2009 Winter; 2(1): 38-45. PubMed PMID: 19399293; PubMed Central PMCID: PMC2672996.
174. Peralta E. Overview of topical hemostatic agents and tissue adhesives. Available From: <https://www.uptodate.com/contents/overview-of-topical-hemostatic-agents-and-tissue-adhesives>
175. Khoshmohabat H, Paydar S, Kazemi H M, Dalfardi B. Overview of Agents Used for Emergency Hemostasis. *Trauma Mon*. 2016 Feb 6; 1(1):e26023. doi: 10.5812/traumamon.26023. eCollection 2016 Feb. Review. PubMed PMID: 27218055; PubMed Central PMCID: PMC4869418.
176. Chiara O, Cimbanassi S, Bellanova G, Chiarugi M, Mingoli A, Olivero G, Ribaldi S, Tugnoli G, Basilicò S, Bindi F, Briani L, Renzi F, Chirletti P, Di Grezia G, Martino A, Marzaioli R, Noschese G, Portolani N, Ruscelli P, Zago M, Sgardello S, Stagnitti F, Miniello S. A systematic review on the use of topical hemostats in trauma and emergency surgery. *BMC Surg*. 2018 Aug 29; 18(1): 68. doi: 10.1186/s12893-018-0398-z. Review. PubMed PMID: 30157821; PubMed Central PMCID: PMC6116382.
177. Barroso J, Osborne B, Teramura G, Pellham E, Fitzpatrick M, Biehl R, Yu A, Pehta J, Slichter S J. Safety evaluation of a lyophilized platelet-derived hemostatic product. *Transfusion*. 2018 Dec; 58(12): 2969-2977. doi: 10.1111/trf.14972. Epub 2018 Nov 18. PubMed PMID: 30450601; PubMed Central PMCID: PMC6301028.
178. El Hassan M, Elnour A A, Farah F H, Shehab A, Al Kalbani N M, Asim S, Shehab O A, Abdulla R. Clinical pharmacists' review of surgical antimicrobial prophylaxis in a tertiary hospital in Abu Dhabi. *Int J Clin Pharm*. 2015 Feb; 37(1): 18-22. doi: 10.1007/s11096-014-0045-4. Epub 2014 Dec 9. PubMed PMID: 25488316.
179. Weber A, Schneider C, Grill E, Strobl R, Vetter-Kerkhoff C, Jauch KW. [Interventions by clinical pharmacists on surgical wards - impact on antibiotic therapy]. *Zentralbl Chir*. 2011 Feb; 136(1): 66-73. doi: 10.1055/s-0030-1247469. Epub 2011 Feb 18. German. PubMed PMID: 21337293.
180. Martin C M, Hawkins L. Wound care basics for the pharmacist. *Consult Pharm*. 2013 Jun; 28(6): 344-6, 348-52. doi: 10.4140/TCP.n.2013.344. PubMed PMID: 23748122.
181. Watret L, White R. Surgical wound management: the role of dressings. *Nurs Stand*. 2001 Jul 18-24; 15(44): 59-62, 64, 66. Review. PubMed PMID: 12212002.

