



GLOBAL JOURNAL OF MEDICAL RESEARCH: E
GYNECOLOGY AND OBSTETRICS
Volume 19 Issue 4 Version 1.0 Year 2019
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Mozart Compilation during Pregnancy Gave Higher Number of Neurons of *Rattus Norvegicus* Offsprings' Cerebrum Compared with Jazz, Blues, and Rock Compilations

By Syania M O, Hermanto T J & Mudjiani B

Abstract- Aim: To analyze the difference of the number of neurons in the cerebrum of *Rattus norvegicus* offsprings exposed to Mozart, Jazz, Blues, and Rock compilations during pregnancy.

Method: Experimental study with single-blind randomized post-test only control group design using *Rattus norvegicus* as animal subjects that were divided into four groups: Mozart, Jazz, Blues, and Rock groups. The exposures started from the 10 day of gestation for 1 hour in a dark atmosphere with an intensity of 60 dB, in a distance of 25 cm from the exposure box, during March-May 2019. After cesarean delivery at day 19, the offsprings brain were prepared and stained by Hematoxylin-Eosin, then analyzed in 5 fields in each hemisphere with 1000x magnifying microscope. The study was conducted after ethical clearance and used a comparison statistical test chosen accordingly.

Result: There were significant differences of the number of neurons between Mozart ($71,96 \pm 14,44$) with Jazz ($41,06 \pm 7,65$), Blues ($40,92 \pm 6,36$), and Rock group ($31,31 \pm 8,19$) with $p=0,000$.

Conclusion: The number of neurons in the cerebrum of *Rattus norvegicus* offsprings exposed to Mozart compilation during pregnancy proved to be higher than Jazz, Blues, and Rock compilations.

Keywords: *rattus norvegicus*, pregnant, mozart, jazz, blues, rock, offsprings, cerebrum, neuron.

GJMR-E Classification: NLMC Code: WQ 240



Strictly as per the compliance and regulations of:



Mozart Compilation during Pregnancy Gave Higher Number of Neurons of *Rattus Norvegicus* Offsprings' Cerebrum Compared with Jazz, Blues, and Rock Compilations

Syania M O ^α, Hermanto T J ^σ & Mudjjani B ^ρ

Abstract- Aim: To analyze the difference of the number of neurons in the cerebrum of *Rattus norvegicus* offsprings exposed to Mozart, Jazz, Blues, and Rock compilations during pregnancy.

Method: Experimental study with single-blind randomized post-test only control group design using *Rattus norvegicus* as animal subjects that were divided into four groups: Mozart, Jazz, Blues, and Rock groups. The exposures started from the 10 day of gestation for 1 hour in a dark atmosphere with an intensity of 60 dB, in a distance of 25 cm from the exposure box, during March-May 2019. After cesarean delivery at day 19, the offsprings brain were prepared and stained by Hematoxylin-Eosin, then analyzed in 5 fields in each hemisphere with 1000x magnifying microscope. The study was conducted after ethical clearance and used a comparison statistical test chosen accordingly.

Result: There were significant differences of the number of neurons between Mozart (71,96 ± 14,44) with Jazz (41,06 ± 7,65), Blues (40,92 ± 6,36), and Rock group (31,31 ± 8,19) with p=0,000.

Conclusion: The number of neurons in the cerebrum of *Rattus norvegicus* offsprings exposed to Mozart compilation during pregnancy proved to be higher than Jazz, Blues, and Rock compilations.

Keywords: *rattus norvegicus*, pregnant, mozart, jazz, blues, rock, offsprings, cerebrum, neuron.

I. INTRODUCTION

At least there are seven phenomenon/awareness that lead to this effort, i.e. modify the structure and function of the fetal brain to have more neurons, more glia cells and more dendritic density as follow 1. The need for the next better generation, especially with the better brain and its breakthrough¹⁻¹⁴ 2. The findings from Gardner on multiple intelligences¹⁵ 3. The revelation that brain growth and development start in the womb¹⁶⁻²⁰ 4. The environment enrichment by M. Diamond, including analysis of the Einstein brain²¹⁻²³ 5. The findings of Tomatis: brain growth and development needs sound and music²⁴⁻²⁶ 6. Rauscher and Shaw,

findings²⁷⁻³¹ 7. The FOAD and DOHAD hypothesis³²⁻³³. There were already series of studies in Surabaya that analyzed this effort in what kind of (musical) stimulation, the orders, duration, gestational age, what kind of nutrition – with the dependent variables: BDNF, numbers of neurons, glia cells, dendritic density and neuronal apoptotic index. This team also had analyzed the influence of frequency, color, intensity, beat of various compilations including traditional, jazz, rock, blues and pop music compilations and its influence on some plants³⁴⁻⁴⁸.

In this study, Mozart compilation during pregnancy were compared to Jazz, Blues and Rock compilations to more deeply understand the neurophysiology impact of musical exposure to the structure and the function of the fetal brain.

II. MATERIALS AND METHODS

This research was an experimental laboratory study with a single-blind randomized post-test only control group design. The sample size was calculated using the Federer formula: for each group was seven. The subjects were pregnant healthy *Rattus norvegicus*, weighing 130-180 grams and never giving birth. The subjects were divided into 4 groups randomly, which are the treatment groups that were exposed Mozart, Jazz, Blues, and Rock compilations. All exposure carried out in day 10 of pregnancy until delivery at day 19, for 1 hour in a dark atmosphere (represent the night atmosphere) with an intensity of 60 dB and at of 25 cm distance from the exposure box. At cesarean delivery, two offsprings with the heaviest weight were taken in each group, sacrificed by chloroform, weighed, and the brain were prepared and then stained by Hematoxylin-Eosin methode. The number of neuron cells calculated by 1000x magnifying microscope: 5 right and 5 left hemispheres. Statistical analysis was chosen accordingly. The study was conducted in the Animal Cages and Pathology Laboratory of the Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, during March-May 2019 after ethical clearance.

Author α: e-mail: syaniamegaactariyandra@yahoo.com)

Author σ: Dr., Sp. OG (K). e-mails: hos_hermanto@yahoo.com, hermanto.tri@fk.unair.ac.id

Author ρ: Dr. Sp. S. e-mail: mudjjani.basuki@fk.unair.ac.id



Fig. 1: The Exposure Box

III. RESULTS

a) Offsprings Birthweight

Table 1: *Rattus norvegicus* offsprings birthweight

Groups	N		Mean \pm SD
	Mother	Offspring	
X1	5	10	4,42 \pm 1,15
X2	7	14	4,78 \pm 0,42
X3	6	12	4,11 \pm 0,81
X4	7	14	4,62 \pm 0,51

Note: X1 (Mozart) X3 (Blues)
 X2 (Jazz) X4 (Rock)

b) The Number of Neurons in Cerebrum Cortex

The number of cerebrum neuron cells was known by a dark blue cell nucleus, then the preparation was counted by 1000x magnifying microscope 10 visual fields consisting of 5 right and 5 left hemispheres. There were 6 preparations that could not be counted because 4 cortex of Mozart group had meningitis and 2 cortex of Blues group were not found.

Table 2: The Number of Neurons in Cerebrum Cortex

Groups	N Preparation	Mean \pm SD
X1	10	71,96 \pm 14,44
X2	14	41,06 \pm 7,65
X3	12	40,92 \pm 6,36
X4	14	31,31 \pm 8,19

Note: X1 (Mozart) X3 (Blues)
 X2 (Jazz) X4 (Rock)

This table shows the average number of neurons of the *Rattus norvegicus* offsprings in cerebrum cortex. The highest mean value was in the Mozart group (X1) followed by Jazz, Blues, and Rock (X2, X3, X4) groups.

Visualization of the distribution of the number of neurons will give better impression that shows the highest number of Mozart group.




Fig. 2: Scatter Diagram of the Number of Neurons in the Cerebrum Cortex

Table 3: Post-Hoc Test of the Number of Neurons in the Cerebrum Cortex

Groups	P Value		
	X2	X3	X4
X1	0,000*	0,000*	0,000*
X2	-	0,938	0,000*
X3	-	-	0,000*

Note: X1 (Mozart) X3 (Blues)
 X2 (Jazz) X4 (Rock)

This table shows the results of the analysis with post-hoc LSD test: Mozart group with Jazz, Blues, and Rock groups shows the value of $p = 0,000$ ($p < 0.05$), which means that there were significant differences. Analysis of Jazz group with Blues group showed the value of $p = 0.938$ ($p < 0.05$) which means there was no significant difference. Analysis of Jazz group with Rock group shows the value of $p = 0,000$ ($p < 0.05$) which means that there was a significant difference. Furthermore, the analysis of the Blues group with the Rock group shows the value of $p = 0,000$ ($p < 0.05$) which means that there was also significant difference.

Table 4: Analysis of the Number of Neurons in the Right and Left Hemispheres

Groups	Mean \pm SD		P	
	Right	Left	Right	Left
X1	70,80 \pm 15,75	73,12 \pm 13,46	0,177	0,200
X2	39,34 \pm 7,37	42,77 \pm 7,64	0,200	0,200
X3	39,92 \pm 6,01	41,92 \pm 6,66	0,200	0,200
X4	31,46 \pm 6,78	31,17 \pm 9,49	0,200	0,101

Note: X1 (Mozart) X3 (Blues)
X2 (Jazz) X4 (Rock)

This table shows that the average number of neurons were not much differences between right and left hemispheres even the number a little bit higher in the left one.

IV. DISCUSSION

Developing countries with more than 100 orders in Human Development Index (HDI) should have a program to increase their levels, and education is one of the single solution of choice.⁴⁹ Education needs – best education program and brain capacity: multiple biopsychosocial potencies. These countries can only catch up higher HDI rank not by better education program which is very expensive but through a breakthrough program related to the brain structure and function.⁵⁰ We believe in “From Neurons to Nations” premise and the idea of environment enrichment which have been mentioned in the introduction.

Environment enrichment by combination of certain musical stimulation and nutrition during pregnancy have been studied here in the last twenty years. This program in line with the Harvard program which address the influence of early sensory exposure to the brain growth and development; but not with Brain Decade premise which address mostly the brain disease.¹⁷ We understand that medicine especially Obstetrics and Maternal Fetal Medicine deal with pregnancy, birth and puerperium period, also their diseases, not with fetal brain growth and development. Only in pediatrics there is subspecialties in Growth and Development.

Our standard operating procedure is default Mozart compilation which is exposed one hour in duration, at night 8-11 PM, start 20 weeks of pregnancy or day 10, 65 dB, 25 cm distance. This procedure has been used for almost twenty years with the same results in variables studied (BDNF, synaps, number of neurons, glia, dendritic density).

There have been many studies regarding the effect of music on brain development. For example, Kuhlmann, Mariana, Tomatis, Campbell, Kendrick and other research groups. They assume that different types and genres of music have different influences. Music with classical genres has a positive effect on listeners, while rock music has a negative effect on listeners.^{24,26,51-53} The statement is proven in studies with

plant and animal subjects. On the subject of plants, researcher Dorothy Retallack started the determination by using the control variables of light, temperature, and air it turns out some types of plants with rock music exposure for four weeks stopped growing and damaged.⁵⁴ While research with animal subjects was conducted by researchers Harvey Bird and Farleigh Dickinson. The two experts conducted their research on mice as experimental animals given exposure to Strauss music from Strauss, voodoo drum rhythms, and then noiselessly measured their ability to complete the game 'maze'. The results showed that the rats given drum exposure had difficulty completing the game. Other evidence was carried out by Hermanto et al., with the title "The Influence of 11 Mozart Compositions during Pregnancy to The Perinatal Outcome and BDNF Umbilical Cord Blood". The purpose of this study was to compare BDNF levels in infants exposed to Mozart music while in the womb and without exposure. It turned out that there were significant differences at the BDNF level in the two groups.⁹

Based on several studies conducted by experts, shows that the pre-natal period is the most amazing period in the fetal development phase. This phase does not only depend on genetics, but environmental factors also play an important role in the development of the functional capacity of the body's organs. So it can be said that during pregnancy is the right time in preparing the potential for fetal intelligence early on. The studies of Brent Logan, Rene van de Carr and Beatriz Manrique show the existence of environmental interference on fetal intelligence. They discovered the premise of "stimulation induced morphological changes" which means that the structure of the brain is formed by external stimuli.⁵⁶⁻⁵⁸

Music exposure given during prenatal has an influence on the process of proliferation, migration, differentiation, myelination, synaptogenesis, and apoptosis of brain cells. The human brain is one of the most complex organ systems. Cellularly, the nervous system consists of two types of cells, namely neuron and glia cells. These two cells work in harmony so that the brain's commutation ability goes well. Fetal neuron cells stop proliferating until 32 months of gestation, while glia cells can proliferate until post-natal. The formation of these two cells can be stimulated through exposure to music during pregnancy. The more the number of neuron cells formed, it is hoped that intelligence will increase.^{4,12}

This research is a continuation of a series studies with the same aim in preparing a smarter generation from the womb. This research was conducted to find out the differences in the number of neurons in the brain of new born *Rattus norvegicus* in the cerebrum between those who were exposed to Mozart, Jazz, Blues, and Rock compilations during pregnancy.

In this study we compared musical exposure of 4 kinds of western music compilations, we also compared with traditional Indonesian music and religious music compilations in other studies.

The mechanism/s how the music affect/s the brain are not completely understood. It is correlated with BDNF, different in frequency, color, beat, timing, and duration have detected probably explain in parts the mechanism.

We can find that differences in analyzing the frequency and major minor proportion.




Fig. 3: The Different Frequency of Mozart, Jazz, Blues and Rock Compilations. Mozart Shows the Lowest Frequencies

This research was carried out since the 10th day pregnancy of *Rattus norvegicus* due to ear formation in *Rattus norvegicus* was complete at 9-10 days of gestation. Ernawati in her study said that there was no significant difference in the apoptotic index of Mozart's music exposure at the beginning of pregnancy compared to 10 days of gestation. This is possible because the stimulus in the form of sound is received through the ear, then these mechanical waves are converted into electrical pulses and transmitted to the auditory cortex through the auditory nerve when the ear is fully formed. So that the provision of music stimuli will begin to affect after the ear is formed and functioning and begin to form synapses in the 20-24th weeks of pregnancy or equivalent to the 10th day of pregnancy in mice.⁵⁹

Music stimulation during pregnancy is reported to improve fetal brain development, increase spatial-temporal abilities in newborn mice, and trigger rapid progress in motor abilities such as sitting and walking in infants. Research conducted by Kim et al., found that prenatal noise exposure resulted in stunted growth, decreased neurogenesis in the hippocampus, and

disruption of spatial abilities in newborn rats. In contrast, prenatal music exposure can improve neurogenesis in the hippocampus and spatial ability in newborn mice.⁶⁰⁻⁶¹

Brains that grow in a stimulus-rich environment have thicker cortex, larger nucleus neuron cells and more glia cells. Brain neurons that grow in there have more dendritic sites, allowing more synapses to form. Rees also stated that the number of cells undergoing apoptosis depends on synapses, the more synapses the less apoptosis occurs. The richer the neuron cells with more dendritic sites, the more synapses are formed so that the number of cells undergoing apoptosis will also be reduced. Brains that grow in a stimulus-rich environment will experience less apoptosis, thereby increasing brain capacity.²⁰ Ismudi in his study compared 3 Mozart music compilations consisting of several random song sequences and analyzed using computer software "Cool Edit Pro 2.0". As a result of the three compilations, compilation 1 has a high frequency tone, a wider amount of energy area, and regular intensity. Ismudi believes that these characteristics are an effective composition for stimulating neurons. While Mozart 2 and 3 compilation tends to have irregular fluctuations in intensity, so that it will produce an amount of energy that changes in every seconds. In that study showed that exposure to Mozart 1 music compilation during rat pregnancy produced the lowest apoptotic index in neuronal cells of newborn mice compared to Mozart 2 compilation, compilation 3, and control. For this reason, compilation of Mozart 1 according to Ismudi is a standard Mozart compilation which forms the basis of research and application of Mozart's music exposure during pregnancy as a prenatal stimulation.³⁷

The results of our study in the four groups showed the average number of neurons in the cerebrum in the Mozart group compared to the Jazz, Blues, and Rock groups. The average number of neurons of Mozart group is higher among others. Meanwhile, the average number of neurons in the Jazz group is higher than Blues, and Rock. And the average number of neurons in the Blues group is higher than Rock. In the cerebrum, the mean number of neurons in Mozart was 71.96 ± 14.44 , in Jazz 41.06 ± 7.65 , in Blues 40.92 ± 6.36 , and in Rock 31.31 ± 8.19 .

We also counted the average number of neurons in right and left hemisphere. It was done by looking at 10 fields of view (consisting 5 fields each hemisphere). The highest average number of neurons was found in the left hemisphere compared to the right hemisphere in the Mozart, Jazz, and Blues Group. Otherwise, in the Rock group the average number of neurons in the right hemisphere was higher compared to the left hemisphere. Based on the theory, right and left brain have different functions and communicate each other through a band of nerves. Left-brain has specialization in language and logic, meanwhile right-

brain has specialization in creativity and intuition. Commonly in adults, we believed that music is processed in the right-brain. One of the right-brain function is for controlling the ability to play instrument with ease, recognize a song from melody, and play it back upon "hearing it". In this study, we know that it is different if we look from the number of neurons in each hemisphere of *Rattus norvegicus* offsprings.^{62,63} The probable cause is still unknown.

In the analysis of variance using the ANOVA test, there was a significant difference in the data of the number of neurons in the cerebrum between all groups ($p < 0.05$). That means there are significant differences in the number of neuron cells in the cerebrum of *Rattus norvegicus* offsprings.

Furthermore, Post-Hoc LSD (Least Significant Difference) analysis was performed to determine all possible differences in the comparison of each group. If $p < 0.05$, there was a significant difference. Table 3 shows the results of the Post-Hoc LSD test. The comparison of each group shows that there were significant differences with the p value < 0.05 , between the Mozart group with the Jazz, Blues, and Rock groups with a value of $p = 0.000$, $p = 0.000$, and $p = 0.000$ ($p < 0.05$). Analysis was also carried out between groups, the results obtained from the analysis of the Jazz with the Blues groups showed a value of $p = 0.938$ ($p < 0.05$) which means no significant difference between the number of neurons in the Jazz and Blues groups. Analysis of the Jazz with the Rock groups showed the value of $p = 0.000$ which means that there was a significant difference between the number of neurons in the Jazz and Rock groups. Furthermore, the analysis of the Blues group with the Rock groups showed the value of $p = 0.000$ ($p < 0.05$) which means that there was a significant difference between the number of neurons in the Blues and Rock groups.

From this study it can be concluded that Mozart exposure gives the highest number of neurons compared to other western music exposure which are Jazz, Blues, and Rock with significant differences. In addition, Jazz exposure gave a higher number of neurons compared to Blues and Rock with a significant difference from Rock, but didn't make a significant difference from Blues. Exposure to Blues compilation gave a higher number of neurons than Rock with significant differences. This might be caused by every compilation (music) has multidimensional properties and attributes of perception that affect the apoptosis index of brain cells. Other studies conducted by Kauser et al., and Sanyal found that giving a prenatal stimulus with music that has regular rhythm can improve postnatal spatial ability and also memory function, but giving music with complex rhythms and arithmetic noises has the opposite effect.⁶⁴⁻⁶⁷

Study conducted by Sanyal et al., about the effect of music and noise on changes in the number of

neuron and glia cells in several brain areas of newborn chicks. The results showed an increase in the number of neuron cells in the brain area studied due to music stimulation. Music stimulation will have an effect on increasing neurogenesis or decreasing brain cell death.⁶⁸

If traced from several literature reviews, Mozart has a frequency of around 8000 Hz, different from Jazz, Blues, and Rock which has a frequency of ≈ 15000 Hz. Campbell suspected that Mozart's stimulation in general can affect neurochemical changes in which the clarity, majesty, rhythm, melody and high frequency which are able to stimulate the creative and motivational areas of the brain so as to calm its listeners, improve concentration, memory and spatial perception. The results of the Borner et al., 2000 study revealed that Mozart K488 increased the activity of the dorsolateral prefrontal cortex, occipital cortex and cerebellum compared to piano music of the 1990s and Beethoven. The same thing is supported by Ningsih in her research which shows Mozart is better than the music of Beethoven and Chopin. This opinion was proven from the calculation of the number of neurons in the cerebrum and cerebellum of *Rattus norvegicus* offsprings exposed by Mozart's music during pregnancy proved to be higher than those exposed to the music of Beethoven, Chopin, and not exposed to music and there were significant differences.^{26,46}

Research conducted by Kirchberger and Russo comparing the dynamic range of various music stated that Classical and Jazz music has a wider dynamic range than Rock, Rap, Schlager, and Pop music. Dynamic range is the ratio between the lowest and highest volume of an instrument. Narrow dynamic range can make hearing fatigue for the listener.⁶⁹

We recognize that the slow tempo can make an individual feel calmer and lighter. Classical music with a tempo of 60x / minute has been proven to reduce anxiety levels and also improve memory. While Jazz music is estimated to have a tempo of about 240x / minute.⁶⁵ Based on How Music Affects Us and Promotes Health, Mozart's Classical Music has the ability to activate brain areas in processing information.⁷⁰

Research conducted by Poikonen et al., comparing Event Related Brain Potential measured through EEG produced by stimulation of Jazz and Rock music, it was found that Jazz music activates the brain more strongly than Rock music. Whereas Jazz music with Blues music has almost the same characteristics in terms of frequency. The differences also come from the characteristic of music, Jazz music has a distinctive character that is blue notes, improvisation, polyrhythms, syncopation, and shuffle notes, while Blues music tends to represent feelings of sadness, somber, and depressed.⁷¹

This shows that Mozart proved to be better than western music such as Jazz, Blues, and Rock music

according to the results of our research supported by the theory of Hermanto that the brain needs energy and the best energy is sound. The type of season that can help increase the number of brain cells is classical music from Mozart with a frequency of 5000-8000 Hz where the frequency is not too high so it is considered suitable for the fetal environment during pregnancy.⁸⁻⁹

V. CONCLUSION

The number of neurons in the cerebrum of *Rattus norvegicus* offsprings in the Mozart exposure was higher than the Jazz, Blues, and Rock with significant differences.

VI. RESEARCH EXCELLENCE

This research is the first study conducted by comparing the differences in the number of neurons to four types of Western music, namely Mozart, Jazz, Blues, and Rock compilation.

VII. RESEARCH NOVELTY

This research was conducted by looking at 10 fields of view of *Rattus norvegicus* offsprings neuron cell preparations, including in 5 fields of the right and 5 fields of the left hemispheres of the *Rattus norvegicus* cerebrum. So it can be known which part of the brain is thought to have a higher influence in thought processes and intelligence.

REFERENCES RÉFÉRENCES REFERENCIAS

- Hermanto T J. (2002). New Role of Obstetricians: Developing Smarter Fetus in the Womb. Malang: POGI Annual Meeting.
- Hermanto T J. (2002). Prenatal University. Indonesiatrek: Toward a New Brighter Generations. Surabaya: Symposium on Medical Education Innovation.
- Hermanto T J, Esteopangesti ATS., and Widjiati, (2002). The influence of various musical exposure to pregnant (*Rattus norvegicus*) Rat to the number of offsprings rat brain cells. Surabaya: 3rd Scientific meeting on Maternal-Fetal Medicine and AOFOG Accredited Ultrasound Workshop.
- Hermanto, T J. (2004). Smart babies through Prenatal University. Mission: Impossible?', M O G I, 28(1), 1-14.
- Hermanto T J. (2010). Antenatal Care in the Third Milenium. Jakarta: POGI Annual Meeting (replaces Padang).
- Hermanto T J. (2011). Neurostimulation for Fetus in the Womb to Prepare the Next Smarter Generation. Surabaya: Seminar on IINS- From Neuron to Character Builidings.
- Hermanto T J. (2011). Nutrition and Stimulation in the Womb for the Next Better Generation. Jakarta
- Ibis Tamarin: Seminar by Centre for Intelligences MOH.
- Hermanto T J. (2012). Prostration in the Womb. Smarter Baby through Combination of Mozart compilations and Nutrition in the Womb. Surabaya: Global Persada Press.
- Yulia P P, Widjiati, Hermanto T J. (2018). Effect of prenatal Mozart composition on Brain Derived Neurotrophic Factor expression in cerebrum and cerebellum of *Rattus norvegicus* offspring from Food Restriction 50 model. M O G I 26(1): 6.
- Verny T. (1988). The Secret Life of the Unborn Child. New York: Dell Publishing.
- Verny T. (2002). Pre-Parenting. Nurturing Your Child from Conception. New York: Simon & Schuster.
- Story, L. (2003). A Head Start in Life? Prenatal Parenting and Discourse of Fetal Stimulation. Atlantis vol 27(2), 27-72.
- Logan B. Learning Before Birth: Every Child Deserves Giftedness. Author house.
- Hopson J L. (1998). Fetal Psychology. Psychology Today, October.
- Gardner H. (1999). Intelligence Reframed: Multiple Intelligences for the 21st Century. Cambridge: Basic Book.
- Shonkof J P. (2000). The Developing Brain – From Neurons to Neighborhoods. <http://www.ncbi.nlm.nih.gov/books/NBK225562>
- Centre on the Developing Child. (2007). The Science of Early Childhood Development. Closing The Gap Between What We Know and What We Do. Harvard University.
- Johnson M H, de Haan M. (2015). Developmental Cognitive Neuroscience. 4th ed. Malden: Wiley Blackwell – p 36.
- Stiles J, Jernigan T L. (2010). The Basics of Brain Development. Neuropsychol Rev Dec; 20(4): 327–348. Nov 3. doi: 10.1007/s11065-010-9148-4
- Rees S., & Walker D. (2001). Nervous and Neuromuscular Systems. In Harding R & Docking A D. (eds.), Fetal growth and Development (Vol. 1, pp. 154-185). Cambridge: Cambridge University Press.
- Diamond M. & Hopson J L. (1999). Magic Trees of the Mind: How to Nurture Your Child's Intelligence, Creativity, and Healthy Emotions from Birth through Adolescence. New York: Plume.
- Diamond M C. 2001. Response of the Brain to Enrichment. An. Acad. Bras. Cienc., 73 (2).
- Bures J, Buresova O, Krivanek J. (1988). Brain and Behaviour. Paradigms for Research in Neural Mechanisms. Chisester: John Wiley & Sons.
- Leeds J. (1997). Therapeutic Music and Sound in Health care: Tomatis@ Method, Frequency Medicine for 21st Century. American J of Acupunture 25(4): 299-305.

25. Levitin D. (2006). This Is Your Brain on Music. London: Atlantic Books.
26. Campbell D. The Mozart Effect. (1997). The Mozart Effect. Tapping the Power of Music to Heal the Body, Strengthen the Mind, and Unlock the Creative Spirit. New York: Avon Books.
27. Rauscher F H, Shaw G L, Ky K N. (1995). Listening To Mozart Enhances Spatial-Temporal Reasoning: Towards A Neurophysiological Basis. *Neurosci Lett*. Feb 6; 185(1): 44-7.
28. Rauscher F H, Shaw G L, Levine L J, Ky K N, Wright E L. (1994). Music and Spatial Task Performance: A causal relationship. Paper presented at American Psychology Association Annual Meeting Los Angeles.
29. Rauscher F H. (2006). The Mozart Effect in Rats: Response to Steele. *Music Perception*, 23, 447-453.
30. Bodner M L, Muftuler L T, Nalcioğlu O and Shaw G L. (2001). fMRI Study Relevant to the Mozart Effect: Brain Areas Involved in Spatial-Temporal Reasoning. *Neurol Res*; 23: 683-690.
31. Thomson B M, Andrews S R. (2000). An Historical Commentary on the Physiological Effects of Music: Tomatis, Mozart and Neuropsychology. *Integrative Physiological and Behavioral Science*, July-September, Vol. 35, No. 3, 174-188.
32. Barker DJP, Bergmann R., Ogra P L. (2007). The Window of Opportunity: Pre-Pregnancy to 24 Months of Age. Bali, Indonesia, April 1-5.
33. Gluckman P, & Hanson M. (2003). Mismatch. efile accessed 2011.
34. Jamil, Hermanto T J, Komang, Dyah. (2003). Sound Attenuation in Pregnant Sheep Measured By Intrauterine Microphone. Unpublished.
35. Didi D, Agus S, Hermanto T J. (2005). The Influence of Mozart's Twinkle-Twinkle Little Star (K 265) to the Fetal Biophysical Profile. Unpublished.
36. Indra P K, Agus S, Hermanto T J. (2005). Comparison of the Influence of Mozart's Twinkle-Twinkle Little Star (K 265) to the Fetal Biophysical Profile in the Day and the Night. Unpublished.
37. Ismudi H P, Widjati, Hermanto T J. (2007). Comparison of the Brain Neuronal Apoptotic Index of *Rattus norvegicus* Offsprings exposed to 3 Sequences of Mozarts compositions During Pregnancy. Unpublished.
38. Rizarina S, Hermanto T J, Widjati. (2005). Comparison of the Brain Neuronal Apoptotic Index of *Rattus norvegicus* Offsprings Exposed and Non Exposed to Sequence of Mozarts Composition During Pregnancy. Unpublished.
39. Niken W S, Margarita M., M. Dikman A, Hermanto T. J. (2009). Comparison of the Brain Derived Neurotrphic of *Rattus norvegicus* Offsprings Exposed and Non Exposed to Sequence of Mozart's Composition during Pregnancy. Unpublished.
40. Toni W, Hermanto T J. (2012). Comparison of the Newborn Umbilical Brain Derived Neurotrophic Factor Exposed and Nonexposed to Sequence of Mozarts Compositions during Pregnancy. Unpublished.
41. Rino A, Widjati, Hermanto T J. (2014). Comparison of the Brain Neuronal Apoptotic Index of *Rattus norvegicus* Offsprings exposed at 30, 60 and 120 minutes to Sequence of Mozarts Compositions during Pregnancy. Unpublished.
42. Harya N, Widjati, Hermanto T J. (2015). The Influence of Sequence of Mozarts Compositions during Pregnancy to Expression of Brain Derived Neurotrophic Factor and Protein kinase at Cerebrum and Cerebellum *Rattus norvegicus* Offsprings. Unpublished.
43. Nareswari ICM, Widjati, Hermanto T J. (2015). The Influence of Sequence of Mozarts Compositions during Pregnancy to Expression of Brain Derived Neurotrophic Factor and Dendritic Density at Cerebrum and Cerebellum *Rattus norvegicus* Offsprings. Unpublished.
44. Rozi A A, Widjati, Hermanto T J. (2015). The Influence of Sequence of Mozarts Compositions during Pregnancy to Expression of Brain Derived Neurotrophic Factor, Number of Neuron and Glia cells at Cerebrum and Cerebellum *Rattus norvegicus* Offsprings. Unpublished.
45. Eka N M., Widjati, Windhu P, Hermanto T J. (2017). Comparison of the Brain Neuronal Apoptotic Index of *Rattus norvegicus* Offsprings Exposed to Sequence of Mozarts Compositions during Pregnancy in the Light and Dark Conditions. *MOGI* vol 25 no 3.
46. Ningsi H A. (2018). Differences in the Effects of Music Exposure on Mozart, Beethoven, and Chopin during Pregnancy to the Number of Neuron Cells in Cereberum and Cerebellum. Thesis. Unpublished.
47. Habibie PH., Widjati, Hermanto TJ. (2018). Neuronal apoptotic index of *Rattus norvegicus* cerebrum and cerebellum exposed to reversed sequence of default Mozart compositions during pregnancy. Unpublished.
48. Angelucci F, Ricci E, Padua L, Sabino A, Tonali P A. (2007). Music exposure differentially alters the levels of brain-derived neurotrophic factor and nerve growth factor in the mouse hypothalamus. *Neuroscience Letters* 429: 152-155.
49. UNDP. (2018). Human Development Index Report. UNDP.
50. Gardner H. (2004). "How Education Changes." In M. Suárez- Orozco & D. B. Qin-Hilliard, eds. *Globalization, Culture, and Education in the New Millennium*. Berkley, CA: University of California Press.
51. Xing Y, Xia Y, Kendrick K, Liu X, Wang M, Wu D, Yang H, Jing W, Guo D, Yao D. (2016). Mozart,

- Mozart Rhythm and Retrograde Mozart Effects: Evidences from Behaviours and Neurobiology Bases. Scientific Reports | 6:18744 | doi: 10.1038/srep18744.
52. Kühlmann, AYR., Rooij, A D, Hunink, MGM., Zeeuw, CID, Jeekel, J. (2018). Music Affects Rodents: A Systematic Review of Experimental Research. Frontiers in Behavioral Neuroscience, 12. doi: 10.3389/fnbeh.2018.00301.
53. Mariana P, Veronika D A., Camila C, Nadia J. (2017). Rock influences spatial memory in adult rats, while classical music do not. Psicológica 38, 177-193.
54. Retallack, D. L. (1973). The sound of music and plants. Santa Monica, CA: De Vorss.
55. Chaudhury S, Nag T C, Jain S, Wadhwa S. (2013). Role of sound stimulation in reprogramming brain connectivity. J. Biosci. 38 605–614.
56. Logan B. (2003). Learning before birth: every child deserves giftedness. U.S.: 1st Books Library.
57. Van De Carr FR & Lehrer M. (1997). While You're Expecting... Your Own Prenatal Classroom. Atlanta: Humanics Limited.
58. Manrique B. Make Way for Baby! Babies Can Actually Learn Before Birth. Make Way for Baby website.
59. Ernawati, Hermanto T J, Widjiati. (2008), Comparison of *Rattus norvegicus* offsprings neuronal apoptosis index exposed to Mozart compositions since day 0, day ten of pregnancy and no exposure. Unpublished.
60. Kim C, Lee S, Shin J W, Chung K, Lee S, Shin M, Baek S, Sung Y, Kim C, Kim K. (2013). Music and Noise Influence Neurogenesis and Thickness in Motor. And Somatosensory Cortex of Rat Pups. Int Neurourol J; 17: 107-113.
61. Kim H., Lee M H., Chang H K, Lee T H, Lee H H., Shin M C., Shin M S, Won R., Shin H S & Kim C J. (2006). Influence of prenatal noise and music on the spatial memory and neurogenesis in the hippocampus of developing rats. Brain & Development, 28, 109-114. doi: 10.1016/j.brain dev.05.008.
62. Corballis M C. (2014). Left Brain, Right Brain: Facts and Fantasies. PLoS Biology, 12(1). doi: 10.1371/journal.pbio.1001767.
63. Right Brain vs. Left Brain – What's the Difference? (n.d.). Retrieved from https://www.superduperinc.com/handouts/pdf/202_RightBrainLeftBrain.pdf.
64. Kauser H, Roy S, Pal A, Sreenivas V, Mathu R, Wadhw S, Jain S. (2011). Prenatal Complex Rhythmic Music Sound Stimulation Facilitates Postnatal Spatial Learning but Transiently Impairs Memory in the Domestic Chick. Dev Neurosci, 33(1), 48-56. doi: 10.1159/000322449.
65. Pope K. (2017). The Effects of Jazz and Classical Music on Recall. Journal of Health Education Research & Development, 05(02). Indonesia.
66. Purwanto A H. (2010). Becoming a Jazz Musician: Pola Sosialisasi Musik Jazz pada Beberapa Musik Balejazz. 22-42. in Indonesia.
67. Sanyal T, Kumar V, Nag T C, Jain S, Sreenivas, Wadhwa, S. (2013). Prenatal Loud Music and Noise: Differential Impact on Physiological Arousal, Hippocampal Synaptogenesis and Spatial Behavior in One Day-Old Chicks. PLoS ONE, 8(7), 1-16. doi: 10.1371/journal.pone.0067347.
68. Sanyal T, Palanisamy P, Nag T C, Roy T S, Wadhwa, S. (2013). Effect of prenatal loud music and noise on total number of neurons and glia, neuronal nuclear area and volume of chick brainstem auditory nuclei, field L and hippocampus: a stereological investigation. International Journal of Developmental Neuroscience, 31(4), 234-244. doi: 10.1016/j.ijdevneu.2013.02.004.
69. Kirchberger M, Russo F. (2016). Dynamic Range across Music Genres and the Perception of Dynamic Compression in Hearing-Impaired Listeners. Trends in Hearing, 20, p.23312165166 3054.
70. How Music Promotes Health. (n.d.). Retrieved from <https://www.emedexpert.com/tips/music.shtml>.
71. Poikonen H, Alluri V, Brattico E, Lartillot O, Tervaniemi M, Huotilainen M. (2016). Event-related brain responses while listening to entire pieces of music. Neuroscience, 312, pp.58-73.