

Breast Metastasis from Lung Cancer: A Case Report

L. Mezouar

Received: 6 December 2018 Accepted: 4 January 2019 Published: 15 January 2019

Abstract

Secondary breast tumors are rare. Breast metastases accounted for 0.4 to 2

Index terms—

1 Introduction

Secondary breast tumors are rare. Breast metastases accounted for 0.4 to 2% of all mammary cancers. Melanomas, lymphomas, and lung cancers are most often the primary tumors [1]. Up to now more than 400 cases of mammary metastasis have been reported in the literature [2,3]. We report in our work the clinical case of metastatic lung carcinoma into breast.

2 II.

3 Case Report

A 51-year-old father of two, consulted for headache, vomiting, and right hemiplegia as part of an intracranial hypertension syndrome.

The initial physical examination objectified a fixed mammary mass of 5cm in size in the right breast, the axillary level was free. A mammogram showed the presence of an irregular opacity measuring 45 mm hypoechogenous on ultrasound without microcalcifications, and without axillary adenopathy classified BIRADS 4.

A biopsy of the breast mass was made, whose anatomopathological results showed an undifferentiated tumor process. Thus, an immunohistochemical study was used which showed a low differentiated squamous cell carcinoma compatible with a lung cancer.

A cerebral and cervicothoraco-abdominal CT scan was performed, showing a right apical pulmonary tumor associated with Baret's lymphadenopathy, and also revealed the presence of a right breast mass (already described in mammography.) and bilateral adrenal lesions in favor of secondary lesions. At the cerebral level there were several bilateral temporoparietal lesions with right verminal cerebellar lesion in favor of secondary lesions.

The patient was admitted to radiotherapy department where he received total brain irradiation at a dose of 30 Gy in 10 fractions of 3 Gy /fr over 2 weeks with marked clinical improvement. Afterwards he was treated with palliative chemotherapy based on Carboplatin AUC 6, Paclitaxel 175 mg / m² every 21 days. After the 3rd cycle there was an incomplete remission estimated at more than 70%. The evaluation after the sixth cycle treatment was in favor of stable disease.

The patient was monitored, and 2 months later there was local, mammary and cerebral progression, so he was treated with second line chemotherapy Gemcitabine monotherapy but after 3 courses the patient presented in a critical state with a WHO score of 3 for whom he was referred for palliative care. The patient died 40 days later.

4 III.

5 Discussion

Breast cancer is the most common cancer in the world, however mammary metastases from extramammary cancer are extremely rare accounting for less than 2% of all breast cancers [2,3,6,22].

Breast metastases may be the first revealing symptom in 8 to 33% of cases [7]. These metastases may be associated with ipsilateral axillary adenopathies in more than 50% of cases [7].

The most frequent primary cancers are melanoma, lung cancer, and lymphoma [4,5]. So far the largest study published is that of Stavros Georgiannos, et al which included 60 patients collected over a period of a century

60/14000 presented with breast metastasis (0.43%), of which 95% are female, their age varies between 12 and 90 years, the primary cancers involved in most patients were cutaneous melanomas, large and small cell lung cancers, but there were other tumors, such as endometrial adenocarcinoma, pancreatic adenocarcinoma, retinal melanoma. Other localization may be responsible of these metastases, including uterine, gastric, rectal cancers, and non-Hodgkin's lymphoma [6,8], the latter represents 17% of all mammary metastases [8] As for metastases from gastric cancer, they were described by Krichen et al. in a case of a metastatic gastric adenocarcinoma in the ovary, the primary tumor and ovarian metastasis was resected, but after 4 months a metastatic relapse was detected as an isolated mammary metastasis of the same histological type as the gastric cancer, it was resected but the evolution was marked by the apparition of cutaneous lymphangitis of the chest wall, and emergence of lymph nodes and bone metastases [9]. And this is not the only case of gastric cancer, Boutis et al have also reported a series of 25 cases of gastric cancer associated with mammary metastases of which 13/25 has a form of signet cell ring carcinoma [10]. One of the tumors that can metastasize at the breast level are neuroendocrine tumors, the largest series of secondary breast neuroendocrine tumors reported by Upalakalin et al. [15] including 24 cases, in nine cases mammary metastasis was the revealing site of the primary tumor.

The clinical presentation of these tumors is most often as a well-rounded, hard, and painless lump located at the level of the supero-external quadrant [6,9,12,13,14]. This lump may be associated with axillary metastases [14,15,16]. As well as other metastatic lesions [15].

The mammography shows aregular round or slightly irregular nodule, not containing microcalcifications except in cases of metastases of ovarian origin [13,14,15,16] and at the mammary ultrasound they appear a hypoechogenic, without acoustic shadowing.

Histologically, intramammary metastases have different characteristics of a primary breast cancer, in fact most intra-mammary metastases have histology similar to that of the tumor from which it comes [22].

But one can be mistaken in making the diagnosis of a primitive breast cancer [17,18,19,20]. To overcome this problem we must use immunohistochemistry that will allow us to differentiate a primary cancer from a metastasis, usually for primitive breast cancer the CK7 is positive, CK20 is negative and the hormone receptors are positives. In the case of metastasis, CK 7 (-) and hormone receptors are negatives [14,15]. To guide the diagnosis to the primary tumor, we can look for certain tumor markers such as TTF1 (lung cancer or thyroid cancer) and desmin (soft tissue tumors) [21]. On the other hand, the elevation of some serum markers can be useful to make the diagnosis: ACE (colon cancer), CA19-9 (pancreatic cancer), CA 125 (ovarian cancer, or gastrointestinal cancer)

The management of mammary metastases must be part of the treatment of the primary cancer in question and consists of palliative chemotherapy, and it may be combined with palliative radiotherapy if needed [22]. Surgical treatment is not indicated in these cases except in cases of primary surgery to make the diagnosis [15] or more rarely in the case of a very advanced mammary metastases [17]. Until now there are no studies that support axillary dissection in cases of mammary metastasis associated with axillary adenopathies, and this will only increase postoperative morbidity [23,24].

IV.

6 Conclusion

Breast metastases are exceptional tumors. Their diagnoses are difficult and require a radio-clinicopathological arguments. Their treatment consists of palliative chemotherapy and surgical treatment is to be avoided (mastectomy). Their prognoses depend on the primary tumor but are usually of poor prognosis.

7 bibliography

For example in the case of metastases from ovarian cancer we often found the presence of intrametastasis microcalcifications [14,16].

The prognosis is generally poor and depends on the primary cancer. In the Lee et al study the mean survival in 32 patients with breast metastasis was 13.9 months [15].¹

¹© 2019 Global Journals

[Tanaka et al. ()] 'A case of malignant eccrine spiradenoma metastatic to intramammary lymph node'. Y Tanaka , E Bhunchet , T Shibata . *Breast Cancer* 2008. 15 (2) p. .

[Alva and Shetty-Alva ()] 'An update of tumor metastasis to the breast data'. S Alva , N Shetty-Alva . *Arch Surg* 1999. 134 (4) p. 450.

[Assoulinea et al.] P Assoulinea , * , M.-B Léger-Ravetb , R Saffroyc , J Hamelinc , A Bénissadd , P Husleaga . *Métastases mammaires des cancers broncho-pulmonaires avec mutation d e l'EGFR*,

[Mohaned and Abass ()] 'Axillary Lymph Node Dissection for Breast Cancer: Efficacy and Complication in Developing Countries'. O Mohaned , Abass . JGO.18.00080. *J Glob Oncol* 2018. 4.

[Hajdu and Urban ()] *Cancers metastatic to the breast*, S I Hajdu , J A Urban . *Cancer* 1972. 29 p. .

[Upalakalin et al. ()] 'Carcinoid tumors in the breast'. J N Upalakalin , L C Collins , N Tawa , S Parangi . *Am J Surg* 2006. 191 p. .

[Lee et al.] 'Characteristics of metastasis in the breast from extramammary malignancies'. S K Lee , W W Kim , S H Kim , S M Hur , S Kim , J H Choi . *J Surg Oncol*

[Yeh et al. ()] 'Clinical and ultrasonographic characteristics of breast metastases from extramammary malignancies'. C-N Yeh , C-H Lin , M-F Chen . *Am Surg* 2004. 70 (4) p. .

[Silverman et al. ()] 'Fine needle aspiration cytology of neoplasms metastatic to the breast'. J F Silverman , P S Feldman , J L Covell , W J Frable . *Acta Cytol* 1987. 31 p. .

[Boutis et al. ()] 'Gastric signet-ring adenocarcinoma presenting with breast Metastasis'. A L Boutis , C Andreadis , F Patakliouta , D Mouratidou . *World J Gastroenterol* 2006. 12 p. .

[Alexander et al. ()] 'Isolated breast metastases from gastrointestinal carcinomas: report of two cases'. H R Alexander , A D Turnbull , P Rosen . *J Surg Oncol* 1989. 42 p. .

[Charfia et al. ()] 'Les métastases mammaires: étude anatomoclinique de six cas'. S Charfia , S Maknia , * , A Khanfirb , K Abbes . *J Gyn Obst et Biologie de la Reproduction* 2008. 37 p. .

[Cangiarella et al. ()] 'Malignant melanoma metastatic to the breast: a report of seven cases diagnosed by fine-needle aspiration cytology'. J Cangiarella , W F Symmans , J M Cohen , A Goldenberg , R L Shapiro , J Waisman . *Cancer* 1998. 84 (3) p. .

[Shaw et al. ()] 'Malignant pheochromocytoma metastasising to the breast'. C Shaw , D M O'hanlon , O 'keane , C Kerin , MJ . *Ir J Med Sci* 2003. 172 (1) p. .

[Domanski ()] 'Metastases to the breast from extramammary neoplasms: a report of six cases with diagnosis by fine needle aspiration cytology'. H Domanski . *Acta Cytol* 1996. 40 p. .

[Williams et al. ()] 'Metastases to the breast from nonbreast solid neoplasms: presentation and determinants of survival'. S A Williams , Ehlers 2nd R A , K K Hunt , Yi M Kuerer , H M Singletary , SE . *Cancer* 2007. 110 (4) p. .

[Ravdel et al. ()] 'Metastatic melanoma in the breast: a report of 27 cases'. L Ravdel , W A Robinson , K Lewis , R Gonzalez . *J Surg Oncol* 2006. 94 (2) p. .

[Maknia et al. ()] 'Metastatic signet ring cell carcinoma to the breast from stomach'. S Maknia , K Abbasa , A Khanfirb , * , M Frikhab , T Sellami , Boudawara . *Cancer/Radiothérapie* 2007. 11 p. .

[Duncan et al. ()] 'Non-Hodgkin's lymphoma of the breast: a review of 18 primary and secondary cases'. V E Duncan , V V Reddy , N C Jhala , D C Chhieng , D Jhala . *Ann Diagn Pathol* 2006. 10 p. .

[Khalifeh et al. ()] 'Primary peritoneal serous carcinoma presenting as inflammatory breast cancer'. I Khalifeh , M T Deavers , M Cristofanilli , R L Coleman , A Malpica , M Gilcrease . *Breast J* 2009. 15 (2) p. .

[Whelan (2015)] 'Regional Nodal Irradiation in Early-Stage Breast Cancer'. T J Whelan . 10.1056/NEJ-Moa1415340. *N Engl J Med* 2015 Jul 23. 373 (4) p. .

[Georgiannos et al. ()] 'Secondary neoplasms of the breast. A survey of the 20th century'. S N Georgiannos , J C Aleong , A W Goode , M Sheaff . *Cancer* 2001. 92 p. .

[Recine et al. ()] 'Serous carcinoma of the ovary and peritoneum with metastases to the breast and axillary lymph nodes: a potential pitfall'. M A Recine , M T Deavers , L P Middleton , E G Silva , A Malpica . *Am J Surg Pathol* 2004. 28 (12) p. .

[Lee ()] 'The histological diagnosis of metastases to the breast from extramammary malignancies'. Ahs Lee . *J Clin Pathol* 2007. 60 (12) p. .