

Successful Management of Spondylodiscitis in a three Years Old Girl: A Case Report

Ali ALSharqi

Received: 7 December 2019 Accepted: 4 January 2020 Published: 15 January 2020

Abstract

Spondylodiscitis considered one of the rare diseases that cause back pain. The disease pathology is not yet been clearly known, however, in most patients the disease thought to be spreading hematogenously from a previously existing site of infection. We report two years and 11 months old child, previously healthy girl, presented to the emergency department with two weeks' history of weakness of the lower extremities and lumbar back pain with slightly arched back. She had a complete recovery with early intervention and complete course of antibiotics.

Index terms— spondylodiscitis, MRI, Staph. aureus, antibiotics, case report, Oman.

I. Introduction podylodiscitis (SD), is infectious process of the spine involving vertebral bodies and intervertebral discs. It remains a rare condition with an estimated incidence of around one to two cases in 30000. 2 this case report describes spondylodiscitis in tow years old girl who presented with acute back pain, irritability and inability to walk. SD although it's rare disease, it should be kept as one of the deferential diagnosis in children present with non-traumatic back pain. Staphylococcus aureus is the causative agent of SD accounting for 80% of the cases. 2,7,14,15,16,17,18 Treatment of SD is usually a combination of both pharmacological and non-pharmacological.

1 II. Case Presentation

A two years and 11 months old toddler, previously healthy girl, presented to the emergency department in a tertiary center in Muscat, Oman in 2019, with a two weeks' history of weakness of the lower extremities with back pain and slightly arched back. There was no history of trauma, unexplained weight loss, or any other systemic manifestation. There was no history of fever, joint pain or skeletal deformity, skin rash, seizure, or photophobia. She was not known to have any chronic diseases. She was up to date with her vaccinations. Her parents reported no exposure to individuals with similar symptoms. In addition, none of her family members and neighbors had recently suffered from chronic cough or unexplained weight loss. There was no history of previous admission.

On admission, she was irritable, her vital signs were within normal ranges. Her physical examination revealed normal gait with slight hyperextension of lower back. In addition, she was bearing weight with support due to pain and there was slight pain in lower back while flexion and extension of the back. Otherwise, no muscle wasting, full range of movement of all joints actively and passively, with normal tone, power and reflexes. A blood investigations revealed complete blood count: Hemoglobin of 9.2 g/dl, increase in platelet count of 736 * 10⁹ /L, normal white cell counts and slight increased level of acute phase reactants C-reactive protein(CRP) of 11.1 mg/L and erythrocyte sedimentation rate of 42mm/hour. Pelvic X-RAY was done and was reported as normal and Ultrasound hip showed no fluid in hip joint. She was started on non-steroidal anti-inflammatory medications and on vancomycin and ceftriaxone but the next day vancomycin was switched to flucloxacillin and child showed clinical improvement. Lumbar puncture was done as she was inactive and it showed normal microscopy, count, protein and glucose, with negative CSF and blood culture. On day five of admission, MRI done which showed bone marrow changes of L4 and L5 vertebrae associated with endplates irregularities and mild destruction, with loss of intervertebral space and indentation of thecal sac suggestive of spodylodiscitis due to pyogenic or granulomatous infection. Child was tested for Q fever, tuberculosis and brucellosis in which all test were negative. Child was treated with ceftriaxone and flucloxacillin for two weeks as she was responding

to the treatment and finally discharged on oral cefdinir for four weeks after consultation with infectious disease doctor. On subsequent follow up as outpatient she showed marked improvement and she started to regain her full movement of lower limbs.

2 III. Discussion

We report almost three years old girl with two weeks' history of lower back pain and lower limb weakness in whom MRI spine showed destructive changes of L4 and L5 with high ESR of 42 mm/hour and all other tests were not significant. When lumbar pain is accompanied by significant irritability, (as was evident in our case), this should lead pediatrician to include infectious discitis among the differential diagnosis. Kang et al 1 reported that approximately 60% of discitis and SD cases were diagnosed in children <3 years old showed that irritability was the most common among all other symptoms at the time of disease presentation. Discitis and SD are infectious processes of the spine involving vertebral bodies and intervertebral discs. It remains a rare condition with an estimated incidence of around one to two cases in 30,000 2 . In our case, there was no delay in establishing the correct diagnosis where it has been established in the fifth day of admission. Delays of diagnosis for four to six months have been reported 2,3,4 . These delays are attributed to the often non-specific clinical presentation of children with discitis or SD and their inability to describe the site of discomfort 2,5,6 . This delay can lead to an increase risk of permanent abnormalities 7 . Its pathophysiology has not yet been clearly established, but in most patient, pathogens reach the spine hematogenously, starting from a previously existing site of infection 2,7 . A prodrome with a distant focus of infection has been identified in most cases. Mylona et al 8 , described these to include the genitourinary tract 17%, skin and soft tissue 11%, intravascular devices 5%, gastrointestinal tract 5%, respiratory tract 2% and the oral cavity 2%. A wide range of pathogens can cause this disease and many studies showed it is primarily monomicrobial bacterial infection. Many attempts to identify the causative pathogen of Discitis and SD of children through blood and/or vertebral aspiration cultures have failed to identify the organisms; causing related problems in selecting the most appropriate antibiotic therapy 9,10,11,12,13 . When positive, pyogenic bacteria are usually detected, with Staphylococcus aureus being the cause of discitis and SD in approximately 80% of the cases that occur in first months of life and in most of those that develop in older children 2,7,14,15,16,17,18 . The most specific imaging method to diagnose discitis is MRI 19 . Intravenous antibiotic treatment, analgesia and physical rehabilitation treatment showed complete recovery in most cases. Treatment include pharmacological like antibiotics and non-pharmacological such as physiotherapy and bed case back pain, irritability, and walking difficulties are rest 20 . Mortality has dropped from 25-26% 21,22 to less than 5% 23 with antibiotics treatment.



Figure 1: FFigure 1 :

76 [J Antimicrob Chemother ()] , 10.1093/jac/dkq303. *J Antimicrob Chemother* 2010. 65 (3) p. . (Suppl)

77 [Cornett et al. ()] ‘Bacterial spine infections in adults: Evaluation and management’. C A Cornett , S A Vincent
78 , J Crow , A Hewlett . *J. Am. Acad. Orthop. Surg* 2016. 24 p. . (PubMed)

79 [Fernandez et al. ()] ‘Discitis and vertebral osteomyelitis in children: An 18-year review’. M Fernandez , C L
80 Carrol , C J Baker . *Pediatrics* 2000. 105 p. . (PubMed)

81 [Karabouta et al. ()] ‘Discitis in toddlers: a case series and review’. Z Karabouta , I Bisbinas , A Davidson , L
82 L Goldsworthy . *Acta Paediatr* 2005. 94 p. .

83 [Brown et al. ()] ‘Discitis in young children’. R Brown , M Hussain , K Mchugh , V Novelli , D Jones . *J. Bone
84 Jt. Surg* 2001. 831 p. .

85 [Crawford et al. ()] ‘Diskitis in children’. A H Crawford , D W Kucharzyk , R Ruda , H C Smitherman . *Clin.
86 Orthop. Relat. Res* 1991. 266 p. . (PubMed)

87 [Cushing ()] ‘Diskitis in children’. A H Cushing . *Clin. Infect. Dis* 1993. 17 p. . (PubMed)

88 [Cottle and Riordan ()] ‘Infectious spondylodiscitis’. L Cottle , T Riordan . *J. Infect* 2008. 56 p. . (PubMed)

89 [Rocco and Eyring ()] ‘Intervertebral disc infections in children’. H D Rocco , E J Eyring . *Am J Dis Child* 1972.
90 123 p. .

91 [Ryoppy et al. ()] ‘Nonspecific diskitis in children’. S Ryoppy , J Jaaskelainen , J Rapola , A Alberty . *Clin.
92 Orthop. Relat. Res* 1993. 297 p. .

93 [Garron et al. ()] ‘Nontuberculous spondylodiscitis in children’. E Garron , E Viehweger , F Launay . *J
94 PediatrOrthop* 2002. 22 p. .

95 [Ceroni et al. ()] ‘Osteoarticular infections in young children: What has changed over the last years’. D Ceroni
96 , G Kampouroglou , R Valaikaite , R Anderson Dellallana , D Salvo . *Swiss Med. Wkly* 2014. w13971. 144.
97 (PubMed)

98 [Bauman and Stifel ()] ‘Osteomyelitis of the spine’. G I Bauman , R E Stifel . *Ann Surg* 1923. 78 p. .

99 [Fuster et al. ()] ‘Prospective comparison of whole-body (18)F-FDG PET/CT and MRI of the spine in the
100 diagnosis of haematogenous spondylodiscitis’. D Fuster , X Tomás , M Mayoral , A Soriano , F Manchón ,
101 C Cardenal , A Monegal , U Granados , S Garcia , F Pons . *Eur. J. Nucl. Med. Mol. Imaging* 2015. 42 p. .
102 (PubMed)

103 [Digby and Kersley ()] ‘Pyogenic non-tuberculous spinal infection: An analysis of thirty cases’. J M Digby , J B
104 Kersley . *J. Bone Jt. Surg. Br* 1979. 61 p. .

105 [Kulowski ()] ‘Pyogenic osteomyelitis of the spine: an analysis and discussion of 102 cases’. J Kulowski . *J Bone
106 Joint Surg Am* 1936. 18 p. .

107 [Sapico and Montgomerie ()] ‘Pyogenic vertebral osteomyelitis: report of nine cases and review of the literature’.
108 F L Sapico , J Z Montgomerie . *Rev Infect Dis* 1979. 1 p. .

109 [Kayser et al. ()] ‘Spondylodiscitis in childhood: Results of a longterm study’. R Kayser , K Mahlfeld , M Greulich
110 , H Grasshoff . *Spine* 2005. 30 p. . (PubMed)

111 [Rubio Gribble et al. ()] ‘Spondylodiscitis in the autonomus community of Madrid (Spain)’. B Rubio Gribble , C
112 Calvo Rey , J Garcia-Consuegra , L Ciria Calabria , M L Navarro Gomez , J T Ramos Amador . *An. Pediatr*
113 2005. 62 p. . (PubMed)

114 [Tapia Moreno et al. ()] ‘Spondylodiscitis: Diagnosis and medium-long term follow up of 18 cases’. R Tapia
115 Moreno , M G Espinosa Fernandez , M I Martinez Leon , J M Gonzalez Gomez , P Moreno Pascual . *An.
116 Pediatr* 2009. 71 p. . (PubMed)

117 [Kang et al. ()] ‘The etiology, clinical presentation and long-term outcome of spondylodiscitis in children’. H M
118 Kang , E H Choi , H J Lee , K W Yun , C K Lee , T J Cho , J E Cheon , H Lee . *Pediatr. Infect. Dis. J*
119 2016. 35 p. . (PubMed)

120 [Wenger et al. ()] ‘The spectrum of intervertebral disc-space infection in children’. D R Wenger , W P Bobechko
121 , D L Gilday . *J Bone Joint Surg [Am]* 1978. p. .

122 [Modic et al. ()] ‘Vertebral osteomyelitis: Assessment using MR’. M T Modic , D H Feiglin , D W Piraino , F
123 Boumphrey , M A Weinstein , P M Duchesneau . *Radiology* 1985. 157 p. . (PubMed)