

Pathological Proximal Femoral Fracture in Mc-Cune Albright Syndrome: A Case Report

Apoorv Sehgal¹ and Pratyush Shahi²

¹ University College of Medical Sciences

Received: 7 December 2019 Accepted: 5 January 2020 Published: 15 January 2020

Abstract

A 25-year-old female present to us in the emergency department with atraumatic fracture of the proximal femur. She had a cafe-au-lait spot on the face and gave history of precocious puberty. A skeletal survey showed polyostotic fibrous dysplasia. A clinico-radiological diagnosis of McCune-Albright syndrome was made which was further confirmed on biopsy. An extra-medullary fixation was done and at 6 months follow-up, the fracture had united and the patient could walk without support.

Index terms—

1 Introduction

fibrous dysplasia/McCune-Albright syndrome (FD/MAS) occurs due to somatic gain-of-function mutations of the GNAS gene. These mutations cause over activity in the target tissues leading to clinical features that vary in severity and age of onset. Since the disease is rare and can have atypical presentation, the diagnosis can be missed. We report the case of a young female with McCune-Albright syndrome presenting to us with pathological fracture of the proximal femur and highlight the challenges one can face in fixation of such fractures.

2 II.

3 Case Presentation

A 25-year-old female patient presented to us with a history of sudden onset of pain in her right hip and inability to bear weight over the affected limb. There was no history of trauma, fever or weight loss. On examination, the right lower extremity was externally rotated and shortened. Movements at the right hip elicited pain. The patient was of short stature and had a café-au-lait skin macule with jagged, irregular borders (coast of Maine) on her face with short stature. (Figure ??) On further inquiry, she gave a history of precocious puberty.

Radiographs revealed lytic expansile lesion with ground-glass appearance and thinning of cortex in metaphyseal region of the right proximal femur associated with a pathological subtrochanteric fracture. Other sites like skull, radius and phalanges also had similar lytic lesions. (Figure 2)

Based on clinical and radiological features, a diagnosis of McCune-Albright syndrome with pathological right subtrochanteric fracture was made.

The patient was put on skin traction for the right lower limb and advised to undergo surgery for the subtrochanteric fracture of right femur in view of the nature of the fracture and the requirement of early ambulation. After proper consent, the patient was taken up for surgery. Fracture was reduced on a traction table after opening the fracture area. Biopsy was taken from the fracture site followed by reduction and fixation with distal femoral locking plate of opposite side in reverse configuration. The initial plan was of internal fixation with cephalomedullary nail, but due to very narrow diameter of the medullary canal, an extramedullary fixation had to be done. (Figure ??) The patient was discharged on postoperative day 5 after confirming healthy state of the wound. She was started on partial weight-bearing with crutches at 2 weeks continuing to full weight-bearing at 6 weeks. At 6 months follow-up, the patient could walk without support and with no residual complaints.

4 Discussion

McCune-Albright syndrome is a rare sporadic disease caused by somatic postzygotic activating mutations in the GNAS gene. It is characterised by polyostotic fibrous dysplasia, café-au-lait skin spots and variable hyperfunctional endocrinopathies. 2 A fibrous dysplasia lesion predisposes a person to have a pathological fracture and if involving a large area can lead to deformity of the bone. Although a small lesion can be left as such, a larger lesion causing pain or threatening to fracture should be curetted and bone grafted even though there is a high tendency for it to recur. If the site and extent of the lesion necessitates, prophylactic fixation should be done. The management of fibrous dysplasia becomes more complicated when it is associated with extra-skeletal manifestations. 1,3 A lesion of the proximal femur causes more pain, fractures and deformity than any other skeletal localization of the disease. [2][3][4][5] A pathological fracture of the proximal femur secondary to fibrous dysplasia poses several intra-operative challenges. A surgeon should do proper pre-operative planning and have backup fixation devices. Also, good post-operative wound care and early rehabilitation should be done.



Figure 1: Figure1:Figure 2 :

1

-
- 56 [Leet et al. ()] ‘Fracture incidence in polyostotic fibrous dysplasia and the McCune-Albright syndrome’. A I Leet
57 , C Chebli , H Kushner , C C Chen , M H Kelly , B A Brillante , P G Robey , P Bianco , S Wientroub , M
58 T Collins . 10.1359/JBMR.0301262. *J Bone Miner Res* 2004. 19 p. .
- 59 [O’sullivan and Zacharin ()] ‘Intramedullary rodding and bisphosphonate treatment of polyostotic fibrous dys-
60 plasia associated with the McCune-Albright syndrome’. M O’sullivan , M Zacharin . *J Pediatr Orthop* 2002.
61 22 p. .
- 62 [Lichtenstein ()] *Polyostotic fibrous dysplasia. Arch urg*, L Lichtenstein . 10.1001/archsurg.1938.01190230153012.
63 1938. 36 p. .
- 64 [Harris et al. ()] ‘The natural history of fibrous dysplasia. An orthopaedic, pathological, and roentgenographic
65 study’. W H Harris , H R Dudley , Jr , R J Barry . 10.2106/00004623-196244020-00001. *J Bone Joint Surg*
66 *Am* 1962. p. .
- 67 [Stanton et al. ()] ‘The surgical management of fibrous dysplasia of bone’. R P Stanton , E Ippolito , D Springfield
68 , L Lindaman , S Wientroub , A Leet . 10.1186/1750-1172-7-S1-S1. *Orphanet J Rare Dis* 2012. 7 (1) p. S1.
69 (Suppl)