



GLOBAL JOURNAL OF MEDICAL RESEARCH: B
PHARMA, DRUG DISCOVERY, TOXICOLOGY & MEDICINE
Volume 20 Issue 1 Version 1.0 Year 2020
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Evaluation of Methanolic Extract of *Hypericum Mysorensense* Ointment for its Wound Healing Activity

By Sankar C, Muthukumar S, Arulkumaran G, Vinesha R, Manimekalaim
& Sandeep George Simson

KMCH College of Pharmacy

Abstract- The main objective of the present study is to formulate and evaluate semi-solid dosage forms of *Hypericum mysorensense* for its wound healing activity. The stem of *Hypericummysorensense* was extracted by continuous hot percolation/ soxhletation using methanol as solvent and was evaluated for its phytochemical property, *in-vitro* antimicrobial and *in-vitro* antioxidant activity. Using this methanolic extract, semi-solid dosage form (ointment) was formulated. Ointment was prepared using 2% concentration of the extract by fusion method using emulsifying ointment. Hence, the ointment containing 2% methanolic extract of *Hypericummysorensense* was evaluated for its wound healing activity using excision wound model. The effect produced by the extract ointment, in terms of wound contracting ability, wound closure time and histopathological characteristics were comparable to that of a standard drug povidone iodine ointment. Therefore, the present study concluded that the *Hypericum mysorensense* at 2% concentration of the methanolic extract formulated as an ointment showed a better wound healing formulation for better patient care and pharmacoeconomical.

Keywords: *hypericum mysorensense*, excision wound model, ointment.

GJMR-B Classification: NLMC Code: QV 55



Strictly as per the compliance and regulations of:



© 2020. Sankar C, Muthukumar S, Arulkumaran G, Vinesha R, Manimekalaim & Sandeep George Simson. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Evaluation of Methanolic Extract of *Hypericum Mysorensense* Ointment for its Wound Healing Activity

Sankar C ^α, Muthukumar S ^ο, Arulkumaran G ^ρ, Vinesha R ^ω, Manimekalaim [¥]
& Sandeep George Simson [§]

Abstract- The main objective of the present study is to formulate and evaluate semi-solid dosage forms of *Hypericum mysorensense* for its wound healing activity. The stem of *Hypericummysorensense* was extracted by continuous hot percolation/ soxhletation using methanol as solvent and was evaluated for its phytochemical property, *in-vitro* antimicrobial and *in-vitro* antioxidant activity. Using this methanolic extract, semi-solid dosage form (ointment) was formulated. Ointment was prepared using 2% concentration of the extract by fusion method using emulsifying ointment. Hence, the ointment containing 2% methanolic extract of *Hypericummysorensense* was evaluated for its wound healing activity using excision wound model. The effect produced by the extract ointment, in terms of wound contracting ability, wound closure time and histopathological characteristics were comparable to that of a standard drug povidone iodine ointment. Therefore, the present study concluded that the *Hypericum mysorensense* at 2% concentration of the methanolic extract formulated as an ointment showed a better wound healing formulation for better patient care and pharmacoeconomical.

Keywords: *hypericum mysorensense*, excision wound model, ointment.

I. INTRODUCTION

a) Overview of Traditional Herbal Medicine

India is a birthplace of indigenous medicine such as Siddha, Ayurveda, and Unani where many herbs have been used for treatment of human ailments. About 65% of total global population remains dependent on traditional medicines for their primary healthcare. Herbs are occupying a comeback and an 'Herbal Renaissance' is blooming across the world. They have been evidently prized for their medicinal, flavoring and aromatic qualities for centuries, yet for a while they were over shadowed by synthetic products of modern civilization. Folk medicine is generally defined as traditional medicine that is practiced by non-professional healers or embodied in local custom or lore, generally involving the use of natural and especially herbal remedies.

Corresponding Author α: M. pharm., Assistant Professor, Department of Pharmaceutics, KMCH college of Pharmacy, Coimbatore, Tamil Nadu, India, (Affiliated to the Tamil Nadu Dr. M. G. R. Medical University). e-mail: pharmmuthu@gmail.com

Author α ρ ω ¥ §: Department of Pharmaceutics, KMCH College of pharmacy, Coimbatore, Tamil Nadu, India (Affiliated to the Tamil Nadu Dr. M. G. R. Medical University).

Once having realized their sources and adverse effects, people are going back to nature with hopes of safety and security. The rich treasure of herbal drugs is forming a boon for our society. Plant derived compounds, apart from their nutritive values, could serve as important therapeutic weapons to fight various human and animal diseases, thereby making them indispensable in traditional medicine for treating a number of diseases. Plant drugs, popularly known as herbal medicines have since been unabatedly used to treat various diseases. The major challenge is to protect traditional knowledge and will prove to be a beneficial asset to our human surrounding. For all the ailments herbal formulations are proved to be effective without any side effects commonly seen with allopathic drugs.

II. OINTMENT

An ointment is a homogeneous, viscous, semi-solid preparation, most commonly greasy, thick oil (oil 80% - water 20%) with a high viscosity which is intended for external application to the skin or mucous membranes. Ointments have a water number that defines the maximum amount of water that it can contain. They are used as emollients or for the application of active ingredients to the skin for protective, therapeutic, or prophylactic purposes and where a degree of occlusion is desired. Ointments are used topically on a variety of body surfaces. These include the skin and the mucous membranes of the eye (an eye ointment), vagina, anus, and nose. An ointment may or may not be medicated. Ointments are usually very moisturizing, and good for dry skin. They have a low risk of sensitization due to having few ingredients beyond the base oil or fat, and low irritation risk. There is typically little variability between brands of generics and name brand drugs. They are often disliked by patients due to greasiness.

III. WOUND HEALING ACTIVITY

Wounds are inescapable events in life. Wounds may arise due to physical, chemical or microbial agents. Wound healing involves a complex series of interactions between different cell types, cytokine mediators, and the extracellular matrix. The phases of normal wound

healing include hemostasis, inflammation, proliferation and remodeling. Each phase of wound healing is distinct, although the wound healing process is continuous, with each phase overlapping the next. Because successful wound healing requires adequate blood and nutrients to be supplied to the site of damaged tissue.

IV. MATERIALS AND METHODS

a) Collection And Extraction

The leaves of *Hypericum mysorensense* was collected freshly during the month of October from Ooty, India, identified and authenticated. The collected plant, *Hypericum mysorensense* was extracted by continuous hot percolation (soxhletation). 200g of powdered stem of *Hypericum mysorensense* was defatted using petroleum ether. The marc obtained from the powdered plant part was successfully extracted with 250 ml of methanol by using soxhlet apparatus. The extraction was carried out for 48 hours. After extraction, the solvents were distilled out; the concentrated residues were analyzed by chemical tests.

b) Preparation of Semi-Solid Formulation

i. Formulation of Ointment

Preparation of Ointment Base(Emulsifying Ointment)

Emulsifying wax - 50g, White soft paraffin - 20g, Liquid paraffin - 30g

Procedure

Required quantities of emulsifying wax, liquid paraffin and white soft paraffin were weighed and melted. To this, adequate quantity of methanolic extract of plant was added and stirred well until a homogeneous mass were obtained. The composition of the prepared herbal ointment (F1) is listed in Table I.

Table I: Composition of Ointment

Sl. No.	Ingredients	F1
1	Methanolic extract of <i>Hypericum mysorensense</i> stem	2gm
2	Emulsifying Ointment	q.s to 100gm

c) Pharmacological Evaluation of the Formulated Ointment

i. Acute Skin Irritation Study

The primary skin irritation test was performed on albino rats and weighing about 150-200gm. The animals were maintained on standard animal feed and had free access to water *ad libitum*. The animals were kept under standard laboratory condition. The total mass was divided into four batches, each batch containing six animals. Two batches of each were used for control and test. Dorsal hairs at the back of the rats were clipped off one day prior to the commencement of the study.

Animals showing normal skin texture were housed individually in cages with meshes to avoid contact with the bedding. 50mg of the each formulation of different concentrations were applied over one square centimeter area of intact and abraded skin to different animals. Aqueous solution of 0.8% formalin was applied as standard irritant. The animals were observed for seven days for any signs of oedema and erythema.

ii. Wound Healing Activity

a. Animals

Wister rat of male sex, weighing 150-200g were used. All animals were obtained from KMCH College of Pharmacy. All animals were housed for at least one week in the laboratory animal room prior to testing. The selected animals were housed in polypropylene cages in standard environmental conditions (20-25°C). Fed with standard rodent diet and water *ad libitum*. The experiments on animals were conducted in accordance with the international accepted principles for laboratory animal use and the experimental protocols duly approved by the institutional ethical committee, KMCH college of pharmacy, Coimbatore, Reg No. 685/Po/02/a/CPCSEA.Dt:21st Aug 2002, IAEC No: KMCRET/M.Pharm/9/2013-14

V. RESULTS

The colour of the ointment was dark green with characteristic odour. The result clearly indicated that the ointment showed good spreadability and extrudability. The ointment showed a pH that lie in the normal pH range of human skin and is easily diffusible.

Pharmacological evaluation of the formulated ointment

a) Wound healing activity

Table 7: Percentage Wound closure

Test Compounds	0 th day	3 rd day	7 th day	13 th day	17 th day	19 th day
	% WC	% WC	% WC	% WC	%WC	%WC
STD	0	15.7 ±0.29	39.8±0.77	81.8±1.09	88.5±1.25	95.5±1.10
Ointment Base	0	13.5 ±0.49	27±0.83	52.5±1.24	66.8±1.24	88.2±1.88
Control	0	27.5±0.58	23.6±2.03	74.2±0.99	72±0.99	90.1±0.99
Ointment Containing 2%Extract	0	10±0.42	13.8±1.48	55.4±1.08	84.1±1.08	96.9±1.08

Results were expressed as mean (mm) ± SEM and were compared with the corresponding control group (simple ointment B.P.) by applying ANOVA test. P value was set <0.05 for all analyses.

Table 8: Figure showing percentage wound reduction

Test compounds	0 th day	3 rd day	7 th day	13 th day	17 th day	19 th day
	%WR	%WR	%WR	%WR	%WR	%WR
STD	100±0.00	84.3±0.29	60.2±0.77	18.2±1.09	11.5±1.25	4.42±1.10
Ointment Base	100±0.00	86.5±0.49	73±0.83	47.5±1.24	33.2±1.24	11.8±1.88
Control	100±0.00	72.5±0.58	76.4±2.03	25.8±0.99	28±0.99	9.81±0.99
Ointment Containing 2% Extract	100±0.00	90±0.42	86.1±1.48	44.5±1.08	15.8±1.08	3.02±1.08

Results were expressed as mean (mm) ± SEM and were compared with the corresponding control group (simple ointment B.P.) by applying ANOVA test. P value was set <0.05 for all analyses.

Excision Wound On 0th Day

Only Wound Wound + Ointment Base



Wound + Standard



Wound +Oint Containing 2% Meoh.Hms

Figure: Figure showing the wound healing activity from 0thday — 21th

Excision Wound on- 21st Day

Only Wound



Wound + Ointment Base



Wound + Oint Containing 2% Meoh.Hms



Wound+Standard

VI. DISCUSSION

Hypericum mysorensense is an ornamental plant belonging to the family Hypericaceae, having antibacterial activity, against both gram-positive and gram-negative bacteria. Leaf and flower have strong antioxidant potential and used for liver disorders. Its stem is having strong antitumor, antipsychic and antiviral activities. Among the various indications where traditional herbal medicines are used, skin and skin related disorders are ranked top. Thus, the main objective of the present study is to formulate and evaluate semi-solid dosage forms of *Hypericum mysorensense* for its wound healing activity. The stem of *Hypericum mysorensense* was extracted by continuous hot percolation/ soxhletation using methanol as solvent. Using this methanolic extract, semi-solid dosage form (ointment) was formulated. Ointment was prepared with 2% methanolic extract using emulsifying ointment base. Hence, the ointment containing 2% Methanolic Extract of *Hypericum mysorensense* stem was evaluated for its wound healing activity using excision wound model. The formulation did not produce any skin irritation for about a week when applied over the skin. Wister rats weighing around 150-200g were used for the study. They were divided into 4 groups consisting of 6 rats each. The 4 groups include standard (povidone iodine ointment), control (only wound), rats treated with ointment base and rats treated with 2% ointment. The effect produced by the extract ointment, in terms of wound contracting ability, wound closure time and histopathological characteristics were comparable to that of a standard drug, povidone iodine ointment.

Thereby, the ointment formulated using 2% methanolic extract of *Hypericum mysorensense* was found to possess wound healing activity.

VII. CONCLUSION

The effect produced by the extract ointment, in terms of wound contracting ability, wound closure time and histopathological characteristics were comparable to that of a standard drug, povidone iodine ointment. Therefore, the present study concluded the wound healing activity of *Hypericum mysorensense* stem at 2% concentration of the methanolic extract formulated as an ointment.

ACKNOWLEDGEMENT

The authors are grateful to the authorities of KMCH College of Pharmacy, Coimbatore for the facilities.

Conflict of Interests

The authors declare that they have no conflicts of interest.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Yogesh Sharma, G. Jeyabalan and Ramandeep Singh: Current Aspect of Wound Healing Agents From Medicinal Plants. A Review, Journal of Medicinal Plants Studies, 2013;1(3):1-11.
2. Meher-Homji VM: Phytogeography of the South Indian Hill Stations. Bulletin of the Torrey Botanical Club, 94(4), 1967: 230-242.

3. Laurence M. Harwood, Christopher J. Moody: Experimental organic chemistry. Principles and Practice (Second edition), 1999: 122–125.
4. Dr. Mohammed Ali: Textbook of Pharmacognosy. CBS publishers and Distributors, New Delhi, 2nd edition, 1998: 20-22.
5. Dr. K.M. Nadakarni: Indian Meteria Medica. Popular Prakashan Private Limited, Mumbai, Third edition, VOL(I), 2002: 1317.
6. P.K. Warriar: Indian medicinal plants. Orient Longman Ltd, Chennai, Vol (5), 1996:445.
7. Pulk K. Mukherjee: Quality control of Herbal Drugs(An approach to Evaluation of Botanicals). Business horizon Publishers, 1st edition, 2002: 131-159,177,246-378,446-447.
8. Dr. Kokate C.K: Practical Pharmacognosy. Vallabh Prakashan Publishers, Fourth edition, 2004: 107-111.
9. Sundarajan.: Practical manual of Microbiology. Vol (1), 1st edition, 1998:34-35.
10. Ashok Ratan : Antimicrobial Activity in Laboratory Medicine. 1st edition, 2000: 67-90.
11. Almagboul, A.Z., Bashir, A.K, Farouk, A, Salih A.K.M: Antimicrobial activity of certain Sudanese plants used in folkloric medicine. Screening of antibacterial activity, *Fitoterapia*, 1985:331-337.
12. Raghu Chantrashekhar H: Antioxidant activity of Hypericum hookerianum. Wight & Arn. Manipal College of Pharmaceutical Science, 2009; Vol 23(13):1240-51.
13. Roberta R: Antioxidant activity applying an improved ABTS radical cation decolourization assay. Free radical biology and Medicine, 1999; Vol 26(9/10): 1231-1237.
14. Uma Shankar Mishra, P. N. Murthy, Debananda Mishra, Kanhu Charana Sahu: Formulation and Standardisation of herbal gel containing methanolic extract of Calophyllum inophyllum. American Journal of Pharma Tech Research, Vol I, Issue 4, 2011:276-289.
15. Barhate SD, Bavaskar KR, Saoji YS, Potdar M, Gholap TN: Development of Transdermal drug delivery system of Ketoprofen. Int. J. Parma. Res. Develop. 2009; 1(10): 1-7.
16. Purnima Kumari, Poonam Yadav, Priya Rani Verma¹, Sandeep Kumar and Arvind Arya A, Review on wound healing properties of Indian medicinal plants. Indian journal of fundamental and applied life sciences, 2013; 3(1):220-232.