



GLOBAL JOURNAL OF MEDICAL RESEARCH: B  
PHARMA, DRUG DISCOVERY, TOXICOLOGY & MEDICINE  
Volume 20 Issue 2 Version 1.0 Year 2020  
Type: Double Blind Peer Reviewed International Research Journal  
Publisher: Global Journals  
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

## *In-Vitro* Antioxidant, Anti-Inflammatory and Anti-Microbial Activity of *Carica Papaya* Seeds

By Pooja G. Singh, Madhu S. B, Shailasreesekhar, Gopenath TS,  
Kanthesh M. Basalingappa & Dr. Sushma BV

**Abstract-** Papaya seeds are reported to have higher therapeutic potential in comparison to the fruits in which they reside. Thus, the present in-vitro study aimed to evaluate and compare the anti-oxidant, anti-inflammatory and anti-microbial effect of seed extracts on *Carica papaya* L. (Caricaceae). The bioactive from the seeds were sequentially fractionated with hexane, chloroform, diethyl ether, and methanol in the increasing order of polarity. Total phenolic and flavonoid contents were estimated. These extracts were assessed for an antioxidant property by 1, 1-diphenyl-2-picryl-hydroxyl (DPPH) method and reducing power assay was carried out using the FeCl<sub>3</sub> method. Inhibition of 15-lipoxygenase (LOX) by these extracts at 5 - 25µg to assess anti-inflammatory capacity was studied.

**Keywords:** *carica papaya* L., seed extracts, phytochemical analysis, antioxidant, lipoxygenase inhibition, antimicrobial activity.

**GJMR-B Classification:** NLMC Code: QV 738



*Strictly as per the compliance and regulations of:*



# In Vitro Antioxidant, Anti-Inflammatory and Anti-Microbial Activity of *Carica Papaya* Seeds

Pooja G. Singh <sup>α</sup>, Madhu S. B <sup>σ</sup>, Shailasreesekhar <sup>ρ</sup>, Gopenath TS <sup>ω</sup>, Kanthesh M. Basalingappa <sup>¥</sup>  
& Dr. Sushma BV <sup>§</sup>

**Abstract-** Papaya seeds are reported to have higher therapeutic potential in comparison to the fruits in which they reside. Thus, the present in-vitro study aimed to evaluate and compare the anti-oxidant, anti-inflammatory and anti-microbial effect of seed extracts on *Carica papaya* L. (Caricaceae). The bioactive form the seeds were sequentially fractionated with hexane, chloroform, diethyl ether, and methanol in the increasing order of polarity. Total phenolic and flavonoid contents were estimated. These extracts were assessed for an antioxidant property by 1, 1-diphenyl-2-picryl-hydroxyl (DPPH) method and reducing power assay was carried out using the FeCl<sub>3</sub> method. Inhibition of 15-lipoxygenase (LOX) by these extracts at 5 - 25µg to assess anti-inflammatory capacity was studied.

Antibacterial activity against some human pathogenic bacteria was tested by agar disk diffusion method. Among all the organic solvent extracts, methanolic extracts exhibited good antioxidant and antibacterial activity. Methanolic extract with an IC<sub>50</sub> value of 48mg for LOX inhibition is reported. The extracts showed inhibition of human pathogenic bacteria in the order: *Escherichia coli* > *Pseudomonas vulgaris* > *Klebsiella pneumonia*. Significant and positive linear correlations were found between total antioxidant capacities and phenolic contents indicating that phenolics were the dominant antioxidant constituents in tested seeds. Methanol extracts of *C. papaya* were subjected to LC-MS metabolite profiling. The LC-MS analysis identified 6 metabolites *p*-hydroxybenzoic acid, salicylic acid, hyperoxide, gental alcohol, triallyl glucose, kaempferolhexoside as the main constituents for the first time from this seed extract. Our study demonstrated that the selected papaya seeds have good antioxidant, anti-inflammatory, and antibacterial properties.

**Keywords:** *carica papaya* L., seed extracts, phytochemical analysis, antioxidant, lipoxygenase inhibition, antimicrobial activity.

## I. INTRODUCTION

Papaya (*Carica papaya* L.) is a member of the family Caricaceae. This plant family has four genera including Jarilla, Cylicomorpha, Cylicomorpha, and Carica. *Carica papaya* L. is common papaya and extensively grown over the world. The plant is herbaceous, soft tissue and fast-growing. Common names include papaya, papaw or pawpaw, Papeete (Pakistan), paper (French), Tenenbaum (German), chose (Spanish), mamao, mamoeiro (Portuguese), mugu (Chinese) and Malakal (Thailand). Papaya is a fruit plant with a soft stem, commonly and erroneously referred to as a tree. The papaya seeds contain balance-nutrients which consist of protein (24.3%), fatty oil (25.3%) and total carbohydrate (32.5 %). Although it contains a significantly high level of unsaturated fatty acids, papaya seeds seem not to be good oil seeds. Papaya seeds are used generally as an anti-parasitic agent by humans' plant is properly a large herb growing at the rate of 1.8-3 m in the first year and reaching 6-9 m in height[1]. The lower trunk is conspicuously scarred where leaves and fruit are borne. In some parts of the world, papaya leaves are made into tea as a treatment for malaria, dengue but the mechanism is to be scientifically proven. Papaya contains about 6% of the level of beta carotene. Excessive consumption of papaya may cause carotenemia, the yellowing of soles and palms[15]. Papaya releases a latex fluid when not ripe, possibly causing irritation and an allergic reaction in some people.

**Author α:** Division of Molecular Biology, Faculty of Life Sciences, JSS AHER, Mysuru, Dept of Nutrition & Dietetics, Faculty of Life Sciences, JSS AHER, Mysuru.

**Author σ:** Division of Molecular Biology, Faculty of Life Sciences, JSS AHER, Mysuru. e-mail: kantheshmb@jssuni.edu.in

**Corresponding Author ¥:** Assistant Professor, Division of Molecular, Biology, Faculty of Life Sciences, JSS Academy of Higher Education & Research, Mysore, Division of Molecular Biology, Faculty of Life Sciences, JSS AHER, Mysuru.

**Author ρ:** Institution of Excellence, Vijnana Bhavana, Mysuru.

**Author ω:** Division of Biotechnology, Faculty of Life Sciences, JSS AHER, Mysuru.

**Author §:** Assistant Professor, Division of Nutrition and Dietetics, School of Life Sciences, JSS AHER, SS NAGAR, Mysuru-15. e-mail: sushmabv@jssuni.edu.in



Figure 1: Papaya Fruits with Seeds

## YYP

### a) Chemicals and Reagents

Linoleic acid, 1, 1-diphenyl-2-picrylhydrazyl (DPPH), catechin were purchased from SIGMA ALDRICH (USA, MO). Sodium bicarbonate, 15-lipoxygenase, aluminum chloride, gallic acid, ascorbic acid, trichloroacetic acid (TCA), potassium ferricyanide, ferric chloride, folic-ciocalteu (FC) potassium buffer, borate buffer, nutrient agar, peptone, beef extract, hexane, chloroform, diethyl ether, methanol, borate salt, sodium dihydrogen phosphate, disodium hydrogen phosphate were laboratory chemicals.

### b) Processing of Plant Samples

*Carica papaya* fruits were collected from Mysore district, Karnataka, India. The pawpaw fruits were washed in tap water and then rinsed in sterile distilled water. The seeds were removed and shade-dried for about a week and were crushed using liquid nitrogen using mortar and pestle. Seeds were ground into a coarse powder. Fractionation of bioactive compounds was carried out using a solvent to increase the polarity of the solvents like hexane, chloroform, diethyl ether and methanol for 48 h in dark with constant stirring at room temperature. After each fractionation, the respective solvents were carefully filtered using a muslin cloth to prevent contamination by seed residue. The clear extract was air-dried to get a fine paste. The extract was weighed and stored at 4°C in dark until further analysis.

### c) Extraction Of Plant Material

About 25 g of coarsely powdered papaya seeds were weighed and suspended into 200mL of the solvent (hexane, chloroform, diethyl ether, and methanol) based on the increasing order of polarity. The extraction was

carried out at room temperature for 48 h using rotatory shaker at 30°C 60 rpm for 48 h. The extracts were first filtered with a clean muslin cloth and then suction filtered using flash operator at 44°C 160 rpm and finally dried it in glass Petri dishes at RT. The final drying process was carried out using by collecting the filtrates in the Eppendorf tube and dried in speed vacuum for 3 h at 40°C and the extracts were stored in dark at 4°C till further use.

## YYYP

## P

### Qualitative Analysis of Saponins and Tannins

#### a) A Test for Saponins

About 0.1 g of methanolic extract was diluted in 1ml of methanol. Extract (0.5 mL) was taken in a test tube and solubilized using 4.5mL of distilled water[2]. The formation of stable foam indicated the presence of Saponins.

#### b) A Test for Tannins

About 0.1 g of methanolic extract was diluted in 1ml of methanol. Extract (0.5 mL) was taken in a test tube and mixed with 10mL distilled water and ferric chloride reagent (3 drops,) added to the filtrate. A blue-black green precipitate confirmed the presence of Gallic tannins or catechol tannins[2].

#### c) Determination of Total Phenolics

The total phenolic content was estimated using the Folin-Ciocalteu (FC) calorimetric method. Gallic acid (20-100mg) standard was prepared. Extract (0.1 g) was weighed and diluted to make 100mg in 100mL. The extract (20-100mL) reacted with FC reagent (250mL) and was incubated at RT for 5min[3]. The reaction was neutralized with saturated sodium bicarbonate (1.5mL,

20%) that was added to the mixture and allowed to stand for 1 h[6]. The absorbance of the resulting blue color was measured at 765 nm (BECKMAN COULTER, DU 730 LIFE SCIENCE UV/VISIBLE SPECTROPHOTOMETER). Total phenolics content in the methanol extract of seeds was quantified by the calibration curve obtained from measuring the absorbance of known concentrations of gallic acid standard. The total phenolic contents were expressed as gallic acid equivalence (GAE) in  $\mu\text{g}$ .

#### d) Determination of Total Flavonoids

The total flavonoid content was determined by the aluminum chloride colorimetric method[5]. In brief, 10- 50 mL of extract were made up to 1mL with methanol, mixed with 4mL of distilled water and then 0.3mL of 5%  $\text{NaNO}_2$  solution.  $\text{AlCl}_3$  (0.3mL of 10%) solution was added after 5min of incubation and the mixture could stand for 6min. Then, 2mL of 1 mol/L NaOH solution was added, and the final volume of the mixture was brought to 10 mL with double-distilled water. The mixture could stand for 15min, and absorbance was measured at 510nm. The total flavonoid content was calculated from a calibration curve, and the result was expressed as  $\mu\text{g}$  catechin equivalent per g dry weight[9].

#### e) Antioxidant Activity

##### i. DPPH radical scavenging assay

The free radical scavenging property of the methanol extracts of papaya seeds was determined by the DPPH method. An aliquot of the extract was dissolved in a solvent and was plated out in duplicate in a 96- well microtiter plate. The DPPH radical solution (50mM; 2.9mg in 25mLmethanol) was added to alternating columns of the test samples and methanol was used as control. The percent of decolorization was recorded spectrophotometrically at 517nm using the Thermo Scientific Varioskan Flash Microtiter Plate Reader. The reaction for scavenging DPPH radical was in dark and the absorbance was recorded at 517 nm (Spectra Max, Molecular devices). Percent radical scavenging activity was determined by comparing with a solvent added as a control. The  $\text{IC}_{50}$  values were determined, which denote the Concentration of extracts required to scavenge 50% DPPH radicals[4]. Ascorbic acid (0.1 g in 5mL) was used as positive control at least three independent tests were performed for each sample. Solvent extracts of hexane, diethyl ether and methanol were tested[6]. Percent scavenging effect was determined by the following equation: % inhibition =  $[(\text{Absorbance of control} - \text{Absorbance of the test sample})/\text{Absorbance of control}] \times 100$ [7].

##### f) Reducing Power Assay

This estimation of reducing power was carried for papaya seeds with slight modifications. Test solution (0.1mL, 1mg/mL) was mixed with equal volume of

phosphate buffer (0.2M, pH 6.6) and potassium ferricyanide (2.5mL, 1%) and was incubated at 50°C for 20 min. Trichloroacetic acid (TCA; 10%, 2.5mL) was added to the mixture, which was then centrifuged at 3000 rpm for 5 min. After centrifugation, the supernatant solution (1.5mL) was taken in a test tube and was mixed with an equal volume of distilled water and ferric chloride (0.5mL, 0.1%). Ascorbic acid was used as a standard and phosphate buffer was used as a blank solution. Absorbance was measured at 700nm (Beckman-Colter, Du 730 Life Science Uv/Visible Spectrophotometer). Increased absorbance of the reaction mixture indicates stronger reducing power[8].

#### g) Anti-Inflammatory Activity

##### i. Lipoxygenase assay (LOX)

*Carica papaya* seeds were extracted and solubilized in methanol and tested for in vitro anti-inflammatory activity spectrophotometric assay for determination of LOX activity for papaya seed. Weight of empty Eppendorf tube was noted and 5mg of methanol extract was taken in empty Eppendorf tube extract was diluted using 1ml of methanol and shaken well. 15-LOX (5mg) activity with linolenic acid(0.2mM) in borate buffer (0.2M, pH 9.0) was carried out methanol extract (5, 10, 15, 20ml). The inhibition of LOX by the extracts was recorded by the time scan method at 234 nm[10]. Ascorbic acid inhibiting LOX was as recorded at 234nm using UV- Visible Spectrophotometer (Beckman-Coulter, Du 730 Life Science Uv/ Visible Spectrophotometer). The inhibitory effect of the extract was expressed as % of enzyme activity inhibition ( $\text{IC}_{50}$ ) value indicating the concentration required to inhibit 50% LOX activity[14]. It was calculated using the formula % of inhibition =  $[(\text{initial activity} - \text{inhibitor activity})/\text{initial activity}] \times 100$ .

#### h) Antibacterial Activity

##### i. Determination of Antibacterial Activity

Antibacterial activity of methanolic extracts (1, 2.5, 5 10  $\mu\text{L}$ ) of papaya seeds was determined by the disc diffusion method. The bacterial samples tested were *Escherichiacoli*, *Klebsiellapneumonia*, and *Pseudomonasvulgaris*. The media was prepared using peptone (3.75g), beef extract (2.25g), agar (15g) and distilled water (750mL). The contents were transferred to a flask and were plugged with cotton and wrapped using brown paper. Glass Petri plates were washed thoroughly rinsed with methanol and autoclaved at 121°C for 15min for complete sterilization[11]. The agar solution could cool, and 15 mL was poured into sterile glass Petri plates. The plates could set and then incubated at 37°C overnight. Colonies were picked from plates and used as inoculums of test organisms. The plates were incubated at 37°C overnight. Disc of Whatman No.1 filter paper was sterilized by heating in an oven for 30 min at 80°C[12]. Agar plates were inoculated with each organism and after 5 min, 6 filter paper discs,

impregnated with 5mL of the concentrated extracts, streptomycin (0.5mg/mL) were transferred onto the agar plates using sterile forceps. The plates were then incubated at 37°C overnight. The effectiveness of the extract as an antibiotic against the test organism was determined by measuring the diameter of the zone of inhibition.

## ii. LC-MS ANALYSIS

For the qualitative analysis of the metabolites were analyzed by Synapt G2 (UPLC separations with Quant of) according to the manufacturer's protocol. The nebulizer pressure was 60 psi and the nitrogen flow rate 10 L/min at a drying temperature of 350°C. The methanol seed extract was filtered (0.2-micron syringe filters, Millipore, U.S.A) and an aliquot (5 µl) was injected into the system. The mass spectra were acquired from m/z 100-1000 in negative ionization mode. Helium was used as the collision gas for the fragmentation of the isolated compounds in the ion trap[13]. The detection conditions were as follows: capillary voltage, 3500 V; skimmer voltage, -40 V; cap exit voltage, -158.5 V; Oct 1 DC, -12 V; Oct 2 DC, -2.45 V; trap drive level, 45.0; Oct RF, 150 Vpp; Lens 1, 5.0 V; Lens 2, 60 V.

## Y P

*Carica papaya* seed extract was prepared using four different solvents (hexane, chloroform, diethyl ether, and methanol) for the screening of bioactive capacity. The analysis was performed using a generally accepted

laboratory technique for qualitative determinations. Saponins test performed showed a positive result for hexane, diethyl ether and methanol whereas negative for chloroform. The tannins test conducted showed a positive result for hexane and methanol whereas negative results for chloroform and diethyl ether extracts. Thus methanol extract of *C.papaya* seeds contains saponins, tannin compounds. The importance of saponins and tannins in various antibiotics for treating common pathogenic strains has been reported[16].

**Table 1:** Phytochemical screening of methanolic extract of papaya seeds

Bioactives	Hexane	Chloroform	Diethyl Ether	Methanol
Tanins	+	-	-	+
Saponin	+	-	+	+


P

P

## Quantitative Analysis of Total Phenols and Flavonoids

### a) Total Phenolic Content

Total phenolic contents of the methanolic fractions of the seed of *C. papaya* were determined by using the Folin-Ciocalteu reagent and were expressed as gallic acid equivalents (GAE) per gram of seed extract. The total phenolic contents were 147µg for methanol extract of papaya seeds.



**Figure 2:** Total phenol content of 147µg gallic acid equivalent (GAE) was recorded using standard prepared by gallic acid

### b) Total Flavonoid Content

Flavonoids are secondary metabolites, the antioxidant activity of which is dependent on the presence of free -OH group, especially 3-OH. The total flavonoid content was 100mg for methanol extract of

papaya seeds. As this is the report on the antioxidant activity of *C. papaya* through phytochemical analysis, identification of the active phenolic and flavonoid compounds was attempted.






Figure 3: Total flavonoid content of 100 $\mu\text{g}$  and estimated GAE recorded using Standard prepared by catechin

### c) Antioxidant Activity

DPPH radical scavenging activity (%) can be calculated by using the formula as mentioned previously. Based on the results obtained, a graph also was prepared. The graph showed the percentage of radical scavenging activity of methanolic extracts at different concentrations with 50% of DPPH scavenging activity at 32mg.

Similarly, DPPH scavenging capacity for hexane, chloroform and diethyl ether for various extract concentration is reported. A comparison of the ability of various concentrations of seed-extract of hexane and diethyl ether with methanol extract indicated their limited DPPH scavenging capacity.




Figure 4: DPPH scavenging capacity of methanolic seed extract




Figure 5: DPPH scavenging capacity of hexane seed extract




Figure 6: DPPH scavenging capacity of diethyl ether seed extract

The reducing capacity of the papaya seed methanol extracts was compared to standard ascorbic acid. An increase in absorbance at 700 nm indicates the

reducing power of the extracts. The graph shows the concentration of methanol extract to scavenge 50% of added reducing chemical.




Figure 7: Reducing Power of Methanolic Seed Extract

## d) Anti-Inflammatory Activity

## i. Lipoxygenase (15-Lox) Inhibition

The methanol extracts of *Carica papaya* exhibited potent capacity inhibiting 50% LOX activity with an  $IC_{50}$  value of 47  $\mu$ g.





Figure 8: 15- LOX inhibition by methanolic extract of papayaseeds with  $IC_{50}$  of 47  $\mu$ g

## e) Antibacterial Screening

The results of the antibacterial sensitivity of the methanolic extract of *C. papaya* seed by disc diffusion method are depicted for different time intervals of 12, 24 and 48 h in the graph for *Escherichia coli* (Fig. 9) and *Pseudomonas Vulgaris* (Fig. 10). The results reveal that the extract has antimicrobial activity against these pathogenic organisms studied. The antibacterial

activity was screened from the zone of inhibition. The four different concentrations of methanolic extract (1, 2.5, 5 and 10 mg) inhibited *Escherichia coli* (Table 3; Fig. 11) *Pseudomonas Vulgaris* (Table 4; Fig. 12) growth with a maximum inhibition at 10mg. The streptomycin drug used showed maximum growth inhibition (3.94mm) compared to control (methanol, 10 ml). The drug inhibited *Escherichia coli* (2.95mm).








Figure 9: Results of methanolic seed extract inhibiting *E. coli* at different time intervals







Figure 10: Results of methanolic seed extract inhibiting *Pseudomonas vulgaris* at different time intervals

**Table 2:** Antimicrobial activity of papaya seed methanolic extract for *Escherichiacoli*

SI No.	Solvent Used	Concentration	Zone of Inhibition			
			T(0h)	T(12h)	T(24h)	T(48h)
1	Methanol	CONTROL	0.0	0.0mm	0.20mm	0.23mm
	Spteromycin	DRUG	0.0	0.45mm	2.63mm	2.95mm
		1 $\mu$ l	0.0	0.22mm	0.33mm	0.38mm
		2.5 $\mu$ l	0.0	0.4mm	0.48mm	0.53mm
		5 $\mu$ l	0.0	0.28mm	0.33mm	0.35mm
		10 $\mu$ l	0.0	1.35mm	1.42mm	2.86mm

**Table 3:** Antimicrobial activity of papaya seedmethanolic extract for *Pseudomonas vulgaris*

SI No.	Solvent Used	Concent Ration	Zone of Inhibition			
			T(0h)	T(12h)	T(24h)	T(48h)
1	Methanol	Control	0.0	0.96mm	1.43mm	1.65mm
	Streptomycin	Drug	0.0	3.26mm	3.27mm	3.94mm
		1 $\mu$ L	0.0	0.22mm	0.23mm	0.34mm
		2.5 $\mu$ l	0.0	0.0mm	0.34mm	0.36mm
		5 $\mu$ l	0.0	0.24mm	0.29mm	0.39mm
		10 $\mu$ l	0.0	0.7mm	1.56mm	1.67mm







Figure 11: Inhibition capacity of methanolic extracts for *Escherichiacoli* at concentration of 1 ug (c); 2.5 ug (d); 5 ug (e); and 10 ug (f) in comparison to control used methanol (a) and drug, streptomycin (b)















Figure 12: Inhibition capacity of methanolic extracts at concentration of 1 ug (c); 2.5 ug (d); 5 ug (e); and 10 ug (f) in comparison to control used (methanol; a) and drug, streptomycin (b)





- f) Profiling of Compounds in methanol extract of *Carica papaya* seeds determined by UPLC-PDA-ESI/HDMS  
Identification of metabolites based on chromatographic and in-house MS data identified 6 metabolites *p*-hydroxybenzoic acid, salicylic acid, hyperoside, gentisyl alcohol, trigalloyl glucose, kaemferolhexoside (Table 5) by comparison of retention time and MS/MS data.




Table 4: Putative compounds identified by UPLC-PDA-ESI/HDMS

M/Z Ratio	Molecules	Structures
137  138 462-463 407-408	<i>p</i> -hydroxy benzoic acid Salicylic acid Hyperoside Gentisyl alcohol Glucosyl and benzoyl groups intact	 <p>p-hydroxy benzoic acid</p>  <p>Salicylic acid</p>  <p>Hyperoside</p>
		 <p>gentisyl alcohol</p>






279,280,635, 115,116,117, 253,293,353, 359,277,278, 279,377,554, 642,297,265, 269,299,298, 357,358,393, 476,531,595, 688,281,311	Trigalloyl glucose  Luteolin 8-C(2-malonyl glucoside)  Kaempferol hexoside	<div>Trigallavl glucose</div>  <div>Kaempferol hexoside</div> 
353,406,547, 605,639,440, 995,831,738, 116,355,311, 368,402,833		


m/z	Molecules	Structures
253	Chrysin	 Chrysin
353	Chlorogenic acid	
355	Neochlorogenic acid	





	3-O-Caffeoylquinic acid	
	4-O- Caffeoylquinic acid	
	5-O- Caffeoylquinic acid	
269	Apigenin	


299	Hydroxybenzoic acid –O- hexoside	
358	Rosmarinic acid	
477	Isorhamnetin-3-O- hexoside	




595	Cyanidin-3-O- rutinoside	
136	Salicylic acid	
345	Epirosmanol	
461	Luteolin-3-O-glucoronide	
463	Quercetin-3-O- hexoside	


270	Naringenin	
-----	------------	---

354	1-Caffeoylquinic acid	
	3-Caffeoylquinic acid	
311	Caftaric acid	
355	Ferulic acid-0- hexoside	


611	Quercetin-3-rhamnosylglucoside	
-----	--------------------------------	--



P-MS1



P-MS2



P-MS3

### YP T

The constituent of the extract of *C. papaya* (dried) seeds contain compounds and micronutrients which may be responsible for its observed antioxidant activity. This study suggests that the plant possesses antioxidant activities that can counteract the oxidative damage. The total phenol test provides information on the reactivity of the seed extract with a stable free radical. It gives a strong absorption band. The degree of reduction in absorbance measurement is indicative of the radical scavenging (antioxidant) power of the extract. The extract of *Carica papaya* appeared to be as potent as Gallic Acid with maximum inhibition. The extract is found to have broad-spectrum antibacterial activity and used as analgesics and narcotics for pain relief. A report indicates that extracts are more active against Gram-positive bacteria than Gram-negative bacteria while that of the leaf extract

of *C. papaya* was next to the most sensitivity with Gram-negative bacteria[17]. The activity of the extract is comparable to those of antibiotics. The demonstration of activity against the test bacteria provides scientific bases for the local usage of the plant in the treatment of various ailments. The fact that the extract is active against Gram-positive bacteria and Fungi tested may indicate a broad spectrum of activity. This observation is very significant because of the possibility of developing therapeutic substances that will be active against multidrug-resistant organisms.

Lipoxygenases (LOXs) are a family of non-heme iron-containing dioxygenases catalyzing the biosynthesis of leukotrienes. Leukotrienes function as initiators of inflammation and their inhibition is partly responsible for the anti-inflammatory activity. In the present study methanolic extracts, *Carica papaya* showed good anti-LOX activity with an IC<sub>50</sub> value of

47 $\mu$ g.LOX inhibition was used to evaluate the anti-inflammatory activity of a few medicinal plants[10].

Plant phytochemicals with health benefits have been attributed to health as they cannot be synthesized by humans and they have been linked to antioxidant activity. In the present study, UPLC-DAD identified *p*-hydroxybenzoic acid, salicylic acid, hyperoside, gentisyl alcohol, trigalloyl glucose, kaemferolhexoside among others. These are reported as the strongest natural anti-inflammatory agent[13]. The presence of the phytochemicals in the extract could also support the therapeutic property tamarind seed for the mentioned application in the traditional literature of India.

## YP S

*Carica papaya* is a nutraceutical plant having a wide range of pharmacological activities. The whole plant has its own medicinal value. The wide range of enzymes, vitamins present in *Carica papaya* makes it a nutraceutical plant. Antioxidant and antimicrobial properties of methanolic extract of *Carica papaya* have recently been of great interest in both the research and food industry, because of its possible use as natural additives which emerged from a growing tendency to replace synthetic antioxidants with natural ones. Owing to the antioxidant and antibacterial activities exhibited by the seed extract investigated in this study, it could be considered a natural herbal source that can be used in the food and pharmaceutical industries. However, further studies are needed to obtain purified compounds that may be responsible for the activities observed from the tested seeds.

1. Irfan Muhammad Haseeb Tariq, Bilal Ghaffar, Tauqeer Ahmed, Abbas Sultan, Muhammad, Muhammad Junaid Farrukh. (2015) Phytochemical and Microbiological Evaluation of Different Chemical Extracts of Papaya Seeds on Clinical Isolates. *Int j pharm.* 5(1): 122-126.
2. Yusha'u, Onuorah F C. and Murtala Y. (2009) In-Vitro Sensitivity Pattern of Some Urinary Tract Isolates to *Carica papaya* extracts. *Bayero Journal of Pure and Applied Sciences.* 2(2): 75 – 78.
3. Delphin D V, Haripriya R, Subi S, Jothi D and P Thirumal Vasan. Phytochemical Screening of Various Ethanolic Seed Extracts. *Research Publications in Ayurvedic Sciences.* 7(3): 1041-1048.
4. Sekhar Shailasree, Karmakar Ruma, Harishchandra S. Prakash. (2013) Curative properties of *Buchananialanzan*: As evaluated by its antioxidant, anti-inflammatory and DN protective properties. *Jnatpharm* IP: 14.139.155.135.
5. Lisda Hayatie, Agung Biworo, and Eko Suhartono. (2015) Aqueous Extracts of Seed and Peel of *Carica* Papaya Against *Aedes Aegypti*. *Journal of Medical and Bioengineering.* 4(4): ISSN: 2315-442.
6. Chethan j, Sampath kumara k k, Shailasreeseekhar, Prakash H.S. (2012) Antioxidant, Antibacterial and DNA Protecting Activity of Selected Medicinally Important Asteraceae Plants. *Int J Pharm PharmSci.* ISSN-0975-1491.
7. N. Raaman. (2015) Thin layer chromatographic analysis and Anioxidant Activities of Methanol Extract of Leaves of *C.papaya* L. *IJAPBC.* 4(2):ISSN: 2277 – 4688.
8. Aysun OZKAN, Hamide GUBBUK, Esma GUNES, Ayse ERDOGAN. (2009) Antioxidant capacity of juice from different papaya cultivars grown under greenhouse conditions in Turkey. *Turkish Journal of Biology.* 35(5):619-625
9. Kaibing Zhou, Hui Wang, Wenli Meri, Xiaona Li, Ying Luo and Haofu Dai. (2011) Antioxidant Activity of Papaya Seed Extracts. *Molecules.* 16(8): 6179– 6192.
10. Shailasreeseekhar, Sampath-kumara kk, Niranjana s r and Prakash H S. (2013) In Vitro Antioxidant Activity, Lipoxxygenase Cyclooxygenase-2 Inhibition and DNA Protection Properties of memecyl- onspecies. *International Journal of Pharmacy and Pharmaceutical Sciences.* 5(2):257-262.
11. E.i.okoye. (2011) Preliminary phytochemical analysis and antimicrobial Activity of seeds of *Carica papaya*. *American Journal of Pharmacology and Pharmacotherapeutics.* ISSN :2393-8862
12. Augustine Ocloo, Nonye Constance Nwokolo and Nicholas T. K. D. Dayie. (2012) Phytochemical Characterization and Comparative efficacies of crude extracts of *carica papaya*. *Int. J. Drug Res. Tech.* 2(5):399-406.
13. Tumkurramasettybharathi, Shailasreeseekhar, Kiggakadappasampath kumara, Mudalabeeduchandregowda, Madhusudhanand Harishchandra sripathy Prakash. (2016) Metabolite profiling by UPLC-PDA-esi/hdms and antibacterial activity of *memecylontalbotianum*. *Pharmacogn. Commn.* 6(4): 225-231.
14. Sekhar shailasree, K. K. Sampathkumara, S. R. Niranjana and H. S. Prakash. (2014) Bioactive Potential of Medicinal Plants from Western Ghats Region, India. *Journal of Herbs, Spices & Medicinal Plants.* 20:221–234.
15. Onkar Singh and M. Ali (2011) Phytochemical and Antifungal Profiles of the Seeds of *Carica Papaya* L. *Indian J Pharm Sci.* 73(4): 447–451.
16. Kubmarawa D, Ajoku G A. Enworem N M. and Okorie D A. (2007) Preliminary phytochemical and antimicrobial screening of 50 medicinal plants from Nigeria. *African Journal of Biotechnology* 6: 1690-1696.
17. Jigna and Sumitra. (2007). Antibacterial activity and phytochemical screening of some medicinal plants



commonly used in Saudi Arabia against selected pathogenic microorganisms. *Journal of King Saud University-Sciences*.25(2):115–120.

18. Lohiya N K, Goyal R B, Jayaprakash D, Ansari AS, Sharma S.(1994) Antifertility effects of aqueous extract of *Carica papaya* seeds in male rats. *Planta Med*.60(5):400-4. [PubMed: 7997464].
19. Udoh P, Essien I, Udoh F.(2005) Effect of *Carica papaya* (paw paw) seeds extract on the morphology of pituitary-gonadal axis of male Wistar rats. *Phytother Res*.19:1065–8. [PubMed: 16372375].
20. Verma R J, Nambiar D, Chinoy N J.(2006) Toxicological effects of *Carica papaya* seeds extract on spermatozoa of mice. *J Appl Toxicol*. 26:533–5. [PubMed: 17080407].
21. V. Boshra and A. Y. Tajul. (2013) "Papaya - an innovative raw material for food and pharmaceutical processing industry," *Health and the Environment Journal*. (4): ISSN 0936-6768.
22. Rezaeizadeh, A, Zuki A. B. Z., Abdollahi, M Goh, Y. M, Noordin, M. M, Hamid M, &Azmi T. I. (2011) Determination of antioxidant activity in methanolic and chloroformic extracts of *Momordica charantia*. *African Journal of Biotechnology*. 10(24):4932–4940.
23. Boning, Charles R. (2006) Florida's Best Fruiting Plants: Native and Exotic Trees, Shrubs, and Vines. Sarasota, Florida Pineapple Press. Inc. 166–167.
24. K. Kovendan, K. Murugan, C. Panneerselvam, N. Aarthi. (2012) Antimalarial activity of carica papaya leaf extract against plasmodium falciparum. *Asian Pac J. Trop Dis*. Inc. S306-S311.
25. Trease G. E and Evans W. C. (1989) Pharmacology, 11th edn., Bailliere Tindall, London, 45-50.
26. Akah P. A, Oli A. N, Enwerem N. M. and Gamaniel K. (1997) Preliminary studies on purgative effect of *Carica papaya* root extract. *Fitoterapia*. 68 (4): 327-331.
27. Ayoola. (2008) Phytochemical Screening and Antioxidant Activities of Some Selected Medicinal Plants. *Journal of Pharmaceutical Research*. 7 (3): 1019-1024.

