

# Neglected Trans-Scaphoid Volar Perilunate Dislocation with Post-Operative Complex Regional Pain Syndrome: A Case Report

Debasish Meher

*Received: 12 December 2019 Accepted: 5 January 2020 Published: 15 January 2020*

## Abstract

We present the case report of a 30-year-old male who fell from a height of 15 feet on his palmar-flexed right wrist and came to us 1 month after the injury. He had a swollen, stiff wrist with painful movements and no neurovascular deficit. X-rays showed a trans-scaphoid volar perilunate dislocation. We used the volar approach for open reduction and internal fixation with a headless screw for scaphoid fracture and scapho-lunate and capito-lunate K-wires for intercarpal instability. The wrist was immobilised in a below-elbow POP slab for 6 weeks after which the K-wires were removed. The patient unfortunately developed complex regional pain syndrome (CRPS), which can be attributed to his late presentation after the injury. Active and passive range of motion exercises and contrast bath were initiated and he was kept on low dose amitriptyline. Gradually CRPS resolved and at 6-month follow-up, the patient had a wrist dorsiflexion of 30° and palmar-flexion of 45°.

## Index terms—

## 1 Introduction

olar perilunate dislocation (VPLD) is a rare injury accounting for about 3% of perilunate dislocations which comprise of less than 10% of all wrist injuries. It generally occurs due to high-energy trauma and is frequently missed. Open reduction and internal fixation is the ideal treatment due to variable results with closed reduction. Patients presenting late are particularly difficult to manage due to soft tissue contractures and increased chances of 1) avascular necrosis of scaphoid and lunate, 2) tractional injury to neurovascular structures, 3) wrist stiffness, and 4) complex regional pain syndrome (CRPS). We, through this case report, aim to highlight the problems faced in managing neglected VPLDs and importance of prompt restoration of intercarpal alignment in such injuries to prevent complications. We also state that early recognition and treatment improves patient outcomes in CRPS.

## 2 II.

## 3 Case Presentation

A 30-year-old man presented to us with pain in the right wrist and restricted movements. He had fell down from a height of around 15 feet on his palmarflexed wrist a month back. He took the primary treatment from an osteopath where some kind of bandaging was done. He came to us when his symptoms hadn't After taking an informed consent, we took the patient to the operating table. We used the volar approach to the scaphoid. Intra-operatively, we found a comminuted fracture of the scaphoid and gross intercarpal instability. We did open reduction and internal fixation with a headless screw for scaphoid fracture and scapho-lunate and capito-lunate K-wires for intercarpal instability.

The patient was kept on a below-elbow plaster slab for 6 weeks, after which it was removed along with the K-wires and range of motion exercises were started. Unfortunately, the patient developed complex regional pain

syndrome (CRPS) at this point, which was suggested by his complaints of severe burning pain and abnormally increased sweating in the affected area. Xrays showed severe patchy osteopenia around the wrist. Active and passive range of motion exercises and contrast bath were initiated and he was kept on low dose amitriptyline. Gradually CRPS resolved and at 6month follow-up, the patient had a wrist dorsiflexion of 30° and palmar-flexion of 45°. (Figure 2)

## 4 Discussion

Perilunate dislocation comprises volar or dorsal translation of the carpus secondary to ligamentous or bony disruption around the lunate. This was explained by Johnson as greater arc (associated with a fracture around lunate) and lesser arc (pure ligamentous disruption around lunate) injuries. Trans-scaphoid volar perilunate dislocation (VPLD) is an extremely rare injury, generally caused by a fall on palmar-flexed and ulnar-deviated wrist. It is frequently missed and hence, can be neglected. Patients can present with persistent pain and stiffness of the wrist joint with signs of median nerve compression.

Prompt restoration of intercarpal alignment is necessary to prevent complications like median nerve injury, carpal instability, complex regional pain syndrome, avascular necrosis of lunate and scaphoid, and secondary osteoarthritis. The literature suggests that an open reduction is necessary and can be done till 2 months after the injury. In patients where open reduction is not successful or those presenting after 2 months, a reconstructive or salvage procedure is generally required. Patients presenting late with VPLDs have more chances of developing complex regional pain syndrome (CRPS) in the post-operative period. CRPS occurs due to central and peripheral nociceptive sensitisation, altered sympathetic function, inflammatory and immune related factors, brain changes, and psychological factors. It is characterised by excessive burning pain, swelling, altered sweating pattern and warm or cold skin. Diagnosis is solely clinical. Early recognition and treatment is necessary. Various treatment modalities like drugs (corticosteroids, anti-convulsants, analgesic antidepressants), ganglion blocks, spinal cord stimulation and physical therapy have been documented.



Figure 1: Figure 1 :



Figure 2: Figure 2 :



---

65 [An Update on the Pathophysiology of Complex Regional Pain Syndrome Anesthesiology ()] ‘An Update on the  
66 Pathophysiology of Complex Regional Pain Syndrome’. *Anesthesiology* 2010. 113 (3) p. .

67 [Inoue and Shinova ()] ‘Late treatment of unreduced perilunate dislocation’. G Inoue , K Shinova . *J Hand Surg*  
68 *Br* 1999. Bruehl S. 24 p. .

69 [Kammano et al. ()] ‘Palmar lunate transscaphoid fracture dislocation caused by a palmar flexion injury’. M  
70 Kammano , Y Honda , K Kazuki . *J Orthop Trauma* 2001. 15 p. .

71 [Johnson ()] ‘The acutely injured wrist and its residuals’. R P Johnson . *Clin Orthop Relat Res* 1980. (149) p. .

72 [Howard and Dell (1986)] ‘The unreduced carpal dislocation. A method of treatment’. F M Howard , P C Dell .  
73 *Clin Orthop Relat Res* 1986 Jan;(202. p. .

74 [Adkison and Chapman ()] ‘Treatment of acute lunate and perilunate dislocations’. J W Adkison , M W  
75 Chapman . *Clin Orthop Relat Res* 1982. 164 p. .

76 [Lal et al. ()] ‘Two stage procedure for neglected transscaphoid perilunate dislocation’. H Lal , V Jangira , R  
77 Kakran , D Mittal . *Indian J Orthop* 2012. 46 p. .

78 [Youssef and Deshmukh (2008)] ‘Volar perilunate dislocation: a case report and review of the literature’. B  
79 Youssef , S C Deshmukh . *Open Orthop J* 2008 Apr 11. 2 p. .