

1 Atoui Hadi¹, El Haddad Cynthia², Barakat Habib³ and Darido Jessie⁴

2 ¹ Holy Spirit University of Kaslik, Lebanon

3 Received: 7 December 2019 Accepted: 1 January 2020 Published: 15 January 2020

4

5 **Abstract**

6 Renal angiomyolipoma (AML) is the most common benign tumor of the kidney. There are few
7 case reports in the literature, especially those occurring during pregnancy. We, at this moment,
8 are reporting a case of a 32-year-old female patient who presented at 21 weeks of gestation
9 with right-sided flank pain, chills, macroscopic hematuria, and vomiting. On examination, she
10 was hemodynamically stable, with no fever. Renal ultrasound showed the presence of a
11 hyperechogenic vascularized fatty tissue on the right kidney, measuring 7.4 x 5.1 x 6.2 cms,
12 with minimal pelvicalyceal dilatation. The MRI opted for an angiomyolipoma. Discharged
13 home at day 4 of admission, the patient's continued the remaining weeks of her pregnancy
14 uneventfully, until 37 weeks. She delivered her baby vaginally with no further complications
15 during pregnancy or in the post-partum period. In conclusion, due to the insufficient data in
16 the literature supporting the management of patients with AML, the individualization of the
17 treatment is an essential strategy.

18

19 **Index terms**— ?angiomyolipoma? ?renal tumor? ?pregnancy? ?surgery? ?embolization?.

20 **1 Renal Angiomyolipoma during Pregnancy:**

21 A Case Report Her medical, surgical, and obstetrical history consisted of kidney stones, one vaginal delivery, and
22 one dilatation and curettage for incomplete abortion. She was on acetylsalicylic acid (ASA) during her current
23 pregnancy.

24 On examination, she was hemodynamically stable, with no fever. A blood test was ordered and revealed,
25 hemoglobin level at 11.5 (Hematocrit 33.7), White Blood Count (WBC) at 11.9, CRP 3.92, Creatinine 0.48.
26 Urine analysis showed red blood cells at 80 at WBC at 5. Hepatic panel and electrolytes were within normal
27 levels.

28 Abdominal examination revealed tenderness on the right groin. The urologist and infectious disease specialists
29 were also in this case. Her pain was relieved by intravenous analgesics and relative bed rest.

30 The obstetrical ultrasound showed a single intrauterine pregnancy with positive cardiac activity commensurate
31 with the gestational age; however, the renal ultrasound showed the presence of a hyperechogenic vascularized
32 fatty tissue on the right kidney, measuring 7.4 x 5.1 x 6.2 cms, with minimal pelvicalyceal dilatation. There is
33 no lithiasis or subcapsular hematoma (Image 1). A renal MRI completed the investigations.

34 Image 1: Ultrasound of the right kidney, as described above

35 The MRI result showed a well-defined, 75mm, multilocular renal mass occupying the middle segment of
36 the right kidney with an image of a small pelvicalyceal dilatation and an intracavitory hemorrhagic content.
37 Consequently, the MRI report evoked the diagnosis of angiomyolipoma. (Image 2, 3, 4)1 Year 2020 Introduction
38 dilatation.

39 The MRI opted for an angiomyolipoma. Discharged home at day 4 of admission, the patient's continued the
40 remaining weeks of her pregnancy uneventfully, until 37 weeks. She delivered her baby vaginally with no further
41 complications during pregnancy or in the postpartum period.

42 In conclusion, due to the insufficient data in the literature supporting the management of patients with AML,
43 the individualization of the treatment is an essential strategy.

44 **Keywords:** "angiomyolipoma" "renal tumor" "pregnancy" "surgery" "embolization". renal angiomyolipoma
45 (AML) is the most common benign tumor of the kidney. It appears mainly in females during their procreation

7 CONCLUSION

46 age and is affected by the hormonal changes occurring during pregnancy. It could be life-threatening when
47 ruptured, leading to severe bleeding.

48 There are few cases in the literature concerning the optimal management taken in the case of AML, especially
49 during pregnancy.

50 We, at this moment, are going to describe the evolution of AML during pregnancy in a 32 years old female,
51 trying to maintain a normal renal function and a viable fetus until delivery.

52 2 II.

53 3 Case Presentation

54 It is the case of a 32 years-old female who presented at 21 weeks of gestation. She had a one-day history right-
55 sided flank pain, chills, macroscopic hematuria, and vomiting. On day 4 of admission, the patient's condition
56 remained stable, with no fever and less pain. Therefore, she was discharged on analgesics with a medical report
57 of her state, so she can rest at home. The remaining weeks of her pregnancy were completely uneventful, and
58 the patient delivered her baby vaginally at 37 weeks with an APGAR of 9/10, weighting 2500 g. There were no
59 further complications during the pregnancy or in the post-partum period.

60 4 III.

61 5 Discussion

62 Angiomyolipoma is the most common benign mesenchymal tumor of the kidney, composed of adipose and vascular
63 tissue in the association of smooth muscle.

64 Its prevalence varies between 0.12 and 0.14 percent in the general population. There is also a female
65 predominance with a ratio of 4:1. Most of the time, it is the Right kidney that is affected [1].

66 The AML could appear either sporadically or in association with tuberous sclerosis. In the first case, AML
67 is often solitary and accounts for 80% of the AML. Generally, patients present with a mean age of 43 years old.
68 On the other hand, in 20% of the cases, AML is associated with tuberous sclerosis. In the latter case, the mean
69 age at the time of diagnosis is 25 to 35 years. The lesions typically exceeds the isolated angiomyolipoma in size,
70 and they are often bilateral and multiple. Angiomyolipoma occurs in 80% of patients with tuberous sclerosis.
71 Exceptionally, these renal tumors could rupture, leading to massive retroperitoneal hemorrhage and resulting in
72 what we call the Wunderlich syndrome [2].

73 The classical clinical presentation of AML is flank pain, palpable mass, nausea, hematuria, and anemia. AML
74 tends to appear during the pregnancy period, due to the hormonal influence of estrogen and progesterone in
75 addition to the increased receptors on the surface of the AML associated with the expansion of the intraabdominal
76 pressure during gestation.

77 The sonographic features of AML consist of a well-circumscribed and highly echogenic mass because of its
78 high-fat content, multiple nonfatty interfaces, heterogeneous cellular architecture, and numerous vessels. Other
79 renal tumors, such as lipoma, teratoma, Wilms tumor, oncocytoma, and renal cell carcinoma (RCC), may contain
80 fat and can be difficult to be differentiated on imaging studies. A CT scan with thin (less than 5-mm) sectioning is
81 recommended for the confirmation of diagnosis whenever AMLs are suspected [3]. However, MRI does not appear
82 to have an advantage over CT scan, except in pregnancy, and when the intravenous contrast administration is
83 not indicated.

84 In the case of rupture, hemodynamic stability is of critical importance for the selection of an optimal
85 treatment strategy. In the case of hemodynamically unstable patients, emergent surgery (nephrectomy) or Renal
86 Angiomyolipoma during Pregnancy: A Case Report arterial embolization (if available) are the main options of
87 treatment [4]. The Embolization consists of an alternative after 12 weeks of gestation with minimal fetal radiation
88 exposure. Concerning the asymptomatic pregnant patients, the conservative approach may be of choice in these
89 cases [5]. As for the definitive treatment, it may be delayed after the delivery. According to the literature, most
90 of the patients with renal angiomyolipoma, delivered their babies via cesarean section (56%), whereas only 19%
91 delivered vaginally (Table 1). However, vaginal delivery is a safe approach for these patients, and the cesarean
92 does not affect the risk of rupture. Consequently, the mode of delivery should be decided based on obstetrical
93 indications only. Vacuum extraction can also be an alternative in order to reduce the time of the second stage of
94 labor.

95 Seeing that our patient was hemodynamically stable, along with the normal development of her fetus, a
96 multidisciplinary approach decided that the patient proceeds to the term of delivery. She underwent a successful
97 vaginal delivery without complications.

98 6 IV.

99 7 Conclusion

100 Due to the insufficient data in the literature supporting the management of patients with AML, the individual-
101 ization of the treatment is an essential strategy.

102 We need to have more experience with these strategies and to initiate more studies, so it can be the basis of
103 any recommendation for the optimal treatment method.

104 **8 Conflict of Interest**

105 No conflict of interest to declare

106 **9 Consent and Ethical Approval**

107 Obtained from the patient to publish the case.

108 **10 Financial Funding**

No funding was obtained for this publication. ¹



Figure 1:

109

¹© 2020 Global JournalsRenal Angiomyolipoma during Pregnancy: A Case Report

1

Author	Year	Maternal Age	GW	Tumor size (cm)	Ruptur	RA Management	Pregnancy Man- agement
Lee [6]	1994	29	27	NR	Yes	Nephrectomy	Fetal death
Yanai [7]	1996	NR	39	NR	Yes	Embolization	Term delivery
Oka [8]	1999	32	36	NR	Yes	Nephrectomy	Term C/S
Tanaka [9]	2001	23	27	7	Yes	Conservative Embolization	+Later Vaginal delivery
Gimeno Argente [10]	2006	40	33	NR	Yes	Nephrectomy	CS
Raft [11]	2006	40	34	NR	Yes	conservative	Preterm C/S
Storm [1]	2006	32	39	8	Yes	Conservative	Vaginal delivery
Koh [12]	2007	31	12	NR	Yes	Nephrectomy	Term C/S
Illescas Molina [13]	2007	36	28	NR	Yes	conservative	Term C/S
Kontos [14]	2008	28	33	7	Yes	Nephrectomy	PreTerm C/S
Binkowska [15]	2009	26	20	17,4	Yes	Embolization	Term C/S
Komeya [16]	2010	39	38	8	Yes	Embolization	Term C/S
Zapardiel [17]	2011	30	35	11	Yes	Embolization	PreTerm C/S
Gyimadu [18]	2011	21	25	11,5	Yes	Conservative Embolization	+Later Term C/S
Lopater [19]	2011	34	30	4	No	Nephrectomy	Term C/S
Govendik Horny [20]	2011	30	20	8	No	Nephrectomy	NR
Bolufer [21]	2012	26	NR	12	Yes	Nephrectomy	Vaginal delivery
Ferianec [22]	2013	30	9	21	Yes	Nephrectomy	Therapeutic Abortion
Iruloh [23]	2013	23	31	12	Yes	Embolization +Nephrectomy	Term C/S
Pontis [24]	2013	33	34	4,8	Yes	Nephrectomy	Preterm C/S
Davis [25]	2013	NR	NR	NR	NR	NR	NR
dos Santos [26]	2014	40	18	5	Yes	Conservative	Preterm C/S
Preece [27]	2015	45	24	15	Yes	Embolization +Nephrectomy	Term C/S
Bidault [28]	2015	31	NR	9	No	Nephrectomy	-
Cetin [29]	2015	26	44	11	No	Nephrectomy	Vaginal delivery

[Note: NR: Not Reported; GW: Gestational week; RA: Renal Angiomyolipoma; C/S: Caesarean section.]

Figure 2: Table 1 :

110 [Argente ()] , Gimeno Argente . 2006.

111 [Molina ()] , Illescas Molina , T . *Ginecol Obstet Mex* 2009. 77 p. .

112 [Cetin ()] , C Cetin . *J obstet Gynecol* 2015. 12 (02) p. .

113 [Lee ()] ‘angiomyolipoma of the left uterovesical junction’. J D Lee . *J Reprod Med* 1994.

114 [Govednik-Horny and Atkins ()] ‘Angiomyolipoma with vascular invasion during pregnancy’. C Govednik-Horny
115 , M Atkins . *Ann Vasc Surg* 2011. 25 p. 1138.

116 [Tanaka (2001)] ‘conservative management and vaginal delivery following ruptured renal angiomyolipoma’. M
117 Tanaka . *Obstet Gynecol* 2001 Nov. 98 p. .

118 [Storm and Mowad ()] ‘Conservative management of a bleeding renal angiomyolipoma in pregnancy’. D W Storm
119 , J J Mowad . *Obstet Gynecol* 2006. 107 p. .

120 [Gyimadu ()] ‘conservative management of a retroperitoneal hemorrhage following a ruptured renal angioliopoma
121 in pregnancy’. A O Gyimadu . *J obstet Gynaecol res* 2011. 37 p. .

122 [Binkowska ()] ‘embolization of renal angiomyolipoma in pregnancy: case’. M Binkowska . *report ginekol pol*
123 2009. 80 p. .

124 [Komeya M et al rupture of renal angiomyolipoma during pregnancy: a case report Hinyokika Kiyo ()]
125 ‘Komeya M et al rupture of renal angiomyolipoma during pregnancy: a case report’. *Hinyokika Kiyo*
126 2010. 56 p. .

127 [Kontos S et al rupture of renal angiomyolipoma during pregnancy: case report Cases J ()] ‘Kontos S et al rup-
128 ture of renal angiomyolipoma during pregnancy: case report Cases’. *J* 2008. 1 p. 245.

129 [Lopater et al. ()] ‘Management of angiomyolipoma vena cava thrombus during pregnancy’. J Lopater , O
130 Hartung , F Bretelle , C Bastide . *Obstet Gynecol* 2011. 117 p. .

131 [Zapardiel ()] ‘renal angioliopoma during pregnancy: review of the literature’. I Zapardiel . *Gynecol Obstet invest*
132 2011. 72 p. .

133 [Iruloh et al. ()] ‘Renal angiomyolipoma and lymphangioleiomyomatosis in pregnancy’. C Iruloh , R Keriakos ,
134 D J Smith , T Cleveland . *J Obstet Gynaecol* 2013. 33 p. .

135 [Raft (2006)] ‘renal angiomyolipoma rupture during pregnancy’. J Raft . *Gynecol Obstet Fertil* 2006 Oct. 34 (10)
136 p. .

137 [Bidault et al. ()] ‘Renal angiomyolipoma with inferior vena cava thrombosis during pregnancy’. V Bidault , G
138 Pignot , L Rocher , L Glas , J J Patard . *Prog Urol* 2015. 25 p. .

139 [Bolufer et al. ()] ‘Robot assisted partial nephrectomy (Da Vinci) in an angiomyolipoma associated to Wünder-
140 lich Syndrome’. E Bolufer , G Lopez-Fontana , O A Castillo . *Arch Esp Urol* 2012. 65 p. .

141 [Oka (1999)] ‘Rupture of a renal angiomyolipoma in regnancy: a case report’. D Oka . *J Gland Biol Neoplasia*
142 1999 Oct. 4 (4) p. .

143 [Pontis et al. ()] ‘Rupture of renal angiomyolipoma in pregnancy’. A Pontis , B Piras , A Meloni , A De Lisa , G
144 B Melis , S Angioni . *J Obstet Gynaecol* 2013. 33 p. .

145 [Davis et al. ()] ‘Selective arterial embolisation of bilateral angiomyolipomata in a symptomatic pregnant female’.
146 N F Davis , R Kelly , M J Lee , P Mohan . *BMJ Case Rep* 2013. 2013.

147 [Shah et al. ()] ‘Selective Embolization of Bleeding Renal Angiomyolipoma in Pregnancy’. J Shah , J Jones , M
148 A W Miller . *Journal of the Royal Society of Medicine* 1999. 92 p. .

149 [Ferianec et al. ()] ‘Severe retroperitoneal haemorrhage in the first trimester of a multiple pregnancy after
150 spontaneous rupture of renal angiomyolipoma’. V Ferianec , M Gabor , M Cano , P Papcun , K Holoman .
151 *Arch Gynecol Obstet* 2013. 288 p. .

152 [Koh (2007)] ‘simultaneous cesarean section and radical nephrectomy for angiomyolipoma with spontaneous
153 bleeding during pregnancy: A case report’. J L Koh . *J reprod Med* 2007 Apr. 52 (4) p. .

154 [Yanai ()] ‘Spontaneous hemorrhage during pregnancy secondary to renal angiomyolipoma’. H Yanai . *Urol Int*
155 1996. 56 (3) p. .

156 [Santos et al. ()] ‘Spontaneous rupture of renal angiomyolipoma during pregnancy’. Dos Santos , M M Proenca ,
157 S M Reis , M I Viana , R M Martins , L M Colaço , J M Nunes , FM . *Rev Bras Ginecol Obstet* 2014. 36 p. .

158 [Preece et al. ()] ‘Surgical management of haemorrhaging renal angiomyolipoma in pregnancy’. P Preece , B Mees
159 , B Norris , M Christie , T Wagner , P Dundee . *Int J Surg Case Rep* 2015. 7 p. .

160 [Preece et al. ()] ‘Surgical Management of hemorrhaging renal angiomyolipoma in pregnancy’. P Preece , B Mees
161 , B Norris , M Christie , T Wagner , P Dundee . *Int J Surg Case Rep* 2015. 7 p. .

162 [Halpenny et al. ()] ‘The radiological diagnosis and treatment of renal angiomyolipoma-current status’. D
163 Halpenny , A Snow , G Mcneill , W C Torreggiani . *Clin. Radiol* 2010. 65 (2) p. .

164 [Marino et al. ()] *Wunderlich syndrome. Clinical and therapeutic aspects of a long-term experience*, G Marino ,
165 M Pedalino , O G Di Primio , D Piras , R Vella , E Verces . 2010. *Urologia3*. p. .