



GLOBAL JOURNAL OF MEDICAL RESEARCH: E
GYNECOLOGY AND OBSTETRICS
Volume 20 Issue 3 Version 1.0 Year 2020
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Knowledge, Attitudes and Acceptability of Vaccination against Papillomaviruses: A Study on 76 Residents in Gynaecology and Obstetrics in Dakar (Senegal)

By Taliana Stéphanie Gondjout, Omar Gassama, Mouhamed Diadhiou,
Djibril Diallo, Philippe Marc Moreira, Alassane Diouf,
Marieme Gueye Ba & Jean Charles Moreau

Cheikh Anta Diop University

Abstract- The aim of this work was to know the level of knowledge Attitudes and Acceptability of resident in Gynaecology and Obstetrics Knowledge of Vaccination against Papillomaviruses. It was a descriptive prospective cohort study, from March 12 to July 30th, 2019. The study involved 76 residents. The studied parameters included Socio-epidemiological factors including age, education level, occupation, marital status, gynecological and obstetrical history, knowledge of human papillomavirus and knowledge and acceptability of vaccination against human papillomavirus and acceptability oh papillomavirus vaccine. The data has been collected by excel, and the statistical analysis has been performed using Epi-info 7. In this study, collected 76 residents. The mean age of the residents was 30.7 years. Residents were predominantly married (54.2%). The average pregnancy was 0.8. The average age at first intercourse was 27.6. All residents (100%) knew about HPV. Lessons (96.1%) were the principal sources of information.

Keywords: cervical cancer, human papillomaviruses vaccine, acceptability, knowledge-e, Senegal.

GJMR-E Classification: NLMC Code: QW 165.5.P2



Strictly as per the compliance and regulations of:



© 2020. Taliana Stéphanie Gondjout, Omar Gassama, Mouhamed Diadhiou, Djibril Diallo, Philippe Marc Moreira, Alassane Diouf, Marieme Gueye Ba & Jean Charles Moreau. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Knowledge, Attitudes and Acceptability of Vaccination against Papillomaviruses: A Study on 76 Residents in Gynaecology and Obstetrics in Dakar (Senegal)

Taliana Stéphanie Gondjout ^α, Omar Gassama ^σ, Mouhamed Diadhiou ^ρ, Djibril Diallo ^ω, Philippe Marc Moreira [¥], Alassane Diouf [§], Marieme Gueye Ba ^x & Jean Charles Moreau ^v

Abstract- The aim of this work was to know the level of knowledge Attitudes and Acceptability of resident in Gynaecology and Obstetrics Knowledge of Vaccination against Papillomaviruses. It was a descriptive prospective cohort study, from March 12 to July 30th, 2019. The study involved 76 residents. The studied parameters included Socio-epidemiological factors including age, education level, occupation, marital status, gynecological and obstetrical history, knowledge of human papillomavirus and knowledge and acceptability of vaccination against human papillomavirus and acceptability oh papillomavirus vaccine. The data has been collected by excel, and the statistical analysis has been performed using Epi-info 7. In this study, collected 76 residents. The mean age of the residents was 30.7 years. Residents were predominantly married (54.2%). The average pregnancy was 0.8. The average age at first intercourse was 27.6. All residents (100%) knew about HPV. Lessons (96.1%) were the principal sources of information. Among residents who knew about HPV, 93.2% consider HPV to be sexually transmitted infections, and 98.7% consider it to be responsible for cervical cancer. In our study, 44 residents (59.5%) were *known to accept to take the HPV vaccine*.

Keywords: cervical cancer, human papillomaviruses vaccine, acceptability, knowledge-e, Senegal.

Abbreviations: EPI: Expanded Programme on Immunization; HPV: Human Papillomavirus; SPSS: Statistical Package for The Social Sciences.

INTRODUCTION

Cervical cancer is a disease that affects young women regardless of their ethnic origins, and it is a public health problem worldwide. 84% of new cases throughout the world occur in developing countries [1, 2, 3]. However, few types of the cancer can be prevented by two methods of prevention: screening for precancerous lesions and vaccination against the Human Papillomavirus (HPV), which prevents infection and is an effective weapon for its eradication. The vaccination of girls between "9-13" is recommended by The World Health Organization (WHO) against HPV. In

Senegal, the vaccine against this cancer was introduced in the Expanded Programme on Immunization (EPI) on 31 October 2018 and targeted all girls aged 9. The vaccination of HPV raises many questions. The goal of our study is to assess resident's levels in gynecology and obstetrics of knowledge of the HPV and their attitudes towards its vaccination in Dakar Aristide Le Dantec teaching hospital, Senegal.

I. MATERIALS AND METHODS

It was across sectional, descriptive and analytical study conducted in Dakar with the residents in gynecology and obstetrics of the University of Cheikh Anta Diop from March 12 to July 30, that's five months and one day. We included in our study all residents in gynecology and obstetrics from de 1st year to the 4th year of the University of Cheikh Anta Diop. We had obtained the consent of each resident at the beginning of the survey. The study was excluded all residents who refused to take part. An information sheet (appendix) was used, as a basis for collecting data from the residents. Socio-epidemiological factors included age, education level, occupation, marital status, gynecological and obstetrical history, knowledge of human papillomavirus, and knowing of vaccination against human papillomavirus and acceptability oh papillomavirus vaccine.

II. STATISTICAL ANALYSIS

It was carried out using of a structured individual interview using a questionnaire. The Epi info software version 7 saved the data collection and the data analysis with the SPSS software (Statistical Package for the social sciences) version 21. The latter consisted of two parts: descriptive and analytical analysis.

It consisted of a bivariate analysis by comparing the acceptance of HPV vaccine administration with the other variables (previous socio-demographic characteristics...). The statistical tests used were the Chi2 test for the percentage comparison, the student test, or ANOVA for the mean comparison. The difference

Corresponding Author α: Aristide Le dantec Teaching Hospital, Gynaecologic and Obstetrics clinic Cheikh Anta Diop University, Senegal. e-mail: gondjout@gmail.com

was statistically significant when the p-value was strictly less than 0.05. The ORs adjusted with their [95% CI] were allowed to know the strength of the link.

III. RESULTS

At the total, 76 residents of gynecology and obstetrics were enlisted. The average age of the

residents was 30.7 years, with the extremes of 26 and 50 years. More than half of residents have more than 30 years. 54.2% of residents were married, as shown in figure 1.




Figure 1: Distribution of married residents (N=76)

The average pregnancy was 0.8, with extremes of 0 to 4 pregnancies. The average age at first intercourse was 22.7 years, with extreme of 13 and 31 years. The age at first pregnancy was 27.6 years, with

extremeness of 22 and 35 years. A medical history of cervical and breast cancer was found in 10.5% and 5.3% of residents, respectively, as reported in figure 2.




Figure 2: Distribution of medical history of cervical and breast cancer (N=76)

In our series, all residents (100%) knew about HPV. Lessons (96.1%) were the principal sources of information. Among residents who knew about HPV,

93.2% consider HPV to be sexually transmitted infections, and 98.7% consider it to be responsible for cervical cancer, as shown in table 1 and figure 3.

Table I: Distribution of residents according to their sources of information on the HPV vaccine (N=76)

Sources on vaccine information on HPV	Number	Percentage
Lessons	73	96,1%
Medias	20	26,3
Entourage	3	3,9




Figure 3: Distribution of residents according to their knowledge of the HPV (N=76)

In our study, more ¾ of residents (72.4%) knew about HPV vaccines. Only half the residents (55.3%) were aware of the type of HPV vaccine, as shown in table II. Gardasil and Cervarix were the most cited.

Tableau II: Distribution of residents according to their knowledge of vaccine against HPV (N=76)

Vaccine against HPV	Number	Percentage
Gardasil + Cervarix	42	55,3
Don't know the name	5	6,6
Gardasil 9	3	3,9
Gardasil 4	3	3,9
Cervarix	2	2,6
Don't know	21	27,6
Total	76	100

Among residents who were aware of the HPV vaccine, only (30.6%) were aware of the side effects. Pain at the site (45.5%) of injection and skin lesions (45.5%) were the most known side effects of residents, as reported in Table III. Among residents who were aware of the HPV vaccine, only forty-five (59.3%) were

aware of the routes of administration of the vaccine. All had cited intramuscular injection as the route of administration. In our study fifty-five residents (76.4%) knew the targets. Girls under 13 years of age were the main targets mentioned by women as reported in Table IV.

Tableau III: Distribution of residents by HPV vaccine side effects (N=2)

Side effects of HPV vaccine	Number	Percentage
Pain at the site of injection	10	45,5
Skin lesions	10	45,5
Neurological trouble	2	9,1
Headache	3	13,6
Vomiting	3	13,6
Fever	2	9,1
Nausea	2	9,1
Sclerosis	2	9,1
Itching	2	9,1
Diarrhea	1	4,5
Infertility	1	4,5
Lupus	1	4,5
Stenosis	1	4,5
Urticaria	1	4,5
Arthralgia	1	4,5

Table IV: Distribution of residents according to targets effects (N=55)

Targets	Number	Percentage
Virgin girl	55	76,4
Virgin teenagers	10	13,9
All people	1	1,4
All women	6	8,3
Total	72	100

Table V: Shows that the distribution of acceptability of the HPV vaccine is almost the same despite knowledge or not about HPV

Knowledge on the HPV vaccine	Acceptability of vaccine				Total	P value	Ods[IC à 95%]
	Yes		No				
	N	%	N	%			
Sources of information on HPV						0,481	
one source	34	61,8	21	38,2	55		1,5[0,5-4,2]
More than one source	10	52,6	9	47,4	19		Ref
HPV responsible of STD						0,388	
Yes	40	59,7	27	40,3	67		2,2[0,3-14,2]
Non	2	40,0	3	60,0	5		Ref
HPV responsible of breast cancer						0,400	
yes	42	58,3	30	41,7	72		0,0
No	1	100,0	0	0,0	1		
Sources of vaccine information						0,253	
one source	30	54,6	25	45,4	55		Ref
More than one source	10	71,4	4	28,6	14		2,1[0,6-7,4]
Type of vaccine						0,782	
Yes	43	59,7	29	40,3	72		1,5[0,1-26,7]
No	1	50,0	1	50,0	2		Ref
Side effects						0,222	
Yes	11	50,0	11	50,0	22		0,5[0,2-1,5]
No	32	65,3	17	34,7	49		Ref
Way of administration						0,430	
Yes	31	55,4	25	44,6	56		0,6[0,2-2,0]
No	10	66,7	5	33,3	15		Ref
Targets						0,236	
Yes	42	58,3	30	41,7	72		
No	2	100,0	0	0,0	2		
Age of vaccin						0,104	
Yes	37	57,8	27	42,2	64		0,2[0,02-1,7]
No	7	87,5	1	12,5	8		Ref
HPV entourage						0,542	
Yes	11	64,7	6	35,3	17		1,4[0,4-4,4]
Non	31	56,4	24	43,6	55		Ref
Lack of information						0,072	
Yes	11	44,0	14	56,0	25		0,4[0,2-1,1]
No	31	66,0	16	34,0	47		Ref

In table V, 60% think HPV is not an STD, and the main cause of refusal of vaccination is le lack of knowledge 56%.

Table VI: Distribution of HPV vaccine acceptability according to the socio-demographic characteristics

Social-demographic characteristics	Acceptability of vaccine				Total	P value	Ods[Ic à 95%]
	Yes		No				
	N	%	N	%			
Age group						0,776	
<30	17	63,0	10	37,0	27		1,1[0,4-3,2]
≥30	22	59,5	15	40,5	37		Ref
Occupation						0,286	
Married	22	56,4	17	43,6	39		0,6[0,2-1,6]
Single	22	68,7	10	31,3	32		Ref
Level of residency						0,000	
1&2	36	76,6	11	23,4	47		7,8[2,7-22,6]
3&4	8	29,6	19	70,4	27		Ref
Gender						0,271	
Female	28	56,0	22	44,0	50		0,5[0,2-1,6]
Male	16	69,6	7	30,4	23		Ref
Address						0,174	
Dakar Suburb	15	78,9	4	21,1	19		2,4[0,7-9,0]
Dakar Center	20	60,6	13	39,4	33		Ref

In our study, 44 residents (59.5%) were known to accept to take the HPV vaccine. According to socio-demographic characteristics, there were statistically significant differences in the acceptability rate the HPV vaccine. For example, residents in the 1st and 2nd year were 7.8 times more likely to accept the vaccine (Table VI). The distribution of acceptability of the HPV vaccine was almost like according to family history (Table VII).

The distribution of the acceptability of the HPV vaccine was almost similar according to the knowledge of HPV (table V).

Table VI shows that the distribution of acceptability of the HPV vaccine varied, according to the education level of the respondents. Indeed, the acceptability of vaccination against HPV was 7.8 times high among 1st and 2nd-year residents.

Tableau VII: Distribution of HPV vaccine acceptability according to family history

Family history	Acceptability of vaccine					Total	P value	Ods[Ic à 95%]
	Yes		No					
	N	%	N	%				
Cancer du col							0,515	
Yes	3	75,0	1	25,0	4			2,1[0,2-21,4]
No	41	58,6	29	41,4	70			Ref
Breast cancer (age group)							0,852	
≤25	5	62,5	3	37,5	8			1,1[0,2-5,2]
>25	39	59,1	27	40,9	66			Ref

IV. DISCUSSION

1. Socio-demographics characteristics

In our study, the average age of residents was 30.7 years; the majority were married (54.2%). The mean age at first intercourse was 22.7 years; the average was 24 years. The mean age at first pregnancy was 27.6 years with extremes (22 and 35). The average pregnancy rate was 0.9, and the average parity rate of 0.8. A medical history of cervical and breast cancer was found in 10.5% and 5.3% of resident, respectively. The resident survey, and data from Gassama can be superimposed our data.

2. Knowledge

During our study, 100% of student knew about HPV, among which 93.2% consider HPV to be an STI and 98.7% consider it to be responsible for cervical cancer which corresponds to the results of a study among students in china with a slightly lower rate of 67.8% and 86.1% respectively [14] and 85.8% in a study in South Africa [5]. However, in a study in Lagos's knowledge about was very poor, as only 39.8% had good acquaintance of the subject [8]. For the residents, knowledge of the main etiological factor is paramount in the prevention of this scourge.

Regarding the source of information, most of the residents interviewed had heard about it through school course, which was consistent with a finding from a Chinese study followed by an entourage, the media (television, radio) [14]. However, in a study in Lagos, the three highest sources of knowledge and information about HPV vaccination among the respondents were identified as, internet (23.2%), television/radio (14.9%), and teachers (12.4%) [8].

As for vaccination, $\frac{3}{4}$ of residents (72.4%) knew about the HPV vaccine. The main sources of information –generally the lessons (96.1%), the media (26.3%) and the entourage (3.9%) as a study in South Africa the principal sources of knowledge reported by the participants were school (60.1%) and the media (33.0%) [5]. Media plays a significant role in getting information to the youth, which can create and raise awareness about cervical cancer and HPV. Residents have a high percentage (100%) of knowledge of the cervical cancer vaccine, and this is related to the level of education and instruction provided during their teaching, and 62.0% knew that the vaccine was available in South Africa [5]. This finding is made by several authors who have shown that, for example, in a study conducted in China, the knowledge they have about the HPV vaccine is mainly related to the teaching given [6,7]. However, in a study in India, none of the students were aware of cancer-causing HPV types and names of the HPV vaccines, which reflect that they have very restrictive knowledge and understanding of the disease [6].

3. Acceptability

In 44 residents, (59.5%) was noted the acceptability of taking the HPV vaccine, corroborating the findings of one Chinese study (57.2% among male and 78.5% among female) [14]; the main reason was the prevention of cervical cancer and concern about the virus and its health consequences. Actually, there is a correlation between the fact of being sensitized about the HPV, the vaccine, and its acceptability.

On the contrary, in France in 2018, a study revealed that a relatively not enough knowledge of the disease does not constitute a barrier to the acceptability of vaccination [10]. The increased reluctance in France of vaccination can be explained by these contradictions. One of the reason of residents for the no acceptability of the vaccination is the lack of knowledge about the anti-HPV vaccine and the lack of information (Consistently, lack of information about HPV infection and vaccines has been identified as a common barrier to the uptake of HPV vaccines in earlier studies (Kahn et al., 2003; Lee et al., 2007; Iliyasu et al., 2010). These are the reasons why lack or less information can hinder vaccination [14]. In India, the reasons for not getting vaccination are cost, safety, efficacy, no knowledge [7]. Reason for unwillingness to accept the HPV vaccine in Lagos is a lack of adequate information on the HPV vaccine (63.2%), fear of negative consequences of receiving the HPV vaccine (12.4%), and fear of injections (8.0%) [8].

Vaccination against cervical cancer is a controversial subject, especially the side effects such, as autoimmune diseases, multiple sclerosis as reported in literature. These side effects are not attributable to vaccination and yet are major obstacles to the acceptability of vaccination. In our study, the most well-known side effect is the reaction at the injection site (4.5%).

We can, therefore, consider that have already received information on cervical cancer and its prevention methods promotes vaccination. It should, therefore, encourage us to inform as many representatives of the medical community as possible so that they can correctly relegate information to the population on HPV vaccination.

Also, the second barrier to HPV vaccination was related to the vaccine itself, the fear of side effects, and this matches with the literature. Fear of side effects is at the top of the list in most acceptability surveys.

In an American study published in 2013, there was even an increase in this concern, with 4.5% of parents worried about possible side effects in 2008, compared to 16.4% in 2010 [11].

V. CONCLUSION

Acceptability of HPV vaccination requires knowledge of the papillomavirus and vaccination. Awareness remains an essential element in the prevention strategy.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Acceptation vaccinale - Regards croisés, Colloque, Calenda, disponible le mercredi 01 août 2018 sur le site <https://calenda.org/463129> consulté le 03 Mars 2019.
2. Aubin F, Mougin C, Prétet J-L. Papillomavirus humains: biologie et pathologie tumorale. internationales TD-EM ed, DL. 06/2003.
3. Badr.RE. a sensitive and specific marker of HPV-associated squamous lesions of the cervix. American Journal of Surgical Pathology. 2008;32(6):899-906.
4. Beutner KR, Wiley DJ, Douglas JM, Tying SK, Fife K, Trofatter K, Stone KM. Genital warts and their treatment. Clin Infect Dis 1999; 28S1: S37-S56.
5. Nathaniel Mofolo. Knowledge of cervical cancer, human papillomavirus and preventon among first-year female students in residences at the University of the Free State. African Journal of Primary Health Care & Family Medicine ISSN: (Online) 2071-2936, (Print) 2071-2928, Published: 24 May 2018.
6. Shazia Rashid1. Knowledge, Awareness and Attitude on HPV, HPV Vaccine and Cervical Cancer among the College Students in India. PLOS ONE | DOI:10.1371/journal.pone.0166713 November 18, 2016.
7. Sunite A. Ganju. Assessment of knowledge and attitude of medical and nursing students towards screening for cervical carcinoma and HPV vaccination in a tertiary care teaching hospital. International Journal of Community Medicine and Public Health Ganju SA et al. Int J Community Med Public Health. 2017 Nov;4(11):4186-4193 pISSN 2394-6032 | eISSN 2394-6040.
8. Olajumoke Adetoun Ojeleye. Knowledge and acceptance of HPV vaccination among Lagos students. AFRICAN JOURNAL OF MIDWIFERY AND WOMEN'S HEALTH, APRIL-JUNE 2019, VOL 13, NO 2.
9. Ekaterina Shemelova. Facteurs influençant la prise de décision sur la vaccination contre le HPV. Médecine humaine et pathologie. Université Grenoble Alpes, 2017. Français. ffNNT: 2017GREAH015ff. fftel-01691595.
10. Ferenczy A. Epidemiology and clinical pathophysiology of condylomata acuminata. Am J Obstet Gynecol 1995; 172: 1331-9.
11. Gassama.O. bilan des activités de colposcopie au service de gynécologie-obstétrique du CHU A.LeDantec: FMPOS Dakar, Année; 2011 Thèse N° 43, Aristide Le Dantec.
12. Hatch KD. Clinical appearance and treatment strategies for human papillomavirus: a gynaecologic perspective. Am J ObstetGynecol 1995; 172: 1340-4.
13. ICO (Institut Català Oncologia) Information Centre on HPV and cancer Sénégal: Human Papillomavirus and related cancers fact sheet, Dec 15, 2014.
14. Julia Baum-Durrenberger. les connaissances actuelles des étudiants concernant les papillomavirus humains, enquête réalisée auprès d'étudiants lorrains sur leur connaissance du virus, du vaccin, des moyens de prévention et de dépistage. Thèse pour obtenir le grade de docteur en médecine. Université de Lorraine 2012, le 26 septembre 2012.
15. Omar Gassama, Acceptability of Vaccination Against Papillomaviruses: A Study On 115 Women at the Matlaboul Fawzayni Hospital in Touba (Senegal), journal of gynaecology and women's health, Volume 17 Issue 3- November 2019, DOI: 10.19080/JGWH.2019.17.555965.
16. Chun-Jing Fu. Knowledge, Perceptions and Acceptability of HPV Vaccination among Medical Students in Chongqing, China Asian Pacific Journal of Cancer Prevention, Vol 15, 2014.