

Origanum vulgare (Oregano) and its Carvacrol Biocomponent as an Alternative of Antimicrobial Agent

Natalia Kaori Aida, Janaina Priscila Barbosa, Thais Rossini de Oliveira, Vanessa da Silva Cardoso, Simone Nataly Busato de Feiria, Giovana Claudia Boni, Jose Francisco Hofling

Received: 6 December 2019 Accepted: 31 December 2019 Published: 15 January 2020

Abstract

The use of plants as an alternative to medicinal treatments is an old practice. The increased resistance of microorganisms to conventional antimicrobials has made studies with medicinal plants increasingly relevant, and ethnobotanical and ethnopharmacological knowledge is considered essential for the development of new drugs. The essential oil of *Origanum vulgare* and its isolated compound Carvacrol have antimicrobial effects demonstrated in the literature as antibacterial and antifungal activity. Therefore, the present study evaluated the antibacterial and antifungal activity of *O. vulgare* and Carvacrol using the broth microdilution method (CLSI, 2008), determining MIC (Minimum Inhibitory Concentration) and MFC and MBC (Minimum Fungicidal Concentration and Concentration Minimum Bactericide). Used as standard comparative the antimicrobials Fluconazole and Chlorhexidine.

Index terms— *origanum vulgare*; carvacrol; *candida* spp., *streptococcus* spp., *staphylococcus aureus* MRSA.

Introduction the use of plants as an alternative to medicinal treatments is an ancient practice, which provide primary health care to 80% of the world's population. It has also been an important source for new drug discoveries (Wangchuk & Tobgay, 2015). World Health Organization has estimated that around 80% of the population of developing countries use traditional herbal medicines as a source of treatment for diseases as a cheap and alternative source, other factors such as the lack of modern health facilities, Cultural priorities and choices also contribute to the use of medicinal plants as a therapeutic alternative (Aziz et al., 2018). Studies on medicinal plants have become increasingly relevant, ethnobotanical and ethnopharmacological knowledge is considered essential for the development of new drugs. Researches studies the effectiveness and use of traditional plants as alternative medicine and adjuvants in a treatment (Amjad et al., 2017). In this context, *Origanum vulgare* is a plant used since ancient times as a traditional cure for treating infections (Karaman et al., 2017). Studies show that *O. vulgare* essential oil is rich in Carvacrol, has anti-inflammatory, antioxidant, antispasmodic, antimicrobial, and antifungal activity (Król et al., 2019). Also, resistant microorganisms have emerged over the years and have been considered a global health threat. Microbial resistance has brought problems in the treatment of infectious diseases, and the development of new antimicrobial agents is required (Ayaz et al., 2019). The occurrence that strains of the genus *Candida* have been resistant to treatment with commercial antifungals, belonging to the azoles and polyenes family, has been a reason for concern by health professionals. *Candida* infections mainly affect immunosuppressed patients (Barbosa et al., 2019). Given this scenario, the plants are promising antimicrobial agent for the development of new drugs, acting in disease prevention and treatment (Ayaz et al., 2019). Therefore, this work aimed to study the antimicrobial activity of *Origanum vulgare* essential oil and its isolated compound Carvacrol against microorganisms of genus *Candida*, *Streptococci* and, *Staphylococci*.

1 II.

2 Material and Methods

3 a) Essential oil

The essential oil was purchased commercially from Quinari Fragrâncias e Cosméticos LTDA and the isolated compound from Sigma Aldrich. Essential Oil -Lot: 09818DIV Isolated Compound -Lot: S40656V Origanum vulgare (Oregano) and its Carvacrol Biocomponent as an Alternative of Antimicrobial Agent range of 0.08 to 0.1 equivalent to 5.0×10^6 CFU/mL. Then, the inoculum was standardized, diluting to 2.5×10^3 , according to CLSI, 2008.

In sterile microplate was deposited with 100 µl of RPMI, 100 µl of the essential oil or 100 µl of the isolated compound at the initial concentration, followed by serial microdilution and then added with 100 µl of the adjusted inoculum.

Groups of test: CG: positive control group; OL: Essential Oil Treatment Group; IC: Treatment group with isolated compound; AC: Treatment group with the commercial antifungal Fluconazole at an initial concentration of 64 µg/mL (CLSI, 2002). Incubated the plates for 48h at 37°C. Defined the MIC as the lowest concentration of compound that did not exhibit visible growth of the microorganism.

4 d) Inoculum adjustment of the Streptococcus spp. and Staphylococcus aureus MRSA

After growth in BHI liquid culture medium, the inoculum was adjusted by spectrophotometer with a wavelength of 625nm in the range 0.08 to 0.1 absorbance, equivalent to 1.0×10^8 CFU/mL.

In a sterile microplate were deposited 100 µl of BHI, 100 µl of the essential oil or 100 µl of the isolated compound at the initial concentration, followed by serial microdilution, and then added of 100 µl of the adjusted inoculum.

Groups of test: CG: positive control group; OL: Essential Oil Treatment Group; IC: Treatment group with isolated compound; AC: Treatment group with the commercial antifungal chlorhexidine at the initial concentration of 64 µg/mL (CLSI, 2008). Incubation of the plates was made for 24h at 37°C with 10% CO₂ for Streptococcus strains and aerobically for Staphylococcus aureus MRSA. Defined the MIC as the lowest concentration of compound that did not exhibit visible growth of the microorganism.

5 e) Determination of Minimum Fungicidal Concentration (CFM) and Minimum Bactericidal Concentration (CBM)

In a petri dish containing Sabouraud Dextrose Agar (SDA) for yeast and Brain Heart Infusion Agar (BHI) for bacteria tested the determination of minimum fungicidal concentration (MFC) and minimum bactericidal concentration (MBC). Homogenized the wells containing target concentrations and transferred the aliquot to the Petri dish with the solid culture medium. After incubation at 37°C for 24h, established the lowest fungicidal/bactericidal concentration. Determined the MFC/MBC as the lowest concentration of essential oils and isolated compounds that did not allow the growth of any colony of the microorganism on the solid medium after the incubation period. Through visual reading, the inhibition of growth or death provided by the tested substances was confirmed (GULLO et al., 2012).

6 III.

7 Results

8 a) Evaluation of Minimum Inhibitory Concentration (MIC)

and Minimum Fungicidal / Bactericidal Concentration (CFM / CBM) Tested the Origanum vulgare essential oil and its isolated compound Carvacrol against reference strains of the genus Candida, Streptococcus, and Staphylococcus aureus methicillin-resistant strain to determine its inhibitory effect by broth microdilution technique. Both substances showed inhibitory activity on microbial cells (Table ??). For reference strains of the genus Candida, inhibitory concentrations were between 0.125 mg/mL and 0.5 mg/mL for O. vulgare essential oil and 0.125 mg/mL and 0.0625 mg/mL for Carvacrol isolated compound. The antifungal Fluconazole was also tested against strains to determine the minimum inhibitory concentration by broth microdilution technique. Fluconazole has shown inhibitory activity between 1 and 32 µg/ml (Table ??). For bacterial strains of the genus Streptococcus, the essential oil of O. vulgare behaved similarly, varying the MIC between concentrations of 0.5 mg/mL and 0.250 mg/mL, as well as its isolated compound Carvacrol. The essential oil and Carvacrol inhibited the strain Staphylococcus aureus at concentrations of 0.5 and 0.250 mg/mL respectively. Chlorhexidine antimicrobial was also tested against strains to determine the minimum inhibitory concentration by broth microdilution technique. Chlorhexidine demonstrated inhibitory activity between 3.75 to 7 µg/ml (Table ??). After the determined the MIC values, used an aliquot of the susceptibility test to determine the minimum

fungicidal/bactericidal concentration (MFC/MBC) of the strains. For *Candida* spp. Strains, *O. vulgare* essential oil, showed fungicidal activity against *Candida* spp. strains, varying its effect on concentrations between 0.5 mg/mL and 0.250 mg/mL; while Carvacrol showed 0.250 mg/mL fungicidal activity for all *Candida* strains tested (Table ??). *Streptococcus* bacterial strains showed a bactericidal concentration ranging from 0.5 mg/mL to 0.250 mg/mL for the essential oil and ranged from 0.5 mg/mL to 0.125 mg/mL for the isolated compound. *S. aureus*(MRSA) showed a bactericidal concentration of 0.5 mg/mL for both.

Table ??: Visual reading results of MIC and CFM/CBM of the strains tested.

9 IV.

10 Discussion

Excessive and indiscriminate use of antimicrobials is a major determinant of some emerging infections, selection of resistant pathogens, and the continued development of antimicrobial resistance globally. The increasing emergence of multi-drug resistant organisms and the limited development of new agents available to combat them have caused an imminent crisis with alarming implications (Sartelli et al., 2016). In view of the increasing numbers of cases of conventional drug-resistant microorganisms, researchers are looking for alternatives to biocompounds that have antimicrobial properties against microorganisms. Studies with plants as promising agents in the search for new compounds.

In this study, the obtained data showed antimicrobial activity of the tested essential oil, as well as its isolated compound against planktonic cells of *Candida* spp., oral *Streptococcus* species, and *S. aureus* methicillin resistant strain. The essential oil inhibited antimicrobial growth between concentrations of 0.5 to 0.125mg/mL against all strains tested. At the same time, the isolated compound showed antimicrobial activity between concentrations of 0.250 to 0.0625mg/mL against all strains tested. The minimum fungicidal/bactericidal concentration (MFC/MBC) of the essential oil in the strains tested was between 0.5 mg / ml and 0.250 mg / ml. The isolated compound showed MFC/MBC between concentrations of 0.5 mg/mL to 0.125 mg/mL (Table ??).

These data, initially reveal the antimicrobial action of this essential oil, as well as its isolated compound, corroborating with the literature, pointing out

11 Reference strain

12 *O. vulgare* Carvacrol

The data obtained in this study added to the data in the recent literature, suggest that the essential oil of *O. vulgare* and also its isolated compound Carvacrol show antibacterial and antifungal potential.

These data open possibilities for many other studies, such as the performance of these oils and biocomponents in mature biofilms and multispecies biofilms, evaluating cell viability and possible morphological changes, added to cytotoxicity tests, action on cancer cells, in an attempt to add more information about this plant and its use as an alternative agent in the treatment of infections and acting as coadjuvants.

V.

13 Conclusion

? Both the essential oil of *O. vulgare*, and its isolated compound Carvacrol, can inhibit the growth of the tested microorganisms in low concentrations; ? The isolated compound is more effective when compared to the essential oil, inhibiting the microorganisms in a lower concentration; ? About the microorganisms, essential oil and Carvacrol are more effective against *Candida* spp. when compared to bacterial strains; ? Both the EO and its main biocomponent Carvacrol show fungicidal/bactericidal activity against the tested strains

-
- [Karaman et al.] , M Karaman , M Bogavac , B Radovanovi? , J Sudji , K Tesanovic , L Janjusevic .
- [Duarte et al. ()] ‘Anti-Candida activity of Brazilian medicinal plants’. M C Duarte , G M Figueira , Sartoratto Um , V L Rehder , C Delarmelina . *J Ethnopharmacol* 2005. 97 (2) p. .
- [Barbosa et al. ()] ‘Anti-Candida Activity of Essential Oils from Eucalyptus species. A Preliminary Study’. J P Barbosa , T R Oliveira , Dgpb Puppín , A L Teixeira , G C Boni , Snb Feiria , J F Höfling . *Adv Dent & Oral Health* 2018. 8 (3) p. .
- [Barthi et al. (2013)] *Antibacterial activities of Origanum vulgare alone and in combination with different antimicrobials against clinical isolates of Salmonella typhi*. AncSci Life, V Barthi , N Vasudeva , S Sharma , J S Duhan . 2013 Apr-Jun. 32 p. .
- [Gullo et al. ()] *Antifungal activity of Maytenin and Pristimerin*, F P Gullo , Jco Sardi , Vaffm Santos , F Sangalli-Leite , N S Pitangui , S A Rossi , Paula De , Aca Silva . 2012. Hindawi Publishing Corporation.
- [Sartelli ()] ‘Antimicrobials: a global alliance for optimizing their rational use in intra-abdominal infections’. M Sartelli . *World J EmergSurg* 2016. 11 p. 33.
- [Bhat et al. ()] ‘Characterization of Herbal Antifungal Agent, Origanum vulgare against Oral Candida spp. Isolated from Patients with Candida-Associated Denture Stomatitis: An In vitro Study’. V Bhat , S M Sharma , V Shetty , C S Shastry , Vaman Rao , C Shenoy , S Saha , S Balaji , S . *Contemp Clin Dent* 2018 jun. 9 p. .
- [Wangchuk and Tobgay ()] ‘Contributions of medicinal plants to the Gross National Happiness and Biodiscovery in Bhutan’. P Wangchuk , T Tobgay . *J Ethnobiol Ethnomed* 2015. 11 p. 48.
- [Amjad et al. ()] *Descriptive study of plant resources in the context of the ethnomedicinal relevance of indigenous flora: A case study from Toli Peer National Park*, M S Amjad , M F Qaeem , I Ahmad , S U Khan , S K Chaudhari , N Z Malik , H Shaheen , A M Khan . 2017. 12 p. e0171896. Azad Jammu and Kashmir, Pakistan. PLoSOne
- [Cleff et al. ()] ‘In vitro activity of Origanum vulgare essential oil against Candida species’. M B Cleff , A R Meinerz , M Xavier , L Schuch . *Brazilian Journal of Microbiology* 2010. 41 (1) p. .
- [Król et al. (2016)] ‘Investigation of the antifungal activity of carvacrol against strains of Cryptococcus neoformans’. B Król , B Kolodziej , B Kedzia , E Holderna-Kedzia , D Sugier , K Luchowska , R O Nóbrega , A P Teixeira , W A Oliveira , E O Lima , I O Lima . *J Sci Food Agric* 2019. 2016 Nov. 13 (11) p. . (Pharm Biol)
- [Wang et al. (2016)] ‘Membrane Destruction and DNA Binding of Staphylococcus aureus Cells Induced by Carvacrol and Its Combined Effect with a Pulsed Electric Field’. L H Wang , M S Wang , X A Zeng , Z H Zhang , D M Gong , Y B Huang . *J AgricFoodChem* 2016 Aug 17. 64 (32) p. .
- [Clsi ()] *Método de referência para testes de diluição em caldo para a determinação da sensibilidade de leveduras a terapia antifúngica*. Clinicaland laboratorial standards institute, Clsi . 2008. 2008. (CLSI Norma M27-A3, v.28, n.14, 2 ed.)
- [Origanumvulgare essential oil affects pathogens causing vaginal infections Journal of Applied Microbiology ()] ‘Origanumvulgare essential oil affects pathogens causing vaginal infections’. *Journal of Applied Microbiology* 2017. 122 p. .
- [Ayaz et al. (2019)] *Synergistic interactions of phytochemicals with antimicrobial agents: Potential strategy to counteract drug resistance*. ChemBiol Interact, M Ayaz , F Ullah , A Sadiq , F Ullah , M Ovais , AhmedJ , Devkotahp . 2019 May 31. 308 p. .
- [Aziz et al. ()] ‘Traditional uses of medicinal plants practiced by the indigenous communities at Mohmand Agency, FATA, Pakistan’. M A Aziz , Adnan M Khan , A H Shahat , A A Al-Said , M S Ullah , R . *J Ethnobiol Ethnomed* 2018. 14 (2) .