



GLOBAL JOURNAL OF MEDICAL RESEARCH: E
GYNECOLOGY AND OBSTETRICS
Volume 20 Issue 4 Version 1.0 Year 2020
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Obesity in Pregnant Adolescents: Clinical Correlates, Maternal and Fetal Outcomes

By Cecília Ogando Alfama, Amanda Vilaverde Perez, Mariana Sbaraini, Marianna Sperb, Rafaela da Silveira Corrêa, Edimárlei Gonsales Valério, Maria Lúcia Oliveira Rossés, Vera Lúcia Bosa & Janete Vettorazzi

Universidade Federal do Rio Grande do Sul (UFRGS)

Abstract- Objective: To evaluate the relationship between pregestational obesity and maternal and fetal outcomes in pregnant adolescents.

Methods: Cross-sectional study conducted among puerperal teenagers of a university hospital in Southern Brazil. Structured questionnaires on sociodemographic and obstetric data were applied.

Results: We evaluated 500 puerperal adolescents with a mean age of 17.77 ± 1.36 . Of these, 31.2% (n=156) had pregestational weight excess. Pregestational weight excess was significantly associated with gestational hypertension ($p=0.037$) and diabetes ($p=0.045$), cesarean delivery ($p=0.040$), and lower adherence to prenatal care (24.7%, $p=0.007$).

Keywords: obesity, adolescent, pregnancy outcomes.

GJMR-E Classification: NLMC Code: WQ 209



Strictly as per the compliance and regulations of:



© 2020. Cecília Ogando Alfama, Amanda Vilaverde Perez, Mariana Sbaraini, Marianna Sperb, Rafaela da Silveira Corrêa, Edimárlei Gonsales Valério, Maria Lúcia Oliveira Rossés, Vera Lúcia Bosa & Janete Vettorazzi. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Obesity in Pregnant Adolescents: Clinical Correlates, Maternal and Fetal Outcomes

Cecília Ogando Alfama ^α, Amanda Vilaverde Perez ^σ, Mariana Sbaraini ^ρ, Marianna Sperb ^ω,
Rafaela da Silveira Corrêa [¥], Edimárlei Gonsales Valério [§], Maria Lúcia Oliveira Rossés ^χ,
Vera Lúcia Bosa ^ν & Janete Vettorazzi ^θ

Abstract- Objective: To evaluate the relationship between pregestational obesity and maternal and fetal outcomes in pregnant adolescents.

Methods: Cross-sectional study conducted among puerperal teenagers of a university hospital in Southern Brazil. Structured questionnaires on sociodemographic and obstetric data were applied.

Results: We evaluated 500 puerperal adolescents with a mean age of 17.77 ± 1.36 . Of these, 31.2% (n=156) had pregestational weight excess. Pregestational weight excess was significantly associated with gestational hypertension (p=0.037) and diabetes (p=0.045), cesarean delivery (p=0.040), and lower adherence to prenatal care (24.7%, p=0.007). Mothers with pregestational weight excess also gained more weight than recommended during pregnancy and had babies with a higher weight (p=0.02) and length (p=0.018).

Conclusion: Pregestational weight excess in adolescence is related to unfavorable maternal and fetal outcomes. We emphasize the importance of a closer look at this population still in development to identify nutritional imbalances and promote adequate weight gain.

Keywords: obesity, adolescent, pregnancy outcomes.

Author α ω § χ θ: Postgraduate Program in Health Sciences: Gynecology and Obstetrics, Universidade Federal do Rio Grande do Sul (UFRGS). Rua Ramiro Barcelos, 2400, Santa Cecília, 90035-002, Porto Alegre, RS, Brazil. e-mails: cee.alfama@gmail.com, mariannasperb@yahoo.com.br, mariarosses@gmail.com

Author σ ρ § θ: Faculty of Medicine (FAMED), Universidade Federal do Rio Grande do Sul. (UFRGS). Rua Ramiro Barcelos, 2400, Santa Cecília, 90035-002, Porto Alegre, RS, Brazil. e-mails: marisbaraini@gmail.com, amandavperez@gmail.com

Author ¥: Department of Nutrition, Faculty of Health Sciences, Centro Universitário Ritter dos Reis - UniRitter. Rua Orfanotrófio, 555, Santa Tereza, 90840-440, Porto Alegre, RS, Brazil. e-mail: rafaeladscorrea@gmail.com

Author § θ: Department of Gynecology and Obstetrics, Faculty of Medicine (FAMED), UFRGS. Rua Ramiro Barcelos, 2400, Santa Cecília, 90035-002, Porto Alegre, RS, Brazil. e-mails: evalerio@hcpa.edu.br, jvetorazzi@hcpa.edu.br

Author θ: Service of Gynecology and Obstetrics. Hospital de Clínicas de Porto Alegre (HCPA). Rua Ramiro Barcelos, 2350/1125, Santa Cecília, 90035-003, Porto Alegre, RS, Brazil.

Author ν: Department of Nutrition, Faculty of Medicine (FAMED) and Postgraduate Program in Food, Nutrition and Health, Universidade Federal do Rio Grande do Sul (UFRGS). Rua Ramiro Barcelos, 2400, Santa Cecília, 90035-002, Porto Alegre, RS, Brazil. e-mail: vbosa@hcpa.edu.br

I. INTRODUCTION

Adolescence is a period marked by transitions and search for new experiences, such as sexual awakening, making girls more susceptible to unwanted pregnancy. Poverty and socioeconomic inequalities can further increase rates of teenage motherhood, mostly due to knowledge gaps and lack of information about contraception.

In Brazil, 547,564 babies are born annually to adolescent mothers, accounting for 18% of all births and 11.43% of births in the South region¹. Adolescence is a period of woman's life when major physical, cognitive, and social transformations occur, marking the transition from childhood to adulthood. Adolescent pregnancy is a public health problem and may be related to the occurrence of maternal-fetal complications such as preeclampsia, prematurity, infections, low fetal weight, among others^{2,3}.

There is a close relationship between maternal nutritional status before and during gestation and neonatal outcomes⁴. Pregnancy before the age of 15 years old can be harmful to maternal and fetal health since both are under physical development and can compete for nutrients. This competition may increase the odds of low neonatal weight, micronutrient deficiencies, and fetal growth restriction⁵.

Obesity is a complex and multifactorial disease with rising prevalence worldwide⁶. In the last decades, rates of weight excess among Brazilian adolescent girls have nearly tripled⁷. Pregestational obesity is a risk factor for excessive weight gain during pregnancy and is associated with complications such as diabetes, postpartum hemorrhage, infections, thromboembolic events, and disturbs in childbirth⁸. It can also cause hypertensive syndromes, which lead to poor perinatal outcomes⁹. Maternal obesity is also associated with neonatal events such as fetal macrosomia, increased admission in the neonatal intensive care unit, congenital disabilities, hypoglycemia, low Apgar score, and death¹⁰.

The combination of adolescent pregnancy and obesity can further impact health outcomes in both generations. Therefore, this study aims to evaluate the relationship between pregestational weight excess and the occurrence of maternal and fetal outcomes among

pregnant adolescents in a university hospital in Southern Brazil.

II. METHODS

Between November 2014 to March 2016, we conducted a cross-sectional study carried out among puerperal adolescents (10 to 19 years old) that attended a university hospital in southern Brazil. We interviewed the adolescents in the postpartum period, after they signing a free and informed consent form and answered questionnaires on socio-demographic, gestational and perinatal data. Ethics Committee of Hospital de Clínicas de Porto Alegre approved the project. (protocol number 14-0491).

Pregestational nutritional status was analyzed using Anthroplus software, and body mass index (BMI) z-score from 1 to -1 was considered eutrophic and $\geq +1$ weight excess¹⁴. We evaluated gestational BMI according to the Atalah curve published in 1997¹⁵. Weight gain during pregnancy was defined suitable from 12.5-18 kg for those with initial low weight, 11.5-16 kg for eutrophic, 7-11.5 kg for overweight, and 5-9 kg for those obese, as recommended by the IOM (*Institute of Medicine*)¹⁶.

Maternal clinical variables evaluated were menarche, age at first sexual intercourse, number of gestations/parity, type of delivery, number of prenatal visits, previous and gestational diseases such as hypertension, diabetes, gestational hypertension and diabetes, obesity, anemia, and preeclampsia. According to the World Health Organization (WHO) criteria, the birth is preterm when it occurs before 37 weeks¹⁷. We classified hypertensive disorders according to the recommendation of ACOG (*American Congress of Obstetricians and Gynecologists*)¹⁸.

Fetal variables collected were gestational age at birth, sex, weight, length, hospitalization in the neonatal intensive care unit (ICU), and complications such as respiratory dysfunction and Apgar score. We evaluated gestational age at birth using the Capurro method.

Data were analyzed using SPSS (*Statistical Package for the Social Sciences*) version 18.0 (SPSS Inc., Chicago, IL, EUA®). Categorical variables are presented by absolute number (n) and percentage (%), symmetric continuous variables by the mean and standard deviation (SD), and continuous variables asymmetric by the median and interquartile range [25-75 percentile]. We used a Chi-square test for categorical variables and t-test for quantitative variables. We considered, for all analyses, a significance level of 5% ($p < 0.05$) and a 95% confidence interval.

III. RESULTS

We evaluated five hundred adolescents and their respective newborns. Maternal mean age was 17.77 ± 1.36 years, with the mean age at first sexual

intercourse at 14.8 ± 1.5 years. 57.6% (n=235) had a family income of 1.5 to 3 minimum wages, and 63% had white skin color. Primiparous women composed 83.4% of the sample, and 84.8% (n=420) were single with a partner. Regarding pregestational nutritional status, 31.2% (n=156) were weight excess. Table 1 shows other characteristics of the sample.

Table 2 shows the association of pregestational body mass index classification with maternal and fetal outcomes. Mothers with pregestational weight excess were more likely to attend fewer prenatal consultations (24.7% vs. 36.3%, $p=0.007$) and have excessive weight gain during pregnancy than the others (62.2% vs. 32.9%, $p < 0.001$). Prevalence of gestational hypertension (5.1% vs. 1.7%, $p=0.037$) and gestational diabetes (4.5% vs. 1.5%, $p=0.045$) were higher in the weight excess group when compared to the lean/eutrophic group. The proportion of birth by cesarean was also significantly higher in weight excess adolescents (24.4% vs. 17.2%, $p=0.04$).

Children of weight excess mothers had a longer length (48.3 cm vs. 47.5 cm, $p=0.018$) and weight (3.199 kg vs. 3.040 kg, $p = 0.020$) at birth, as observed in figures 1 and 2, respectively. Prematurity, neonatal ICU admission, respiratory dysfunctions, and Apgar score were not associated with pregestational weight excess.

IV. DISCUSSION

Pregnancy in women before the age of 19 years is characterized by poorer outcomes when compared to those aged 20-24 years¹⁹. One explanation for this is that teenage mothers are smaller, due to uncompleted growth at conception. Competition for nutrients can occur between the still developing mother and the fetus, causing developmental impairment in one or both parts. The deficiency of nutrients and energy due to gestation can further impair growth and make adolescent mothers more prone to metabolic imbalances such as obesity and cardiovascular diseases in adulthood²⁰.

Obesity is a growing public health problem. In the United States, the prevalence of overweight in girls aged 12 to 19 years was 4.7% in 1970, increasing to 15.4% in 2002²¹. In Brazil, weight excess affects more than 23% of teenage women⁷. Obesity during adolescence can persist in adulthood, increasing the risk of associated diseases such as hypertension, diabetes mellitus, osteoarthritis, and cardiovascular disease²². Furthermore, maternal obesity is associated with adverse obstetric outcomes, such as increased cesarean rates, preeclampsia and, gestational diabetes. A menarche is a late event in the pubertal process that is related to genetic, environmental, and nutritional factors. Literature shows that adolescents with a higher risk of obesity in adulthood are those with early menarche and those with a short period between

menarche and first parturition³⁷. We also observe this correlation in our study, as weight excess young women had menarche at an earlier age.

During pregnancy, women with overweight and obesity should gain a maximum of 11,5 and 9 kg, respectively²⁶. Adolescent mothers are more likely to have inappropriate weight gain compared with adult women³⁶, and over 60% of teenage mothers have excessive weight gain. In our study, 31.2% of them had pregestational overweight, and the majority of those gained gestational weight above the recommended for their BMI before pregnancy, corroborating with previous findings^{31,34}. The weight gain can be explained because the body of adolescents has to work to both meet their own growth needs and those of the fetus.

Prenatal consultations provide the necessary medical support during pregnancy, and performing them correctly is necessary to prevent possible complications associated with maternal and fetal health. Pregnant women with weight excess were significantly more likely to have inadequate prenatal care, having less than the six visits recommended by the Brazilian Ministry of Health³². Adolescent mothers may not attend gestational care due to fear or non-acceptance of the pregnancy; therefore, professionals should be rightly trained to guide and stimulate them about gestational care³².

Regarding maternal outcomes, we found a significant association of maternal weight excess with hypertension and gestational diabetes mellitus (GDM), corroborating literature previous findings. Sukalich et al. showed that cesarean delivery, preeclampsia, and gestational diabetes were more frequent in pregnant women with a BMI greater than 25, as well as labor induction and cephalic-pelvic disproportion³⁶. Another study also demonstrated that maternal weight excess and excessive weight gain during pregnancy were associated with an increased risk of gestational hypertension⁴⁶. The association between obesity and GDM can be explained by more propensity of decreased insulin sensitivity before and during gestation in these women^{47,42}.

In our study, 24.4% of overweight adolescents had cesarean deliveries. A study with obese teenage mothers at hospitals in the United States⁴² and Colombia²⁰ found similar results. High prevalence of cesarean deliveries may be related to higher fetal weight in mothers with a still-developing body structure⁴³. Furthermore, we stand out that our study obtained a mean number of the cesarean sections below the national average reported by the Brazilian National Health System Database of 2011⁴⁴. This finding may be related to the fact that these adolescents were in a tertiary hospital that prioritizes vaginal delivery and has specific institutional protocols regarding medical indications of cesarean.

Excessive weight gain during pregnancy and pregestational weight excess can result in newborns with bigger weight and length^{46,48,49}. These findings are in agreement with our study since a significant association of maternal weight excess and anthropometric variables of the babies was found. Pregnant women who are overweight, obesity, or underweight can still have babies of adequate birth weight, provided that they gained appropriate weight during pregnancy¹¹.

V. CONCLUSION

Adolescence is a period of transition in which girls are still in the stage of physical and psychological growth. Gestation alone, at this stage, is already a risk factor for adverse outcomes. Our study demonstrates that these adverse outcomes are potentiated when gestation in adolescence is associated with obesity. Excessive weight increase during pregnancy is also associated with adverse maternal and fetal outcomes, such as preeclampsia, increased rate of cesarean section, and macrosomia. These findings highlight the necessity of further studies in the adolescent population, which accounts for 20% of all deliveries.

This study has a few limitations. Among the significant results, such as the association between obesity and gestational hypertension, confusion factors may be involved. Also, we obtained pre-pregnancy information from participants' reports, depending on memories.

Our study contributes to a better understanding of pregestational weight excess in teenage mothers. The assessment of the initial nutritional status is essential to adjust weight gain during pregnancy, to minimize possible harms to the mother-child binomial. Although historically, the concern with teenage motherhood involves low weight gain and nutritional deficits, increasing rates of obese pregnant adolescents make interventions in eating habits necessary to change this paradigm. Therefore, educational programs orientating this population about contraception, pregnancy, and healthy eating habits should be implemented, as well as actions encouraging early prenatal care.



Table 1: Characteristics of pregnant adolescents and their newborns attended at a university hospital.

	Average	Standard deviation
Mean maternal age, years	17.77	1.36
Mean paternal age, years	22.55	5.26
Mean years of study, years	9.29	2.01
Mean age at first sexual intercourse, years	14.78	1.48
Mean menarch age, years	12.15	1.69
	N	%
Marital status		
Single with partner	420	84.8
Single without partner	62	12.5
Married	13	2.6
Family income		
Up to 1.5 minimum wages	113	27.7
1.5 to 3 minimum wages	235	57.6
Over 3 minimum wages	60	14.7
Referred skin color		
White	315	63.0
Non white	185	37.0
Maternal height		
≤150 cm	30	6.0
>150 cm	470	94.0
Pregestational BMI*		
Underweight	7	1.4
Eutrophic	337	6.4
Overweight	100	20
Obesity	47	9.4
Severe obesity	9	1.8
Weight gain during pregnancy		
Insufficient	147	29.5
Adequate	142	28.5
Excessive	210	42.1
Parity		
Primiparous	416	83.4
Multiparous	83	16.6
Delivery type		
Vaginal	403	80.6
Cesarian	97	19.4
Gestational age		
Preterm**	69	13.8
Full term	430	86.2
Newborn		
Female sex	250	50.2
Length at birth, cm	47,72	3.37
Weight at birth, grams	3085,87	575.35

BMI = Body mass index. * According to criteria by WHO. ** Before 37 weeks of gestation.



Table 2: Association between pregestational weight and maternal and fetal outcomes among adolescent mothers attended at a university hospital in Southern Brazil.

	Underweight/Eutrophy	Weight excess ¹	P value
Mean age at menarche, years	12.38 ± 1,67	11.62 ± 1.63	<0.001*
Apgar score			
1st minute	7.85 ± 1.94	7.95 ± 1.57	0.594
5th minute	9.03 ± 1.58	9.06 ± 1.01	0.833
Number of prenatal care consults			
< 6 consults	122 (36.3)	38 (24.7)	0.007
≥ 6 consults	214 (63.7)	116 (75.3)	
Weight gain during pregnancy			
Insufficient	124 (36.2)	23 (14.7)	<0.001*
Adequate	106 (30.9)	36 (23.1)	
Excessive	113 (32.9)	36 (62.2)	
Pathology during pregnancy			
No	95 (27.7)	44 (28.2)	0.494
Yes	248 (72.3)	112 (71.8)	
Gestational hypertension			
No	228 (98.3)	148 (94.9)	0.037*
Yes	6 (1.7)	8 (5.1)	
Preeclampsia			
No	127 (95.1)	146 (93.6)	0.316
Yes	17 (4.9)	10 (6.4)	
Gestational diabetes			
No	339 (98.5)	149 (95.5)	0.045*
Yes	5 (1.5)	7 (4.5)	
Anemia during pregnancy			
No	216 (63)	108 (69.2)	0.104
Yes	127 (37)	48 (30.8)	
Type of delivery			
Vaginal	285 (82.8)	118 (75.6)	0.040*
Cesarean	59 (17.2)	38 (24.4)	
Labor complications*			
No	313 (91)	140 (89.7)	0.385
Yes	31 (9)	16 (10.3)	
Gestational age			
Preterm	45 (13.4)	23 (14.7)	0.393
Full term	297 (86.6)	133 (85.3)	
Newborn complications**			
No	245 (71.2)	108 (69.2)	0.703
Yes	98 (28.5)	48 (30.8)	

Data is presented as average ± standard deviation or n (%). * Hypertension, placental abruption, disseminated intravascular coagulation, Hellp syndrome, uterine atony/hemorrhage, pulmonary edema, hepatic failure, acute renal failure, ICU hospitalization, blood transfusion or maternal death. ** Prematurity, sepsis, infant respiratory distress syndrome, pathological jaundice, central cyanosis, neonatal tachypnea/apnea, shoulder dystocia or neonatal death.




Figure 1: Length at birth in offspring of adolescents with and without pregestational weight excess (overweight and Obesity).




Figure 2: Weight at birth in offspring of adolescents with and without pregestational weight excess (overweight and obesity).

Acknowledgments: We would like to express our deepest gratitude to the participants for their time and patience throughout this study.

Financial disclosure: The authors have no financial relationships relevant to this article to disclose.

Conflict of interest: The authors have no conflicts of interest to disclose.

Contributor statement

JV, EGV, RCS, VLB conceived/ designed the study and worked on data collection. CA and MV worked on data collection. MLOR, JV, VLB and RCS carried out the initial analyses and drafted the initial manuscript. AVP and MS finalized the manuscript, and MS translated it. All authors critically reviewed and revised the manuscript. All authors approved the final manuscript as submitted.

REFERENCES RÉFÉRENCES REFERENCIAS

1. SISTEMA DE INFORMAÇÃO SOBRE NASCIDOS VIVOS. Nascidos vivos brasil. Disponível em: <<http://tabnet.datasus.gov.br/cgi/tabcgi.exe?sinasc/cnv/nvuf.def>>. Acesso em: 10 jul. 2017.
2. Martins LWF, Frizzo GB, Diehl AMP. A constelação da maternidade na gestação adolescente: um estudo de casos. 2014; 25(3): 294-306.
3. Santos NL de AC, Costa COM, Amaral MTR, Vieira GO, Bacelar EB *et al.* Gravidez na adolescência: análise de fatores de risco para baixo peso, prematuridade e cesariana. Ciênc. saúde coletiva. 2014; 19(3): 719-26.
4. Nomura RMY, Paiva LV, Costa VN, Liao AW, Zugaib M. Influência do estado nutricional materno, ganho de peso e consumo energético sobre o crescimento fetal, em gestações de alto risco. Rev. Bras. Ginecol. Obstet. 2012; 34(3): 107-12.
5. Santos MMA de S, Baião MR, Barros DC de, Pinto A de A, Pedrosa PLM *et al.* Estado nutricional pré-gestacional, ganho de peso materno, condições da assistência pré-natal e desfechos perinatais adversos entre puérperas adolescentes. Rev Bras de Epidemiol. 2012;15(1): 143-54.
6. BRASIL, Ministério da Saúde. Série Técnica Redes Integradas a Atenção à Saúde. Perspectivas e desafios no cuidado às pessoas com obesidade no SUS: resultados do laboratório de inovação no manejo da obesidade nas redes de Atenção a Saúde 1.ed. Brasília, 2016.
7. INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA. Pesquisa de orçamentos familiares, 2010. Disponível em: <<http://www.ibge.gov.br/home/presidencia/noticias/imprensa/ppts/0000000108.pdf>>. Acesso em: 17 mar. 2017.
8. Raposo L, Ferreira C, Fernandes M, Pereira S, Moura P. Complicações da Obesidade na Gravidez. Arq Med Porto. 2011; 25(3): 115-22.
9. Oliveira CA de, Lins CP, Sá RAM de, Netto HC, Bornia RG *et al.* Síndromes hipertensivas da gestação e repercussões perinatais. Rev Bras Saúde Materno Infantil. 2006; 6(1): 93-98.
10. Silva JC, Amaral AR do, Ferreira B da S, Willeman IKM, Silva MR *et al.* Obesidade materna e suas consequências na gestação e no parto: uma revisão sistemática. FEMINA. 2014; 42(3): 135-40.
11. Gonçalves CV, Sassi AM, Cesar JÁ, Castro NB de, Bortolomedi AP. Índice de massa corporal e ganho de peso gestacional como fatores preditores de complicações e do desfecho da gravidez. Rev Bras Ginecol Obstet. 2012; 34(7):304-9.
12. MARTINS, M.G. *et al.* Associação de gravidez na adolescência e prematuridade. Rev Bras GinecolObstet, v.33, n.11, p. 354-360, 2011.
13. Gama, S.G.N. *et al.* Fatores associados à cesariana entre primíparas adolescentes no Brasil, 2011-2012. Cad Saúde Pública, v.30, supl.1, p.117-127, 2014.
14. WORLD HEALTH ORGANIZATION. Growth reference data for 5-19 years, 2007. Disponível em: <<http://www.who.int/growthref/en/>>. Acesso em: 23 mar.2017.
15. Atalah SE, Castillo CL, Castro RS. Propuesta de un nuevo estandar de evaluacion nutricional em embarazadas. Rev Med Chile 1997; 125: 1429-36.
16. Institute of Medicine. Weight gain during pregnancy: reexamining the guidelines. Washington, D.C: The National Academies Press; 2009.
17. Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Ações Programáticas Estratégicas. Área Técnica de Saúde da Mulher. Pré-natal e Puerpério: atenção qualificada e humanizada. Brasília-DF, 2006.
18. Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. Obstetrics and gynecology. 2013;122(5):1122-31.Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. Obstetrics and gynecology. 2013;122(5):1122-31.
19. Ganchimeg T, Ota E, Morisaki N, Laopaiboon M, Lumbiganon P, Zhang J, *et al.* Pregnancy and childbirth outcomes among adolescent mothers: a World Health Organization multicountry study. BJOG 2014; 121(Suppl 1):40–8.
20. Restrepo-Mesa SL, Zapata Lopez N, Parra Sosa BE, *et al.* Embarazo adolescente: características maternas y su asociacion con el peso al nacer del neonato. Arch Latinoam Nutr 2014; 64:99
21. Hedley A, Ogden C, Johnson C, Curtin L, Flegal K. Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. JAMA 2004;291:2847-50.
22. Freedman, D.S., Khan, L.K., Dietz, W.H., Srinivasan, S.R., & Berenson, G.S. (2001). Relationship of childhood obesity to coronary heart disease risk factors in adulthood: The Bogalusa heart study. Pediatrics, 108, 712–718.
23. Silva A de AA, Coutinho IC, Katz L, Souza ASR. Fatores associados à recorrência da gravidez na adolescência em uma maternidade escola: estudo

- caso-controle. *Cad Saúde Pública*.2013; 29(3):496-506.
24. Patias ND e Dias ACG. Sexarca, informação e uso de métodos contraceptivos: comparação entre adolescentes. *Psico-USF*. 2014; 19(1): 13-22.
 25. INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA. Pesquisa Nacional de Saúde do Escolar, 2015. Disponível em: <<http://biblioteca.ibge.gov.br/visualizacao/livros/liv97870.pdf>>. Acesso em 08 jul.2017.
 26. Nielsen, J.N., O'Brien, K.O., Witter, F.R., Chang, S.C., Mancini, J., Nathanson, M.S., et al. (2006). High gestational weight gain does not improve birth weight in a cohort of African American adolescents. *American Journal of Clinical Nutrition*, 84, 183– 189.
 27. Barros DC. Avaliação nutricional antropométrica de gestantes adolescentes no município do Rio de Janeiro. 2009. 129f. Tese (doutorado). Fundação Oswaldo Cruz, Escola Nacional de Saúde Pública Sergio Arouca, 2009.
 28. Vitalle MS de S, Tomioka CY, Juliano Y, Amancio OMS. Índice de massa corporal desenvolvimento puberal e sua relação com a menarca. *Rev Assoc Med Bras*. 2003; 49(4): 429-33.
 29. Pinheiro M de M, Oliveira JS, Leal VS, Lira PIC de, Souza NP et al. Prevalência do excesso de peso e fatores associados em mulheres em idade reprodutiva no Nordeste do Brasil. *Rev Nutr*.2016; 29(5):679-689.
 30. Bubach S. Idade da menarca, composição corporal e fatores de risco cardiometabólicos em mulheres adultas jovens: análises nas Coortes de Nascimento de Pelotas de 1982 e 1993. 2017. 203 f. Tese (Doutorado) – Universidade Federal de Pelotas, 2017.
 31. Castilho SD, Pinheiro CD, Bento CA, Barros-Filho A de A, Cocetti M. Tendência secular da idade da menarca avaliada em relação ao índice de massa corporal. *Arq Bras Endocrinol Metabol*. 2012; 56: 195-200.
 32. BRASIL, Ministério da Saúde. Instituto Sírio-Libanês de Ensino e Pesquisa. Protocolos da Atenção Básica: Saúde das Mulheres. Brasília, 2016.
 33. Fonseca MRCC da, Laurenti R, Marin CR, Traldi MC. Ganho de peso gestacional e peso ao nascer do concepto: estudo transversal na região de Jundiaí, São Paulo, Brasil. *Ciência e Saúde Coletiva*. 2014;19(5):1401-1407.
 34. Magalhães EI da S, Maia DS, Bonfim CFA, Netto MP, Lamounier JA, Rocha D da S. Prevalência e fatores associados ao ganho de peso gestacional excessivo em unidades de saúde do sudoeste da Bahia. *Rev bras epidemiol*. 2015; 18(4): 858-869.
 35. Schieve LA, Cogswell ME, Scanlon KS et al (2000) Prepregnancy body mass index and pregnancy weight gain: associations with preterm delivery. *Obstet Gynecol* 96:194–200 15 - 12. Abbey B, Wiemann CM, Rowe TF, Rickert VI.
 36. Sukalich S, Mingione M, Glantz C. Obstetric outcomes in overweight and obese adolescents. *Am J Obst Gyn* 2006;195: 851–5.
 37. Gunderson, E.P., Murtaugh, M.A., Lewis, C.E., Quesenberry, C.P., West, D.S., & Sidney, S. (2004). Excess gains in weight and waist circumference associated with childbearing: The Coronary Artery Risk Development in Young Adults Study (CARDIA). *International Journal of Obesity & Related Metabolic Disorders: Journal of the International Association for the Study of Obesity*, 28, 525–535.
 38. Groth SW. The long term impact of adolescent gestational weight gain. *Res Nurs Health* 2008;31:108–18.
 39. Stevens-Simon, C., & McAnarney, E.R. (1992). Adolescent pregnancy. Gestational weight gain and maternal and infant outcomes. *American Journal of Diseases of Children*, 146, 1359–1364.
 40. Fazio E de S, Nomura RMY, Dias MCG, Zugaib M. Consumo dietético de gestantes e ganho ponderal materno após aconselhamento nutricional. *Rev Bras Ginecol Obstet*. 2011; 33(2): 87-92.
 41. Carvalhaes MA de BL, Gomes C de B, Malta MB, Papini SJ, Parada CMG de L. Sobrepeso pré-gestacional associa-se a ganho ponderal excessivo na gestação. *Rev Bras Ginecol Obstet*. 2013; 35(11):523-9.
 42. Haeri S, Guichard I, Baker AM, Saddlemire S, Boggess KA. The Effect of Teenage Maternal Obesity on Perinatal Outcomes. *American College of Obstetricians and Gynecologists*. 2009; 113(2):300-304.
 43. Seligman LC, Duncan BB, Branchtein L, Gaio DSM, Mengue SS et al. Obesity and gestational weight gain: cesarean delivery and labor complications. *Rev Saúde Pública*. 2006; 40(3):457-65.
 44. BRASIL. Ministério da Saúde. Rede Interagencial de Informações para a Saúde. DATASUS [Internet] (BR). Indicadores e Dados Básicos – Brasil. 2011. Disponível em <<http://tabnet.datasus.gov.br/cgi/tabcgi.exe?ldb2012/f08.def>>. Acesso em 23. Jun. 2017.
 45. Assunção PL de, Melo AS de O, Gondim SSR, Benício MHD, Amorim MMR et al. Ganho ponderal e desfechos gestacionais em mulheres atendidas pelo Programa de Saúde da Família em Campina Grande, PB (Brasil). *Revista Brasileira de Epidemiologia*.2007; 10(3): 352-360.
 46. Gaillard R, Durmus B, Hofman A, Mackenbach JP, Steegers EAP et al. Risk Factors and Outcomes of Maternal Obesity and Excessive Weight Gain During Pregnancy. *Obesity*.2013; 21(5): 1046-55.
 47. Ma RC, Schmidt MI, Tam WH, McIntyre HD, Catalano PM. Clinical management of pregnancy in

the obese mother: before conception, during pregnancy, and post partum. 2016.

48. Kac G, Meléndez GV. Ganho de peso gestacional e macrosomia em uma coorte de mães e filhos. *J Pediatr.* 2005;81(1): 47-53.
49. Li N, Liu E, Guo J, Pan L, Li B et al. Maternal Prepregnancy Body Mass Index and Gestational Weight Gain on Pregnancy Outcomes. *PloS one.* 2013; 8(12): 1-7.

