

Results of Salt Cognition Test using SALSARE since 67 Female University Students

Naomi Katayama, Akemi Ito, Mayumi Hirabayashi¹

¹ Nagoya Women's University

Received: 16 December 2019 Accepted: 31 December 2019 Published: 15 January 2020

Abstract

In Japan, salt reduction is encouraged to prevent high blood pressure. However, it is difficult to reduce salt, and good salt intake per day did not fail to the target value. In this study, we report a saltiness cognitive threshold test using female university students. The participant is 67 female university students. Participants were subjected to a salty cognitive threshold test using SALSARE (manufactured by Advantech). The saltiness test started from a light taste and tried a strong taste in order. The salt concentration is 0.6

Index terms— saltiness test, cognition, threshold, salsare, university student.

1 Introduction

Japanese Dietary Intake Standards in the 2020 version, the daily intake of salt for females is recommended to be 6.5g or less. Salt intake is decreasing with each revision of dietary standards. In the near future, it is expected that salt intake in Japanese dietary intake standards will lower to the international standard of 6.0g. By reducing salt, it is expected to prevent illness from various diseases (high blood pressure, kidney disease, heart disease, etc.). Therefore, the purpose of this study was to conduct a salt concentration cognition test on Japanese people to understand the actual condition of the cognitive threshold for salt, and to use it as future data. To begin with, we report that a healthy female university student underwent a salt concentration recognition test.

2 II.

3 Materials and Methods

4 a) Participants

The participant is 67 female university students. Their average age $\pm$ standard deviation (SD) was 29.6 ± 0.6 , average height $\pm$ SD was 158.9 ± 5.7 cm, and average weight $\pm$ SD was 50.3 ± 4.5 kg (Table 1). This study conducted with the approval of the Ethics Committee (Nagoya women's university 'hitowo mochiita kennkyuuni kansuru iinnkai'). The approval number is 30-14.

5 III.

6 Results

7 a) Saltiness recognition test result

Sixty-seven female university students underwent a saltiness cognitive threshold test using SALSARE. As a result, 62 out of 67 female university students who felt salty at a concentration of 0.6% were 92% of all participants. And, three female university students felt a salt concentration of 0.8%, were 4.5% of all participants. However, two female university students did not feel taste even with a salt concentration of 1.6%. This is 3% of all participants (Table3). Table 4 shows the results of the questionnaire survey. When asked if they had enough saliva, most

female university students replied that they had enough or normal secretions. None of the female university students answered that they had none or no saliva. In response to the question of feeling the taste, most female university students answered that they could well or understand the taste. None of the female university students answered that they didn't understand the question of whether they would feel the taste. When asked about the frequency of use of restaurants and groceries, female university students answered that they would use 2-3 times a week or once a week. When asked if the usual seasoning for food was salty or not too much salty, female university students answered that they were the lightly salted.

8 Discussion

Many researchers have reported the relationship between salt intake and blood pressure 1) and the relationship between hypertension 2,3,4) . There are also reports on diet and salt intake in the younger generation 5) . We also reported the results of the saltiness cognition threshold test conducted by Yakumo Study on the elderly 6,7) . There are also reports on the relationship between salt intake and blood pressure overseas 8,9) , as well as educational effects on salt intake 10,11) . Currently, many new drugs for lowering blood pressure have developed for people with high blood pressure 12) , but dietary food habits also desirable can be improved their blood pressure. The drug also has side effects and that costs a lot to keep taking it. In Alzheimer's dementia, the taste is not known, and the value of the salty cognitive threshold test also deteriorates 13) . Therefore, it is highly likely that the dietary intake will be high, which may raise blood pressure. A low saltiness recognition threshold means that the saltiness of the meal can be reduced. Keeping the salty cognitive threshold low makes sense for a healthy diet. We conducted a saltiness cognition threshold test on 67 young female university students. As a result, they recognized 0.6% saltiness, and with included 0.8% recognized students, the total number of recognized was 96.5%. The result of the dietary habit questionnaire showed that saliva was well secreted, the taste of the meal was well understood, and the dietary intake was light. However, two female university students did not feel taste even with a salt concentration of 1.6%. This is 3% of all participants.

In the future, we would like to conduct a more detailed questionnaire survey on dietary intake and compare it with the salty cognitive threshold test results.

V.

9 Conclusions

The results of this test show that most female university students (92%) feel salty at 0.6%. Also, female university students were eating a light taste in their daily life. Moreover, the use of restaurants and the use of commercially available foods were once or 2-3 times a week. The effect of the salt reduction awareness campaign in Japan, which has continued for over 20 years, is good for the younger generation. However, on the other hand, among the female university students who participated in this study, two students could not recognize 1.6% saltiness, so it is necessary to investigate the dietary habits of the participants in more detail.

1

b) Assessment of salt taste identification

Participants were subjected to a salty cognitive threshold test using SOLSAVE (manufactured by Advantech). The saltiness test started from a light taste and tried a strong taste in order. The saltiness test starts

Figure 1: Table 1 :

2

c) Ethical review board

Figure 2: Table 2 :

3

[Note: b) Questionnaire results]

Figure 3: Table 3 :

4

Year 2020
2

Figure 4: Table 4 :

Average	1.3	1.5	3.8	2.3
??	0.7	0.5	1.0	0.7
IV.				

Figure 5: Saliva secretion Taste perception Use of restaurants and commercial food Favorite food salt taste

.1 Acknowledgements

This study was supported by the research aid of Choju-iryo-kenkyu-kaihatsuhi 30-14 and the Japanese Society of Taste Technology, 2019.

[Malaga et al. ()] , S Malaga , J J Diaz , J Arguelles , C Perillan , I Malaga , M Vijande . *Pediatric nephrology* 2003. 18 (5) .

[Olayemi and Mabadeja ()] *Comparative study of salt taste threshold of jypertensives, their normotensive relatives and non-relatives. The Nigerian postgraduate medical journal*, S O Olayemi , A F Mabadeja . 2003. 10.

[Cho et al. ()] ‘Comparison of salt taste thresholds and salt usage behaviours between asults in Myanmmar and Korea’. H Cho , S M Kinn , Jeong Ss , S B Kim . *Asia Pacific journal of clinical nutrition* 2016. 25 (4) .

[Ferrante et al. ()] ‘Feasibility of salt reduction in prcessed foods in Argentina’. D Ferrante , N Apro , V Ferreira , M Virgolini , V Aguilar , M Sosa , P Perel , J Casas . *Pan American journal of public health* 2011. 29 (2) .

[Volkov et al. ()] ‘Impsired gustatory sensitivity of the tongue to table salt as a risk factor of arterial hypertension’. V S Volkov , O B Poseliugina , S A Nilova , T S Vinogradova , Rokkina Sa , O P Svistunov . *Klinicheskai meditsina* 2010. 88 (1) .

[Katayama et al. ()] ‘Odour and Salt Taste Identification in Older Adults: Evidence from the Yakumo Study in August’. N Katayama , S Kondo , H Ootake . *Acade. J. Med. Plants* 2018. 2018. 7 (3) p. .

[Petrova et al. ()] T S Petrova , N D Bazhenov , Mazur Vv , E S Mazur . *Gustatory sensitivity threshold to table salt*, 2012.

[Nishimoto et al. ()] *Reproducibility of taste examination with Salsave: Control study for healthy volunteers. Japan Society of Stomato-pharyngology*, K Nishimoto , J Ohhori , T Shimomugi , Y Kurono . 2005. p. .

[Okoro et al. ()] ‘Salt taste sensitivity and blood pressure in adolescent school children in southern Nigeria’. E O Okoro , G E Uroghide , E T Jolayemi . *East African medical journal* 1998. 75 (4) .

[Elias et al. ()] ‘Salt-sensitivity in normotensive and hypertensive Nigerians’. S O Elias , E C Azinge , G A Umoren , Jaja St , O A Sofola . *Nigerian quarterly journal of hospital medicine* 2011. 21 (1) .

[Kusabe et al. ()] ‘Sodium restriction improves the gustatory threshold for salty taste in patients with choric kidney sisease’. U Kusabe , Y Mori , M Okagaki , H Neriya , T Adachi , C Sugishita , K Aonomura , T Kimura , N Kishimoto , H Nakagawa , M Okigaki , T Hatta , H Matsubara . *Kidney international* 2009. (6) p. 76.

[Volos et al. ()] ‘Taste receptor response to sodium chloride and naturiuresis in the workers and employees of an industrial enterprise’. B E Volos , ' Vasil , M Lu , L V Masliaeva , I A Snegurskaia . *Likars'ka sprava* 1994. p. .