Artificial Intelligence formulated this projection for compatibility purposes from the original article published at Global Journals. However, this technology is currently in beta. *Therefore, kindly ignore odd layouts, missed formulae, text, tables, or figures.*

CrossRef DOI of original article: 10.34257/GJMRKVOL20IS13PG1

1	Comparison of ATP Values on Meat and Fish Cutting Boards
2	before and after Alcohol Disinfection
3 4	Akemi Ito ¹ , Naomi Katayama ² , Mayumi Hirabayashi ³ , Natuki Sasaki ⁴ and Moe $\rm Inuzuka^5$
5	¹ Nagoya Women?s University
6	Received: 9 December 2019 Accepted: 3 January 2020 Published: 15 January 2020
7	

8 Abstract

- ⁹ Sanitary control of cutting boards in the kitchen is important to prevent food poisoning.
- ¹⁰ Using ATP and microbiological tests, we investigated the cleaning and 70
- 11

12 Index terms— Gender: ATP wiping test, Microbial stamp test, Cutting board, alcohol disinfection.

13 **1** Introduction

anitary control of cutting boards in the kitchen is important to prevent food poisoning. In the past, we reported 14 the results of hygiene management by repeatedly cleaning the cutting board with detergent and running water for 15 30 seconds or more 1). Currently, the COVID-19 epidemic requires stricter hygiene control. To control invisible 16 microorganisms, it is necessary to take measures to avoid the risk of food poisoning accidents due to familiarity 17 with cooking work; as the O-JT education, it is necessary to create a hygiene management manual and protect 18 it with all the cooks 2,3,4,5). However, if the procedure is complicated and difficult, it will not last long. We 19 need easy and reliable procedures and methods that anyone can do. The ATP tests 6,7,8) and HACCP-based 20 microbiological tests 9) are useful in hygiene management to transform invisible bacteria into visible forms and 21 educate them. Therefore, in this study, and the cutting board cleaning method we performed last time, a step of 22 spraying 70% alcohol added. The effects of this alcohol disinfection compared by adding a stamping test (General 23 bacteria, Staphylococcus aureus, Escherichia coli, Salmonella, Vibrio parahaemolyticus) in addition to the same 24 ATP test as in the previous report. 25

²⁶ **2 II.**

²⁷ **3** Materials and Methods

²⁸ 4 a) Kitchen cutting board

The six kitchen meat or fish thick cutting board (cutting board 1) and the six kitchen meat for the fish thin cutting board (cutting board 2) prepared in the kitchen were stored in the sterilization storage the day before cooking.

³² 5 b) ATP inspection procedure

33 Each of the 12 cooks carried a kitchen cutting board for meat or fish at the start of their work and brought it 34 to the cooking table. The work start time depends on the working conditions of the cooks. Still, the inspector 35 always performed an ATP inspection before using meat or fish with a kitchen cutting board. Then, each cook finished the work, washes the cutting board firmly with detergent and sponge, rinse with running water for 30 36 seconds or more. Then, each cook repeated this process twice (as same as the last report 1)). The inspector 37 performed an ATP inspection after using meat or fish with a kitchen cutting board, again. Then, each cook 38 sprayed 70% alcohol on the cutting board after washing. At last, the inspector performed an ATP inspection 39 after using meat or fish with a kitchen cutting board. The ATP test kit used manufactured by KIKKOMAN. 40

⁴¹ 6 c) Stamp test inspection procedure

Five types of stamp test (General bacteria, Staphylococcus aureus, Escherichia coli, Salmonella, Vibrio parahaemolyticus) used. The stamp test conducted by the inspector at the same time as the ATP. The stamp test was colony-counted after culturing in an incubator at 38 degrees for three days. The stamp test made by NISSUI.

⁴⁶ 7 d) Statistical processing

The results obtained compared using statistical methods. The data were statistically processed, was subjected to an F test to determine whether to use a parametric test or nonparametric test. When there is no difference in the F test, the presence or absence of a significant difference was confirmed using the student ttest with or without a correspondence. If there was a difference in the F test, the presence or absence of a significant difference was confirmed using the Wilcoxon test with a pair or the Mann-Whitney test without correlation.

52 **8 III.**

53 9 Results

⁵⁴ 10 a) ATP value results before and after alcohol disinfection

The table ?? and 2 shows the results of ATP wiping tests on cutting board before and after alcohol disinfection. It can see that the average value of the ATP values measured after washing before and after cleaning, after cleaning, the ATP value is low. The ATP value after 70% alcohol spraying was statistically significantly lower than that before alcohol spraying. The ATP value dropped below 100 for both cutting boards.

⁵⁹ 11 b) Stamp test results before and after alcohol disinfection

Tables 3,4,5,6,7,8.9.10.11 and 12 show the results of ATP wiping tests on cutting board before and after 70% alcohol disinfection. Results of general bacteria show in Tables 3 and 4. Results of E. coli show in Tables 5 and 6. Results of Staphylococcus aureus shown in Tables 7 and 8. Result of Salmonella show in Tables 9 and 10. Result of Vibrio parahaemolyticus show in Tables 11 and 12. The number of all microbial bacteria was lower after washing than after cooking and after spraying 70% alcohol. However, there was no statistically significant difference in the number of microbial bacteria.

66 12 Discussion

To manage the hygiene of meat and fish cutting board that has a high risk of causing secondary contamination in 67 cooking. We tried to verify using the ATP test and microbial stamp test by spraying 70% alcohol after cleaning 68 instead of controlling only by the cleaning method 1). The ATP value decreased after washing then after cooking 69 70 and after spraying 70% alcohol than after washing. The ATP value was a statistically significant decrease, which 71 was less than 100 after 70% alcohol spraying. However, the microbial stamp test results were not statistically 72 significant reductions in bacterial counts. The cutting board inspected by spraying 70% alcohol after cleaning. 73 But if 70% of alcohol not sprayed after sufficiently wiping off the water, the alcohol may be dilute, and the bactericidal effect may weakened. In the future, we would like to verify the sterilization of microorganisms by 74 spraying 70% alcohol on the cutting board by thoroughly wiping off the water after cleaning and then spraying 75 70% alcohol. Not all microorganisms are killed even after spraying 70% alcohol, so when using a cutting board 76 left at room temperature (with moist), it is better to wash repeatedly and cook after spraying 70% alcohol. 77 V. 78

79 13 Conclusions

The effects of 70% alcohol spraying investigated using cutting boards for meat and fish. Both cutting boards 80 had high ATP and microbiological test values after cooking. However, although the value of the cutting board 81 decreased after cleaning, the ATP value did not fall below 100. Microbial test values were also high in many 82 cases. After spraying with 70% alcohol, the ATP value was 100 or less, and the value decreased statistically 83 significantly. Microbial test values were decreasing with or without statistically significant reductions. Providing 84 safe and secure meals by further spraying 70% alcohol after cleaning the cooking utensils helps prevent food 85 poisoning. However, since the microorganisms are present even after spraying with 70% alcohol, the bacteria 86 may grow again if the cooking utensils left for a long time. It is advisable to clean and spray 70% alcohol before 87 using the equipment. 88

 $^{^{1}}$ © 2020 Global Journals

	Table 1. ATP test value and statistical pro	cossing result of cutting	r hoor	1 1
	Table1. ATF test value and statistical pro	No alcohol treatmen	-	Alcohol treatment
	For meat	Before washing	After	After wash- ing
	1	8414	50121	50121
Year	2 3 4 5 6 Average value	$210 \ 132205 \ 59141$	50121 56	56
2020		30814 76010 51132.3	103 62 272 70 8447.3	103 62 272 70 8447.3
	??	49166.2	20416	6.0 0416.0
2	F test Student-t [*] neWilcoxon F test Student-t [*] Wilcoxon For	P-0.025*		P=0.0001**
Is- sue XIII Ver- sion I				
DD DD	Student-t [*] Wilcoxon	P=0.028*		P=0.173
(F test		P=0.0	0001**
Medic Re- search	caStudent-t [*] Wilcoxon	*Paired Student-t te	st * P<	<0.05, ** P<0.01 P=0.028*
	l For meat 1 2	Before washing After	r washi	ing After washing 82 40 40 4
01	3	200	46	46
	4	13	0	0
	5	200	9	9
	6	60	0	0
	Average value	93.2	15.8	15.8
	??	87.7	21.4	21.4
	F test	P=0.002**		P=0.003**
	Student-t*	D 0.020*		
	Wilcoxon	P=0.028*		D 0 0001 **
	F test			P=0.0001**
	Student-t* Wilcoxon			P=0.028*
	W IICOXOII	*Paired Student-t te	st * P∢	

Figure 1: Table 3

3

	No alcohol treatment P=0.0001 ^{**} P=0.116	P=0.0001** P=0.075	Alcohol treatme $P=0.035^*$ $P=0.028^*$	ent
	*Paired Student-t test * $P < 0$		1	
For meat	Before washing	After washing	After washing	After alco- hol
1	0	0	0	0
2	2	1	1	0
3	200	8	8	0
4	5	0	0	7
5	200	0	0	0
6	6	0	0	16
Average value	68.8	1.5	1.5	3.8
??	101.6	3.2	3.2	6.6
F test				
Student-t*				
Wilcoxon F test				
r test Student-t*				
Wilcoxon				

Figure 2: Table 4

	No alcohol treatment P=0.0001**		Alcohol treatme P=0.052 P=0.518	ent
	P=0.043*			
		P=0.0001**		
		P=0.418		
	*Paired Student-t test * $P < 0$			
For meat	Before washing	After	After washing	After
		washing		alco-
				hol
1	3	2	2	0
2	0	0	0	1
3	200	23	23	0
4	0	0	0	0
5	15	0	0	0
6	35	0	0	0
Average value	42.2	4.2	4.2	0.2
??	78.5	9.3	9.3	0.4
F test				
Student-t*				
Wilcoxon				
F test				
Student-t*				
Wilcoxon				

Figure 3: Table 5

5

	No alcohol treatment P=0.0001** P=0.068	P=0.0001** P=0.080	Alcohol treatme P=0.0001** P=0.285	ent
	*Paired Student-t test * P<0	0.05, ** P<0.0)1	
For meat	Before washing	After washing	After washing	After alco- hol
1	0	0	0	0
2	5	1	1	0
3	0	0	0	7
4	0	0	0	3
5	152	2	2	0
6	212	0	0	0
Average value	61.5	0.5	0.5	1.7
??	95.3	0.8	0.8	2.9
F test				
Student-t*				
Wilcoxon				
F test				
Student-t*				
Wilcoxon				

Figure 4: Table 6

 $\mathbf{7}$

Year 2020 3 Volume XX Issue XIII Version I $\mathbf{D} \ \mathbf{D} \ \mathbf{D} \ \mathbf{D} \ \mathbf{D}$) K (Medical Research Global Journal of result No alcohol treatment and statistical processing Alcohol treatment $P=0.0001^{**}$ $P=0.004^{**}$ P = 0.109P = 0.465P=0.0001** P = 0.345*Paired Student-t test * P < 0.05, ** P < 0.01

Figure 5: Table 7

1	

	No alcohol treatment P=0.0001** P=0.225	P=0.0001** P=0.144	Alcohol treatme P=0.0001** P=0.655	ent
	*Paired Student-t test * $P < 0$	05, ** P<0.0	1	
For meat	Before washing	After washing	After washing	After alco- hol
1	508	0	0	0
2	80	0	0	0
3	1	0	0	2
4	168	0	0	5
5	1	9	9	0
6	26	15	15	5
Average value	130.7	4.0	4.0	2.0
??	195.4	6.5	6.5	2.4
F test				
Student-t*				
Wilcoxon				
F test				
$Student-t^*$				
Wilcoxon				

Figure 6: Table 8

	No alcohol treatment P=0.0001**	P=0.075	Alcohol treatment P=0.016* P=.0001** P=0.059	P=0.465
	*Paired Student-t test * P	,		A C.
For meat	Before washing	After	After washing	After
		washing		alcohol
1	0	1	1	0
2	0	5	5	0
3	34	1	1	2
4	0	23	23	0
5	21	1	1	0
6	55	0	0	0
Average value	18.3	5.2	5.2	0.3
??	22.8	8.9	8.9	0.8
F test				
Student-t*				
Wilcoxon				
F test				
Student-t*				
Wilcoxon				
W IICOXOII				

Figure 7: Table 9

	No alcohol treatment P=0.019 P=0.463	$P=0.0001^{**}$ P=0.109	Alcohol treatme P=0.0001** P=0.138	ent
	*Paired Student-t test * P<	<0.05, ** P<0.	01	
For meat	Before washing	After washing	After washing	After alco- hol
1	21	0	0	0
2	6	0	0	0
3	1	7	7	0
4	119	0	0	0
5	0	38	38	2
6	18	0	0	0
Average value	27.5	7.5	7.5	0.3
??	45.7	15.2	15.2	0.8
F test				
$Student-t^*$				
Wilcoxon				
F test				
Student-t*				
Wilcoxon				

Figure 8: Table 10

No alcohol treatment P=0.009** P=0.402 P=0.0001** P=0.075 *Paired Student-t test * P<0.05, ** P<0.01 Alcohol treatment $P=0.0001^{**}$ P=0.180

	No alcohol treatment		Alcohol treatm	ent
For meat	Before washing	After	After washing	After al-
		washing		cohol
1	0	0	0	0
2	0	4	4	2
3	5	0	0	0
4	0	0	0	0
5	200	0	0	0
6	256	0	0	0
Average value	76.8	0.7	0.7	0.3
??	118.4	1.6	1.6	0.8
F test	P=0.0001**		P = 0.68	
$Student-t^*$			P = 0.363	
Wilcoxon	P=0.144			
F test		P=0.0001	**	
$Student-t^*$	P=0.144 *Paired Student-t tes	t * $P < 0.05$,	** P<0.01	Year
Wilcoxon				2020
				5

Figure 10: Table 12

Figure 11: K

⁸⁹ .1 Acknowledgments

- ⁹⁰ We would like to thank all the cooks who participated in this experiment. Also, we would like to thank the ⁹¹ inspectors who also performed the ATP inspection.
- ⁹² [Stanley ()] 'A review of bioluminescent ATP techniques in papid microbiology'. P E Stanley . J Biolumin
 ⁹³ Chemilumin 1989. 4 (1) p. .
- 94 [Griffith et al. ()] 'An evaluation of hospital cleaning refimes and standards'. C J Griffith , R A Coooper , J 95 Gilmore , C Davies , M Lweis . J Hosp Infect 2000. 45 (1) p. .
- ⁹⁶ [Lee ()] 'An investigation of Factors that influence Hygiene Practices at a small Day Care Center'. J H Lee . J
 ⁹⁷ Food Prot 2018. 2018. 81 (1) p. .
- 98 [Aycieck et al.] Comparison of results of ATP bioluminescence and traditional ygiene swabbing methods fro the
 99 deteminaton of surface, H Aycieck, U Oquz, K Karci.
- [Nante et al. ()] 'Effectiveness of ATP bioluminescence to assess hospital cleaning: a review'. N Nante , E Ceriale
 , G Messina , D Lenzi , P Manzi . J Prev. Med. Hyg 2017. 58 (2) p. .
- [Stannard and Gibbs ()] 'Rapid microbiology: application s of bioluminescence in the food industry-a review'. C
 J Stannard , P A Gibbs . J Biolumin Chemilumin 1986. 1 (1) p. .
- 104 [Katayama et al. ()] 'Results of Hygiene Education of Kitchen Cutting Board by using ATP Inspection -
- Comparison of vegetable Cutting Board and Meat Cutting Board'. N Katayama, M Hirabayashi, A Ito
 , S Kondo, Y Nakayama, A Naka, N Sasaki, M Inuzuka, T Tamura. *Global Journal of Medical Research*2020. 20 (5) p. .
- [Amodio and Dubi ()] 'Use of ATP bioluminescence for assessing h eclealiness of hospital surfaces: a review of
 the published literature'. E Amodio , C Dubi . J infect Public Health 1990-2012. 2014. 7 (2) p. .