

Prospective Promising Signs of Herd Immunity in COVID-19 Transmission Suppression Via Vaccination

Bayode, Michael Tosin

Received: 7 June 2021 Accepted: 2 July 2021 Published: 15 July 2021

Abstract

Reports of the COVID-19 pandemic show an elevated level of mortality among patients, with some inclining hazards distinguished as age, underlying infection conditions; hypertension, diabetes, and so on, immune-compromised conditions and viral dose during exposure. Different investigations portray an elevated level of super-spreading occasions, which proposes that heterogeneity in infectivity may altogether affect the elements of its transmission. This review is intended to make the perception herd immunity needs to play in COVID-19 transmission concealment inferable from its circumspect viability for the destruction of numerous maladies and indeed, give the premise to vaccines and their applications serving as a proviso for immune individuals that will prompt a huge decrease in disease event and spread. On account of the ebb and flow of the COVID-19 scourge, this may give the perfect viewpoint to totally eradicate the illness in our local communities in the event that a vaccine is before long evolved.

Index terms— COVID-19, transmission, herd immunity, vaccine, immune.

1 Introduction

As of now, the COVID-19 pandemic is a bane to the entire world populace, because of the pace of infectivity, inclined danger of acquiring the airborne viral disease and combined with the way that there is as of now no endorsed corrective control/vaccine for this viral infection. Arrays of investigations have set up that the general population most predisposed and powerless against the ailment are the older individuals, males, people with hidden sickness conditions including; hypertension, diabetic condition, immuno-weakened people, and people dependent on immunosuppressant use [1]. Equally, considering the already infected populace, the possibility of being asymptomatic could vary, contingent upon the conditions and populace, going from 5% to 80% [2]. Three likely method of spread have been portrayed through droplet transmission, express contact spread, via airborne transmission. By and by, of late the digestive tract has additionally been demonstrated as a likely method of spread dependent on abdominal clinical indications and manifestation of diarrhea brought about by the viral infection, just as the viral RNA saw in faeces [3]. Various elements may control the infection's transmission given the infectivity levels of the host, for example, sanitary behaviours including quarantine, hand washing, and appropriate routine care which are needed to soothe the affinity of the viral infection and infectivity. It has been definite that community wellbeing mediation that stifles over 60% of transmissions are productive to deal with the concealment levels of the COVID-19 pandemic [3].

The term herd immunity was initially utilized in 1923 by Topley and Wilson [4]. It subsequently gave the need to vaccines and their purposes, vaccine program expenditure efficacy-breakdown, and the concealment of ailments, for example, smallpox and different irresistible diseases like polio and diphtheria [5,6,7]. Herd immunity is additionally affected by elements, for example, populace immunity and the methods of the spread of causal agents [6,8].

Divergent investigations have broke down the helpfulness of vaccination projects to accomplish herd immunity in, and in this way shield the unvaccinated against different maladies [9]. The primary embodiments are the immunizations for cholera, hepatitis A, hepatitis B, human papillomavirus, haemophilus flu, meningococcal, flu, pneumococcal, polio, challenging hack, measles, chickenpox, rotavirus, and yellow fever [10,11,12,13] a few of

which are represented in Table 1. The ramifications of herd immunity on the viability of infection transmission were as of late observed in America with the Chikungunya and Zika infections, which caused a pandemic in an altogether weak populace. By and by, after the surge of these arboviruses, herd immunity confined their spread [14]. In like manner, the United Kingdom of late executed a plan for the COVID-19 pandemic that caused controversy, giving space for the infection to be transmitted in the populace before charging social seclusion to boost herd immunity [15]. Deriving the orderly model of group immunity as the resistance that a specific populace has against a disease, Fox and his partners itemized four conditions under which such resistance can happen [20]. Initially, the transmittable microbe must be traced, found and restricted to a solitary host. For extreme intense respiratory disorder Covid 2 (SARS-Cov-2), the central method of spread was zoonotic in nature, with its essential host being the bat. Notwithstanding, the intermediary host through which it was transmitted to people is conditional. Transmission is known to have happened because of express contact with the contaminated animals or via their secretory liquids. Viral RNA has additionally been built up in canines and felines living with COVID-19 positive people. However, it has not been uncovered that these creatures can pass on the infection to individuals [21]. Also, the spread must happen predominantly through direct contact. Up till now, it has been standard that the spread of COVID-19 by direct individual-to-individual contact by means of coughing, sneezing, and inward breath of polluted vaporizers and contact spread through the oral, nasal, and ocular membranes [22].

With specific accentuation on the data lacking on the immune reaction incited by COVID-19 in people, it's been difficult to set up the methods by which the immune system make a drawn out response that could fight the viral infection and deflect disease relapse. At last, group immunity is exploited if the populace procure a laid-back integration model. With this situation, everybody is defense-less against getting contaminated with COVID-19 bringing to the fore its inalienable indications. By and by, this plan of laidback blending will rely upon the preventive rules executed by every nation's administration overall which involves quarantine, seclusion, social separation, and fortified pre-emptive rules for in danger gatherings, for example, pregnant ladies, the old people, and youngsters [23]. Hypothetically, it is likely to achieve group immunity under the expressed speculations. In any case, by and by absence of intrinsic comprehension of the adaptive immune response and, in the non-existence of an appropriate vaccine, the moral limitations to achieve such immunity make this cure excessively deceptive to preliminary [24].

2 a) Creating herd immunity within populations

To make herd immunity inside a populace, the immunity achieved by immunization or natural infection must be turn away ahead spread of infections. For SARS-CoV-2, clinical signs are a helpless marker of transmissibility, as asymptomatic hosts can be exceptionally transmittable and add to the spread of an epidemic [25]. For microorganisms in which deep rooted immunity is prompted, similar to the case for measles immunization, herd immunity is significantly efficacious and can turn away microbe spread inside a populace. Nonetheless, this circumstance is sensibly atypical, as immunity for some infectious diseases, for example, pertussis and rotavirus, fade in the long haul [25]. Subsequently, herd immunity is less productive and discontinuous outbreaks can still surface. Eventually, if immunity is unpredictably spread in a populace, gatherings of disease-prone hosts that can frequently interact with one another may continue.

3 b) Herd immunity efficiency for COVID-19 transmission suppression

As indicated by the WHO, herd immunity is basically serviceable for communicable diseases [26]. For transferable infections, however, the immunocompetent people must not exclusively be protected yet C there is the need of likewise decreasing disease among immununo-suppressed hosts through herd safeguard [27]. Attributable to herd protection, various maladies can be eradicated without 100% vaccination presentation. For a valid example Haemophilus influenzae type B vaccine introduction of under 70% in Gambia was adequate to eliminate Hib disease, with equal findings in Navajo populaces [28,29].

For COVID-19, herd immunity, with its relatively inherent effect, probably won't facilitate the infection as the outcomes rely upon the case and adequacy of the control measures and the ability to bring at the same time mounting outbreaks under instant control when required [30]. Antibodies for this infection are not yet accessible, however existing chemotherapeutic synergistic utilization, for example, azithromycin (AZT), hydroxychloroquine sulfate, chloroquine phosphate has been viewed as likely valuable. Regardless, their handiness and safety are built up in extra investigations for this novel malady [31,32,33].

4 III.

5 Specifics of Herd Immunity and Covid-19

The current COVID-19 pandemic has produced over 81 million of clinically established cases and has claimed an absolute number of more than 1.7 million lives worldwide as of 31 st December, 2020. A few exploratory preliminaries to evaluate novel immunization varieties and medication reconstituting approaches for the prevention and fixing of COVID-19 disease are directly in progress. Regardless, it is inconclusive whether these preliminaries will produce valuable cure, and it is dubious how broad these examinations will take albeit a positive

assessment for any immunization preliminary is in any event 12 to year and a half. In the inaccessibility of an immunization, developing of COVID-19 herd immunity through common disease is still theoretically possible.

6 a) Epidemiological considerations for COVID-19 herd immunity

Inconsistency in method of spread between people may assume a key part in COVID-19 dispersal. Super-spreading occasions happen when conditions positive for high paces of transmission emerge. These occasions include a solitary index case infecting countless secondary contacts and are known to be significant in driving episodes of infectious diseases, including SARS, MERS, and measles [34]. Reports of COVID-19 super-spreading occasions have been recorded, recommending that heterogeneity in infectivity may essentially affect the elements of its transmission [35]. At long last, the variables that impact interindividual heterogeneity in COVID-19 vulnerability, clinical pathology, and disease result are not surely known.

7 b) Immunological antibody responses to COVID-19

Following contamination with COVID-19, perceptible IgM and IgG antibodies create inside days to weeks of symptom onset beginning in most infected people [36,37,38]. Why a few patients appear not to build up a humoral immune reaction, as reflected by recognizable antibodies, is dubious. Adding to this vulnerability is the indistinct connection between antibody reaction and clinical improvement. The observations from a small investigation of 9 patients with COVID-19 found that more prominent clinical severity delivered higher antibody titers [36]. Be that as it may, antibody recognition and higher titers have not generally been found to correspond with clinical improvement in COVID-19 [37,38]. Additionally, mild COVID-19 symptoms can resolve earlier detection by IgM and IgG antibodies, albeit distinguishable IgM and IgG antibodies have heralded decreases in COVID-19 viral loads [37,38]. What appears more certain is that viral load regularly peaks from the get-go in infection, and afterward declines as antibodies sprang up and antibody titers ascend over the resulting 2 to 3 weeks [37,38]. The stability of neutralizing antibodies (NAbs, principally IgG) against COVID-19 presently can't seem to be characterized; ingenuity as long as 40 days from symptom onset has been depicted [36].

8 IV. Applications of Herd Immunity in the Current Covid-19 Pandemic

As per various models applied to infectious diseases, herd immunity is mathematically-connected with the spread and infection inclinations of the virus [39,40] as displayed in the herd immunity network in Figure 1, which are the consequence of the relationship set up over a specific timeframe between the number of healthy subjects and those vulnerable to infection, the infected subjects that can no longer add to the transmission of the infection, infected subjects, and the normal or vaccine-immune subjects in a populace, mediated by the infectiousness of the virus, the incubation time frame, the transmissibility period, the virus limit between individuals, the elements of contact among the populace, and the term of the viral disease [41]. Hardly, any investigations describe children as key cases in familial groups [42,43], and plainly, children all the more regularly have asymptomatic diseases when contrasted and grown-ups [33,43,44]. It is of dire significance in demonstrating the pandemic to attempt cautious surveillance, including asymptomatic children and velocity of infection dependent on serology, to more readily describe disease in children and their part in transmission frameworks. Regardless of whether diverse control guidelines decelerate, in the end halting the local spread of COVID-19 infection, the effective containment of this infection actually leaves the populace in danger of resurgence because of deficient acquisition of immunity. In the event that herd immunity can be actuated, it would go about as a boundary to stop the spread of disease [45]. Additionally, vaccines, variable susceptibility and exposure to a great extent establish herd immunity.

Immunization of children to instigate herd immunity has demonstrated efficacy in forestalling the spread of numerous infectious diseases, where children have a critical part in transmission. An elevated level of immunity in one age gathering, who assume a function in transmission, can create herd immunity for others [46], and it is obvious that vaccine inoculation of children is more successful than immunization of old individuals, in specific circumstances, as exhibited in vaccination against flu [46,47], pneumococcal infection [48], rotavirus [19] and numerous others. It is imperative that the overall function of various age cohorts in transmission must be considered. The backhanded advantages of COVID-19 vaccination in children may give or make some assured protection to more established, unvaccinated populaces. At the point when children are immunized, it will be simpler to accomplish enough immunity required for general protection in a given populace.

Coronavirus mortality is emphatically age-ward, and Africa has a similarly more youthful populace than other continents [49]. Consequently, children may be a significant objective for mediations pointed toward decreasing transmission in nations with youthful populaces, particularly since access through school vaccination might be more direct than getting to grown-ups. Despite the fact that the need for COVID-19 immunization would legitimately be for that at the most noteworthy danger of infection, for example, medical services labourers, and those at the most elevated danger of severe infection, for instance, older adults, vaccination of children might be another critical cohort for their own safeguard and to help herd immunity. a) Herd immunity constraints in the current covid-19 pandemic Herd immunity might be accomplished when the populace is exposed to the

infection and develops a characteristic immune reaction and somatic defense system to the infection or when the populace is vaccinated against the COVID-19 malady to accomplish immunity along these lines, by vaccinating certain cohorts of the populace, the spread of the infection will go down. Without an immunization, building herd immunity against COVID-19 through natural infection is hypothetically conceivable. Be that as it may, there is no ethical way to arrive at this objective, as the social results of characteristic natural exposure might be overwhelming [42].

Without a doubt, current mathematical and epidemiological examinations propose that herd immunity through common methods may not be the response to stop the novel coronavirus; exposure to the viral infection ought to be maintained until either an immunization or viable pharmacological medicines are accessible. Hence, pharmacological Interventions (PIs, for example, the utilization of hydroxychloroquine, azithromycin, lopinavir, ritonavir, ribavirin, chloroquine phosphate, and arbidol, have been proposed to be conceivably compelling in fighting COVID-19 once the infection's hereditary arrangement and component of infection are unchangeable [50]. Notwithstanding, the adequacy and efficacy of these competitor drugs in the treatment of COVID-19 should be affirmed in further preclinical and clinical preliminaries, in spite of the in vitro examinations and non-clinical preliminaries previously available [51].

9 V. Group Immunity and Vaccination in Covid-19 Transmission Supression

Vaccine development could help halt the spread of the infection, particularly among the most susceptible populaces. This objective has become the procedure most seriously sought after by worldwide research facilities [52,42]. Immunizations have generally been viewed as a type of a preventive mediation for immediate and aberrant protection for herd immunity in an comprehensive populace Reformist vaccination missions and enhancements in complete disinfection in metropolitan settings have assisted with improving herd immunity [53]. Note that vaccination is additionally balanced by vaccine type, the individual life form reaction, anticipation program adherence, and the age of administration [54].

Vaccines train the body to perceive and battle a particular microbe. The viral spread is confined when the immunization rate or the commonness of a high level of positive serological people in the populace truly limit the transmission of the infectious agent starting with one human then onto the next. Building up a "basic" populace immunity rate to control the expansion of COVID-19 is, with current logical information simply theoretical. Moreover, the mass immunization of billions of individuals could be one of the most significant worldwide challenges of the 21st century [15]. Presently, 25 vaccines are being created to battle COVID-19, with subsidizing chiefly originating from private drug foundations. Figure 2 shows the varying vaccine choices presently being investigated. Some research groups are utilizing inactivated viruses, however most investigations on vaccine varieties center around vaccinations dependent on viral proteins and nucleic acids. As recently detailed, COVID-19 is particularly hazardous in individuals of cutting advanced age or with existing underlying conditions, for example, diabetes, coronary issues, malignancy, or a weakened immune system. Other natural elements, for example, diet quality, inactive way of life, and clinical medicines, may likewise restrict the immune reactions of most weak individuals before future vaccine administration against COVID-19. Thusly, herd immunity might be the most significant "present moment" technique to secure this segment of the population (55). Until a safe vaccine is ultimately developed, research on definite novel treatments (or an efficient blend of existing treatments), together with action plans to contain the spread of the virus, seem to be the only substitute for protecting atrisk populations [56].

Discovering an efficient vaccine will not be without firm challenges to surmount, such as its resultant effects, price and ease of accessibility, limited secondary effects to vulnerable people, long-term immune response, and the keenness of the population to be vaccinated voluntarily [27]. There are certain limits to swaying people to engage in mass vaccination [57], even when direct immunization could safely extend indirect immunity to the most vulnerable populations [58,59]. However, public communication plans on the importance of herd immunity and easily accessible vaccination campaigns will be necessary to increase observance to prevention programs in the fight against the COVID-19 infection [60].


10 VI.

11 Conclusion

Among humans, the novel COVID-19 spread happens by direct person-to-person contact via coughing, sneezing, and inhalation of droplets/infected aerosols and contact spread with the oral and nasal membranes. Herd immunity has expansively been used for the containment of multiple diseases and presents the basis for vaccines and their relevancies. The communicable pathogen has been discovered, but the transitional host is still undecided. Additionally, there should be long-lasting immunity, which is still feasible for COVID-19. With no vaccine in view, cluster immunity is likely to be attained when about 70% of the people has been infected. Diverse forms of vaccines are in advance stages of formulation worldwide in order to curb the scourge of the virus. The COVID-19 pandemic could only end if a clinically-safe, tested and effective vaccine is confirmed, with the concept of herd immunization acquired and embraced. Finally, pending the availability of a vaccine, epidemic

220 spread suppression via exceptionally-concentrated health criterions potentially shows to be the workable and
221 secure plan, implementing quarantine and the application of numerous contact with wherewithal management to
222 suppress the spread of the virus.¹

¹© 2021 Global JournalsProspective Promising Signs of Herd Immunity in COVID-19 Transmission Suppression Via Vaccination



1

Figure 1: Figure 1 :




Figure 2: Figure 2 :

1

Viral vaccines	Age range (yrs) of vaccine coverage	Sample pop-ulation	Vaccination recip-ients	Percentage immuned tion of	herd reduc-	Location/year
Seven-valent pneumococcal conjugate vaccine (PCV7)	<5 (vaccinated) >65(not vaccinated)	851 1812	2,105	98% reduction 85% reduction		UK/2006
Measles conjugate vaccine (MCV)	18 months (neonates)	3,115	2,105	85% reduction 100% reduction		Australia/2008
Live attenuated influenza vaccine (LAIV)	10-13 (congenital) 4-12 86% vaccinated 14% unvaccinated	12,855 15,721	12,853 9021	60% reduction		Canada/2008
Rotavirus vaccine (Rotarix)	0-4 5-14 15-24	66,000	64,000	15% reduction		United States/2008
II.	Elements of Herd Immunity in a Population with Covid-19					

Figure 3: Table 1 :

[Smith ()] , D R Smith . *Herd Immunity. Vet. Clin. Pract* 2019. 35 p. .

[Graham et al. ()] ‘A decade after SARS: Strategies for controlling emerging coronaviruses’. R L Graham , E F Donaldson , R S Baric . *Nat. Rev. Microbiol* 2013. 11 p. .

[Singhal ()] ‘A review of coronavirus disease-2019 (COVID-19)’. T Singhal . *Indian J. Pediatr* 2020. 87.

[Biswas et al. ()] ‘A SEIR model for control of infectious diseases with constraints’. Mha Biswas , L T Paiva , De Pinho , Mdr . *Math. Biosci. Eng* 2014. 11 p. .

[Cortegiani et al. ()] ‘A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19’. A Cortegiani , G Ingoglia , M Ippolito , A Giarratano , S Einav . *J. Crit. Care* 2020. 57 p. .

[Zhao et al. (2020)] ‘Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019’. J Zhao , Q Yuan , H Wang . *Clin Infect Dis* March 28, 2020. (Published online)

[Kinoshita and Nishiura ()] ‘Assessing herd immunity against rubella in Japan: A retrospective seroepidemiological analysis of age-dependent transmission dynamics’. R Kinoshita , H Nishiura . *BMJ Open* 2016. 6.

[Albuquerque et al. ()] ‘Chikungunya virus infection: Report of the first case diagnosed in Rio de Janeiro’. Igcd Albuquerque , R Marandino , A P Mendonça , Rmr Nogueira , Pfdc Vasconcelos , L R Guerra . *Brazil. Rev. Soc. Bras. Med. Trop* 2012. 45 p. .

[Zhen-Dong et al. (2020)] ‘Clinical and transmission dynamics characteristics of 406 children with coronavirus disease 2019 in China: a review’. Y Zhen-Dong , Z Gao-Jun , J Run-Ming , L Zhi-Sheng , D Zong-Qi , X Xiong . *J Infect* 2020. April.

[Khandaker et al. ()] ‘Congenital and neonatal varicella: impact of the national varicella vaccination programme in Australia’. G Khandaker , H Marshall , E Peadon . *Arch Dis Child* 2011. 96 p. .

[Plotkin ()] ‘Correlates of vaccine-induced immunity’. S A Plotkin . *Clin. Infect. Dis* 2008. 47 p. .

[Nicola et al. ()] ‘COVID-19 from veterinary medicine and one health perspectives: What animal coronaviruses have taught us’. D Nicola , M Vito , J S Linda , B Canio . *Res. Vet. Sci* 2020. 131 p. .

[Stebbing et al. ()] ‘COVID-19: Combining antiviral and anti-inflammatory treatments’. J Stebbing , A Phelan , I Grin , C Tucker , O Oechsle , D Smith . *Lancet Infect. Dis* 2020. 20 p. .

[Loeb et al. ()] ‘Effect of influenza vaccination of children on infection rates in Hutterite communities: a randomized trial’. M Loeb , M L Russell , L Moss . *JAMA* 2010. 303 p. .

[Adegbola et al. ()] ‘Elimination of Haemophilus influenzae type b (Hib) disease from the Gambia after introduction of a Hib conjugate vaccine: A prospective study’. R Adegbola , O Secka , G Lahai , N Lloyd-Evans , A Njie , S Usen . *Lancet* 2005. 366 p. .

[Chen et al. ()] ‘Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: A descriptive study’. N Chen , M Zhou , X Dong , J Qu , F Gong , Y Han . *Lancet* 2020. 395 p. .

[Bi et al. (2020)] ‘Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study’. Q Bi , Y Wu , S Mei , C Ye , X Zou , Z Zhang . *Lancet Infect Dis* 2020. April.

[Dong et al. ()] ‘Epidemiology of COVID-19 among children in China’. Y Dong , X Mo , Y Hu , X Qi , F Jiang , Z Jiang . *Pediatrics* 2020. p. 145.

[Mizumoto et al. ()] ‘Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship’. K Mizumoto , K Kagaya , A Zarebski , G Chowell . *Euro Surveill* 2020. 2000180. 25 p. 2020.

[Moulton et al. ()] ‘Estimation of the indirect effect of Haemophilus influenzae type b conjugate vaccine in an American Indian population’. L H Moulton , S Chung , J Croll , R Reid , R C Weatherholtz , M Santosham . *Int. J. Epidemiol* 2000. 29 p. .

[Del Giudice et al. ()] ‘Fighting against a protean enemy: Immunosenescence, vaccines, and healthy aging’. G Del Giudice , J J Goronzy , B Grubeck-Loebenstien , P H Lambert , T Mrkvan , J J Stoddard . *NPJ Aging Mech. Dis* 2017. 4 p. 1.

[John and Samuel ()] ‘Herd immunity and herd effect: New insights and definitions’. T J John , R Samuel . *Eur. J. Epidemiol* 2000. 16 p. .

[Miller et al. ()] ‘Herd immunity and serotype replacement 4 years after seven-valent pneumococcal conjugate vaccination in England and Wales: an observational cohort study’. E Miller , N J Andrews , P A Waight , M Slack , R George . *Lancet Infect Dis* 2011. 11 p. .

[Ali et al. ()] ‘Herd immunity conferred by killed oral cholera vaccines in Bangladesh: A reanalysis’. M Ali , M Emch , Von Seidlein , L Yunus , M Sack , D A Rao , M . *Lancet* 2005. 366 p. .

[Fine et al. ()] ‘Herd immunity”: A rough guide’. P Fine , K Eames , D L Heymann . *Clin. Infect. Dis* 2011. 52 p. .

- [Kwok et al. ()] ‘Herd immunity-estimating the level required to halt the COVID-19 epidemics in affected countries’. K O Kwok , F Lai , W I Wei , Sys Wong , J W Tang . *J. Infect* 2020. 80 p. .
- [Fox et al. ()] ‘Herd immunity: Basic concept and relevance to public health immunization practices’. J P Fox , L Elveback , W Scott , L Gatewood , E Ackerman . *Am. J. Epidemiol* 1971. 94 p. .
- [Fine ()] ‘Herd immunity: History, theory, practice’. P E Fine . *Epidemiol. Rev* 1993. 15 p. .
- [Goncalves ()] ‘Herd immunity: Recent uses in vaccine assessment’. G Goncalves . *Expert Rev. Vaccines* 2008. 7 p. .
- [Randolph and Barreiro ()] *Herd immunity: Understanding COVID-19*, H Randolph , L B Barreiro . 2020. Cell Press. p. .
- [Gautret et al. ()] ‘Hydroxychloroquine and azithromycin as a treatment of COVID-19: Results of an open-label non-randomized clinical trial’. P Gautret , J C Lagier , P Parola , V T Hoang , L Meddeb , M Mailhe . *Int. J. Antimicrob. Agents* 2020. p. 105949.
- [Hochberg ()] *Importance of suppression and mitigation measures in managing COVID-19 outbreaks*. medRxiv, M E Hochberg . 2020.
- [Nymark et al. ()] ‘Inclusion of the value of herd immunity in economic evaluations of vaccines. A systematic review of methods used’. L S Nymark , T Sharma , A Miller , U Enemark , U K Griffiths . *Vaccine* 2017. 35 p. .
- [Gomes et al. (2020)] ‘Individual variation in susceptibility or exposure to SARS-CoV-2 lowers the herd immunity threshold’. M G Gomes , R M Corder , J G King , K E Langwig , C Souto-Maior , J Carneiro . *medRxiv* 2020. January. p. 2020.
- [Fresnadillo-Martínez et al. ()] ‘Mathematical modelling of the propagation of infectious diseases: Where we came from, and where we are going’. M J Fresnadillo-Martínez , E Garcia-Sanchez , E Garcia-Merino , J E García-Sánchez . *Rev. Esp. Quím* 2013. 26 p. .
- [Xun et al. ()] *Neutralizing antibody responses to SARS-CoV-2 in a COVID-19 recovered 2 patient cohort and their implications*, J Xun , L Lu , S Jiang , H Lu , Y Wen , J Huang . Medrxiv 2020.
- [Zhou and Zhao ()] ‘Perspectives on therapeutic neutralizing antibodies against the Novel Coronavirus SARS-CoV-2’. G Zhou , Q Zhao . *Int. J. Biol. Sci* 2020. 16 p. 1718.
- [Pittet and Posfay-Barbe ()] ‘Pneumococcal vaccines for children: a global public health priority’. L F Pittet , K M Posfay-Barbe . *Clin Microbiol Infect* 2012. 18 (5) p. .
- [Lopman et al. ()] ‘Post-licensure experience with rotavirus vaccination in high and middle income countries’. B A Lopman , D C Payne , J E Tate , M M Patel , M M Cortese , U D Parashar . *Curr Opin Virol* 2006 to 2011. 2012. 2 (4) p. .
- [Ahmed et al.] *Preliminary identification of potential vaccine targets for the COVID-19 coronavirus (SARS-CoV-2) based on*, S F Ahmed , A A Quadeer , M R Mckay .
- [Wu et al. ()] ‘Prolonged presence of SARS-CoV-2 viral RNA in faecal samples’. Y Wu , C Guo , L Tang , Z Hong , J Zhou , X Dong . *Lancet Gastroenterol. Hepatol* 2020. 5 p. .
- [Reinfection Becoming an Issue in China, Strategist Says (2020)] *Reinfection Becoming an Issue in China, Strategist Says*, COVID-19. <https://www.cnbc.com/video/2020/03/16/covid-19-reinfection-becoming-an-issue-in-china-strategist-says.html> April 2020. (Available online)
- [SARS-CoV immunological studies Viruses ()] ‘SARS-CoV immunological studies’. *Viruses* 2020. 12 p. 254.
- [Kim ()] ‘Seasonal influenza and vaccine herd effect’. T H Kim . *Clin Exp Vaccine Res* 2014. 3 (2) p. .
- [Liu et al. ()] ‘Secondary attack rate and superspreading events for SARS-CoV-2’. Y Liu , R M Eggo , A J Kucharski . *Lancet* 2020. 395 p. e47.
- [Lloyd-Smith et al. ()] ‘Super-spreading and the effect of individual variation on disease emergence’. J O Lloyd-Smith , S J Schreiber , P E Kopp , W M Getz . *Nature* 2005. 438 p. .
- [To et al. ()] ‘Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study’. K K To , O T Tsang , W S Leung . *Lancet Infect Dis* 2020. 20 (5) p. .
- [Casadevall and Pirofski ()] ‘The convalescent sera option for containing COVID-19’. A Casadevall , L A Pirofski . *J. Clin. Investig* 2020. 130 p. .
- [Mougeni et al. (2020)] ‘The potential effect of the African population age structure on COVID-19 mortality’. F Mougeni , A Mangaboula , B Lell . *medRxiv* 2020. January. p. 2020.
- [Topley and Wilson ()] ‘The spread of bacterial infection. The problem of herd-immunity’. Wwc Topley , G S Wilson . *Epidemiol. Infect* 1923. 21 p. .

[Zhang et al. ()] ‘The use of anti-inflammatory drugs in the treatment of people with severe coronavirus disease 2019 (COVID-19): The experience of clinical immunologists from China’. W Zhang , Y Zhao , F Zhang , Q Wang , T Li , Z Liu . *Clin. Immunol* 2020. 214 p. 108393.

[Peng et al. ()] ‘Transmission routes of 2019-nCoV and controls in dental practice’. X Peng , X Xu , Y Li , L Cheng , X Zhou , B Ren . *Int. J. Oral Sci* 2020. 12 p. 9.

[Metcalf et al. ()] ‘Understanding herd immunity’. C J Metcalf , M Ferrari , A L Graham , B T Grenfell . *Trends Immunol* 2015. 36 p. .

[Korppi ()] ‘Universal pneumococcal vaccination provides marked indirect beneficial effects through herd immunity’. M Korppi . *Acta Paediatr* 2018. 107 p. .

[Ali et al. ()] ‘Unmasking herd protection by an oral cholera vaccine in a cluster-randomized trial’. M Ali , F Qadri , D R Kim , T Islam , J Im , F Ahmmed . *Int. J. Epidemiol* 2019. 48 p. .

[Herrmann and Schwartz] *Using network science to propose strategies for effectively dealing with pandemics: The COVID-19 example*, H A Herrmann , J M Schwartz . medRxiv. 2020.

[Rashid et al. ()] ‘Vaccination and herd immunity: What more do we know?’. H Rashid , G Khandaker , R Booy . *Curr. Opin. Infect Dis* 2012. 25 p. .

[Kim et al. (2011)] ‘Vaccine herd effect’. T H Kim , J Johnstone , M Loeb . *Scand J Infect Dis* 2011. September. 43 (9) p. .

[Colgrove ()] ‘Vaccine refusal revisited the limits of public health persuasion and coercion’. J Colgrove . *N. Eng. J. Med* 2016. 375 p. .

[Wölfel et al. (2020)] ‘Virological assessment of hospitalized patients with COVID-2019’. R Wölfel , V M Corman , W Guggemos . *Nature*. Published online April 1, 2020.

[World Health Organization (WHO) guide for standardization of economic evaluations of immunization programmes ()] *World Health Organization (WHO) guide for standardization of economic evaluations of immunization programmes*, 2008. (WHO/1VB/08.14. This publication is available on the internet at: www.who.int/vaccines-documents)